

Generic Environmental Impact Statement for In-Situ Leach Uranium Milling Facilities

Chapters 5 through 12 and Appendices A through G

Final Report

Office of Federal and State Materials and Environmental Management Programs

Wyoming Department of Environmental Quality Land Quality Division



Generic Environmental Impact Statement for In-Situ Leach Uranium Milling Facilities

Chapters 5 through 12 and Appendices A through G

Final Report

Manuscript Completed: May 2009

Date Published: May 2009

Prepared by:

U.S. Nuclear Regulatory Commission Office of Federal and State Materials and Environmental Management Programs

Wyoming Department of Environmental Quality Land Quality Division

ABSTRACT

The Atomic Energy Act of 1954 and the Uranium Mill Tailings Radiation Control Act of 1978 authorize the U.S. Nuclear Regulatory Commission (NRC) to issue licenses for the possession and use of source material and byproduct material. These statutes require NRC to license facilities that meet NRC regulatory requirements that were developed to protect public health and safety from radiological hazards. *In-situ* leach (ISL) uranium recovery facilities must meet NRC regulatory requirements in order to obtain a source material license to operate.

Under NRC's environmental protection regulations in the Code of Federal Regulations, Title 10, Part 51, which implement the National Environmental Policy Act (NEPA), issuance of a license to possess and use source material for uranium milling requires an environmental impact statement (EIS) or a supplement to an EIS. NRC has prepared a generic environmental impact statement (GEIS) to help fulfill this requirement. The GEIS assesses the potential environmental impacts associated with the construction, operation, aquifer restoration, and decommissioning of an ISL uranium recovery facility in four specified regions in the western United States. The intent of the GEIS is to determine which impacts would be essentially the same for all ISL facilities and which ones would result in varying levels of impacts for different facilities, thus requiring further site-specific information to determine the potential impacts. As such, the GEIS provides a starting point for NRC's NEPA analyses for site-specific license applications for new ISL facilities, as well as for applications to amend or renew existing ISL licenses.

NRC developed this GEIS using (1) knowledge gained during the past 30 years licensing and regulating ISL facilities, (2) the active participation of the State of Wyoming Department of Environmental Quality as a cooperating agency, and (3) public comments received during the preparation of the GEIS. NRC's licensing experience indicates that the technology used for ISL uranium recovery is relatively standardized throughout the industry and therefore appropriate for a programmatic evaluation in a GEIS.

Based on discussions between uranium recovery companies and the NRC staff, future ISL facilities could be located in portions of Wyoming, Nebraska, South Dakota, and New Mexico. NRC is the licensing authority for ISL facilities in these states.

Paperwork Reduction Act Statement

This NUREG contains information collection requirements that are subject to the Paperwork Reduction Act of 1995 (44 U.S.C. 3501 et seq.) These information collections were approved by the Office of Management and Budget, approval numbers 3150-0020; 3150-0014.

Public Protection Notification

The NRC may not conduct or sponsor, and a person is not required to respond to, a request for information or an information collection requirement unless the requesting documents displays a currently valid OMB control number.

CONTENTS

Section					Page
ARSTR	ΔСТ				iii
				SION FACTORS	
SI (IVIO	DEKN	IVIETRI	C) CONVER	510N FACTORS	
1 I	INTRO	DUCTIO	N		1-1
•	1.1	Rationa	le of the GE	IS	1-1
•	1.2	The Pro	posed Fede	ral Action	1-4
				for the Proposed Federal Action	
				Used in the GEIS	
		1.4.1		S	
		1.4.2		ogy	
		1.4.3		of the GEIS	
			1.4.3.1	Describing the ISL Process	
			1.4.3.2	Describing the Affected Environment	
			1.4.3.3	Identifying Environmental Issues and	,
			1.4.5.5	Characterizing Significance	1_7
	1.5	Scope	of the CEIS	Characterizing digrillicance	1-7 1-8
	1.5	1.5.1		Scoping Process	
		1.5.1		Idied in Detail	
		1.5.2		minated From Detailed Study	
		1.5.3			
	4.0	_		tside of the Scope of the GEIS	
	1.6	•		n Uranium ISL Facility Licensing	
		1.6.1		gencies	
			1.6.1.1	NRC	
			1.6.1.2	EPA	
			1.6.1.3	Occupational Safety and Health Administration	
			1.6.1.4	U.S. Department of Transportation	1-16
			1.6.1.5	U.S. Department of Interior, U.S. Bureau of	
				Land Management	
			1.6.1.6	Other Federal Agencies	
		1.6.2	Tribal Age		
		1.6.3	•	ncies	
			1.6.3.1	Wyoming Department of Environmental Quality	
			1.6.3.2	Nebraska Department of Environmental Quality	1-17
			1.6.3.3	South Dakota Department of Environmental and	
				Natural Resources	1-18
			1.6.3.4	New Mexico Environmental Department	1-18
•	1.7	Licensir	ng and Perm	itting Process for a Uranium ISL Facility	1-18
		1.7.1	The NRC	Licensing Process	1-18
		1.7.2	EPA Perm	nitting	1-21
			1.7.2.1	Water Resources	
			1.7.2.2	Air Quality	
		1.7.3		leral Agencies	
		1.7.4		encies	
		1.7.5	State Age	ncies	1-24

Section	n				Page
			1.7.5.1	Wyoming	1-25
			1.7.5.2	Nebraska	
			1.7.5.3	South Dakota	
			1.7.5.4	New Mexico	
	1.8	Use of th	ne GEIS in th	ne NRC Licensing Process	
		1.8.1		or Licensee Environmental Report	
		1.8.2		e Review of the License Application and	
				ntal Report	1-27
		1.8.3		-Specific Environmental Review	
		1.8.4		icipation Activities	
		1.8.5		Final Environmental Review Document and Findings.	
	1.9	Reference			
2	IN-SIT	U LEACH	I URANIUM	RECOVERY AND ALTERNATIVES	2-1
	2.1	Overview	v of ISL Urar	nium Recovery	2-1
		2.1.1		stry of Uranium	
		2.1.2		haracteristics of Uranium Deposits	
		2.1.3		escription of ISL Facilities	
	2.2	Precons			
	2.3				
		2.3.1	Undergrou	nd Infrastructure	2-8
			2.3.1.1	Well Fields	
			2.3.1.2	Pipelines	2-14
		2.3.2	Surface Fa	icilities	
	2.4	Operatio	ns		2-15
		2.4.1		lobilization	
			2.4.1.1	Lixiviant Chemistry	2-16
			2.4.1.2	Lixiviant Injection and Production	
			2.4.1.3	Excursions	
			2.4.1.4	Excursion Monitoring	2-19
		2.4.2	Uranium P	rocessing	
			2.4.2.1	Ion Exchange	2-20
			2.4.2.2	Elution	2-23
			2.4.2.3	Precipitation, Drying, and Packaging	2-23
		2.4.3	Manageme	ent of Production Bleed and Other Liquid Effluents	
	2.5	Aquifer F	Restoration	·	2-26
		2.5.1	Groundwat	ter Transfer	2-27
		2.5.2		ter Sweep	
		2.5.3	Reverse O	smosis, Permeate Injection, and Recirculation	2-29
		2.5.4		n	
	2.6	Deconta	mination, De	ecommissioning, and Reclamation	2-30
	2.7			Management	
		2.7.1	Gaseous o	r Airborne Particulate Emissions	2-33
		2.7.2	Liquid Was	stes	2-36
		2.7.3	•	es	
	2.8	Transpo	rtation		2-38
	2.9	•		and Safety	
	2.10				

Section	on					Page
	2.11	Informa	tion From H	istorical Oper	ation of ISL Uranium Milling Facilities .	2-42
		2.11.1			illing Facilities	
		2.11.2				
		2.11.3				
		2.11.4				
		2.11.5				
		2.11.6			ation	
	2.12	Alternat			uded in the Impact Analysis	
		2.12.1			ive	
	2.13	Alternat			luded From the Impact Analysis	
	2.14					
3	DESC	'DIDTION	I OE THE AI	FECTED EN	IVIRONMENT	3 1-1
3	3.1				TVITO INVICTOR	
	3.1	3.1.1			efining Uranium Milling Regions	
		3.1.1				
		3.1.2	3.1.2.1		 All Uranium Milling Regions and Characteristics of Sandstone-Hos 	
			3.1.2.1			
			2400		Ore Deposits	
			3.1.2.2	Complex L	and Ownership Rights, Responsibilitie	35,
		242	Deference		rtunities	
	2.0	3.1.3				
3.2				Region		
		3.2.1				
		3.2.2 3.2.3				
		3.2.4				
			3.2.4.1		aters	
			3.2.4.2		and Waters of the United States	
			3.2.4.3		ter	
				3.2.4.3.1	Regional Aquifer Systems	3.2-17
				3.2.4.3.2	Aquifer Systems in the Vicinity of	
					Uranium Milling Sites	
				3.2.4.3.3	Uranium-Bearing Aquifers	
				3.2.4.3.4	Other Important Surrounding Aquife	
					for Water Supply	
		3.2.5	• • • • • • • • • • • • • • • • • • • •			
			3.2.5.1			
			3.2.5.2			
			3.2.5.3		d and Endangered Species	
		3.2.6	Meteorolo	gy, Climatolo	gy, and Air Quality	3.2-44
			3.2.6.1	Meteorolog	gy and Climatology	3.2-44
			3.2.6.2	Air Quality		3.2-46
		3.2.7	Noise			3.2-50
		3.2.8	Historical	and Cultural	Resources	3.2-53
			3.2.8.1	Cultural Re	esources Overview	3.2-53
			3.2.8.2	National R	egister of Historic Properties	
					Registers	3.2-58
			3.2.8.3		sultation	

Section					Page
		3.2.8.4	Places of 0	Cultural Significance	3.2-61
	3.2.9	Visual/Scer	nic Resource	es	3.2-63
	3.2.10				
		3.2.10.1		hics	
		3.2.10.2			
		3.2.10.3			
		3.2.10.4		nt Structure	
			3.2.10.4.1		
					3.2-76
				3.2.10.4.1.2 Wyoming	
		00405	3.2.10.4.2	•	
		3.2.10.5		nce	
		3.2.10.6			
	0 0 4 4	3.2.10.7		Social Services	
	3.2.11			al Health	
		3.2.11.1		d Radiological Conditions .	
		3.2.11.2		Ith and Safety	
	0.0.40	3.2.11.3		nal Health and Safety	
0.0	3.2.12				
3.3	•			egion	
	3.3.1				
	3.3.2				
	3.3.3				
	3.3.4				
		3.3.4.1		aters and Waters of the United St	
		3.3.4.2 3.3.4.3			
		3.3.4.3	3.3.4.3.1	ter	
			3.3.4.3.1	Regional Aquifer System	
			3.3.4.3.2	Aquifer Systems in the Vi of Uranium Milling Sites.	•
			3.3.4.3.3	Uranium-Bearing Aquifer	
			3.3.4.3.4	Other Important Surround	
			5.5.4.5.4	Aquifers for Water Supply	
	3.3.5	Ecology		Aquilers for water Supply	
	5.5.5	3.3.5.1			
		3.3.5.2			
		3.3.5.3		d and Endangered Species	
	3.3.6			gy, and Air Quality	
	0.0.0	3.3.6.1		gy and Climatology	
		3.3.6.2			
	3.3.7		,		
	3.3.8			Resources	
	0.0.0	3.3.8.1		esources Overview	
		3.3.8.2		egister of Historic Propertie	
		0.0.0.2		Registers	
		3.3.8.3		sultation	
		3.3.8.4		Cultural Significance	
	3.3.9			98	
	-				

Section						Page
	3.3.10	Socioecono				
		3.3.10.1				
		3.3.10.2				
		3.3.10.3				
		3.3.10.4	• •			
					Colorado	
				3.3.10.4.1.2		
					Wyoming	
				•		
		3.3.10.5				
		3.3.10.6				
		3.3.10.7				
	3.3.11	Public and				
		3.3.11.1	Background F	Radiological C	onditions	3.3-60
		3.3.11.2				
		3.3.11.3			afety	
	3.3.12	References				3.3-61
3.4	Nebrasi				g	
	3.4.1	Land Use				3.4-1
	3.4.2	Transportat	ion			3.4-4
	3.4.3	Geology an	d Soils			3.4-6
		3.4.3.1	The Black Hill			
			Dakota-North	eastern Wyor	ming)	3.4-8
		3.4.3.2	The Crawford	l Basin (North	western Nebraska)	3.4-13
	3.4.4	Water Reso	ources	······	···········	3.4-17
		3.4.4.1	Surface Wate	rs		3.4-17
		3.4.4.2	Wetlands and	Waters of the	e United States	3.4-22
		3.4.4.3	Groundwater			3.4-23
			3.4.4.3.1 F	Regional Aqui	fer Systems	3.4-23
					ns in the Vicinity of	
					g Sites	3.4-26
					ing Aquifers	
		3.4.4.3.4			ng Aquifers for	
	3.4.5	Ecology				
		3.4.5.1				
		3.4.5.2				
		3.4.5.3			ed Species	
	3.4.6				ity	
	0	3.4.6.1			gy	
		3.4.6.2				
	3.4.7		•			
	3.4.8				estern South Dakota	0. 1 00
	5					3.4-57
		3.4.8.1			ew	
		JJ				

Section						Page
			3.4.8.1.1	Cultural Reso	urces of Western	
				and Southwes	stern South Dakota	3.4-57
			3.4.8.1.2	Cultural Reso	urces of	
				Western Nebr	aska	3.4-60
		3.4.8.2	National Re	egister of Histori	ic Properties	
			and State F	Registers		3.4-62
			3.4.8.2.1		erties in Western	
						3.4-62
			3.4.8.2.2	Historic Prope		
				Western Nebr	aska	3.4-67
		3.4.8.3	Tribal Cons	ultations		3.4-68
			3.4.8.3.1	South Dakota	Tribal Consultation	3.4-68
			3.4.8.3.2	Nebraska Trib	al Consultation	3.4-69
		3.4.8.4	Places of C		ance	
			3.4.8.4.1		ural Significance	
					ern South Dakota	3.4-70
			3.4.8.4.2		ural Significance	
					ebraska	3.4-70
	3.4.9	Visual/Scer	nic Resource			
	3.4.10					
		3.4.10.1				
		3.4.10.2				
		3.4.10.3				
		3.4.10.4				
					Nebraska	
				3.4.10.4.1.2		
				3.4.10.4.1.3		
			341042			
			01.11.01.112	3.4.10.4.2.1	Nebraska	
				3.4.10.4.2.2	South Dakota	
				3.4.10.4.2.3	Wyoming	
			3.4.10.4.3		an Communities	
		3.4.10.5				
		0	3.4.10.5.1			
			3.4.10.5.2			
			3.4.10.5.3			
			3.4.10.5.4		an Communities	
		3.4.10.6				
		0.1.10.0	3.4.10.6.1			
			3.4.10.6.2			
			3.4.10.6.3			
		3.4.10.7			3	
	3.4.11					
	J. T . 1 1	3.4.11.1			Conditions	
		3.4.11.2				
		3.4.11.3	Occupation	al Health and S	afety	3.∓-34 ∕ 0_1 ∕
	3.4.12				aicty	

Section					Page				
3.5		Northwestern New Mexico Uranium Milling Region							
	3.5.1 Land Use								
	3.5.2								
	3.5.3								
	3.5.4								
		3.5.4.1		aters					
		3.5.4.2		and Waters of the United States					
		3.5.4.3		ter					
			3.5.4.3.1	Regional Aquifer Systems	3.5-18				
			3.5.4.3.2	Aquifer Systems in the Vicinity of					
				Uranium Milling Sites					
			3.5.4.3.3	Uranium-Bearing Aquifers	3.5-20				
			3.5.4.3.4	Other Important Surrounding					
				Aquifers for Water Supply					
	3.5.5	Ecology							
		3.5.5.1							
		3.5.5.2							
		3.5.5.3		d and Endangered Species					
	3.5.6			ogy, and Air Quality					
		3.5.6.1		gy and Climatology					
		3.5.6.2							
	3.5.7	Noise							
	3.5.8			Resources					
		3.5.8.1		co Historic and Cultural Resources	3.5-40				
		3.5.8.2		egister of Historic Properties					
				Registers					
		3.5.8.3		co Tribal Consultation					
		3.5.8.4		Cultural Landscapes					
	3.5.9			es					
	3.5.10								
		3.5.10.1		hics					
		3.5.10.2							
		3.5.10.3							
		3.5.10.4		ent Structure					
			3.5.10.4.1						
				3.5.10.4.1.1 Arizona					
				3.5.10.4.1.2 New Mexico					
			3.5.10.4.2	County Data	3.5-72				
		3.5.10.5	Local Fina	nce	3.5-72				
		3.5.10.6							
		3.5.10.7	Health and	d Social Services	3.5-77				
	3.5.11	Public and	l Occupation	al Health	3.5-77				
		3.5.11.1	Backgroun	nd Radiological Conditions	3.5-77				
		3.5.11.2	Public Hea	alth and Safety	3.5-78				
		3.5.11.3	Occupation	nal Health and Safety	3.5-78				
	3.5.12	Reference	s		3.5-78				

Section	n						Page
4	ENVIF	RONMEN	TAL IMPAC	rs of cons	STRUCTION,	OPERATION, AQUIFER	
	REST	ORATION	I, AND DEC	OMMISSION	IING ACTIVIT	ΓΙΕS	4.1-1
	4.1	Introduc	tion				4.1-1
	4.2						
		4.2.1					
			4.2.1.1			Land Use	
			4.2.1.2			nd Use	
			4.2.1.3			acts to Land Use	
			4.2.1.4			cts to Land Use	
		4.2.2	Transporta				
			4.2.2.1			Transportation	
			4.2.2.2			ansportation	
			4.2.2.3			acts to Transportation	
			4.2.2.4	•	•	cts to Transportation	
		4.2.3					
		7.2.0	4.2.3.1			Geology and Soils	
			4.2.3.1			eology and Soils	
			4.2.3.3			acts to Geology	4.2-10
			4.2.3.3			acis io Geology	1 2 12
			4.2.3.4				
		4.2.4			• .	cts to Geology and Soils	
		4.2.4					
			4.2.4.1				4.2-15
				4.2.4.1.1		n Impacts to	4045
				10110		ater	4.2-15
				4.2.4.1.2			4040
				40440		ater	4.2-16
				4.2.4.1.3		storation Impacts to	
						ater	4.2-17
				4.2.4.1.4	Decommiss	sioning Impacts to	
				_		ater	
			4.2.4.2		er Impacts		4.2-19
				4.2.4.2.1	Construction	n Impacts	
					to Groundw	/ater	4.2-19
				4.2.4.2.2	Operation I	mpacts to Groundwater	4.2-19
					4.2.4.2.2.1	Operation Impacts to	
						Shallow (Near-Surface)	
						Aquifers	4.2-20
					4.2.4.2.2.2	Operation Impacts to	
						Production and	
						Surrounding Aquifers	4.2-21
					4.2.4.2.2.3	Operation Impacts to	
						Deep Aquifers Below the	
						Production Aquifers	
				4.2.4.2.3	Aguifer Res	storation Impacts to	
						er	4.2-26
				4.2.4.2.4		sioning Impacts to	
					Groundwat	.	4 2-28

Section				Page
	4.2.5	•	Resources Impacts	
		4.2.5.1	Construction Impacts to Ecological Resources	
		4.2.5.2	Operation Impacts to Ecological Resources	4.2-33
		4.2.5.3	Aquifer Restoration Impacts to	
			Ecological Resources	4.2-34
		4.2.5.4	Decommissioning Impacts to	
			Ecological Resources	
	4.2.6		/ Impacts	
		4.2.6.1	Construction Impacts to Air Quality	
		4.2.6.2	Operation Impacts to Air Quality	
		4.2.6.3	Aquifer Restoration Impacts to Air Quality	
		4.2.6.4	Decommissioning Impacts to Air Quality	
	4.2.7	Noise Imp	acts	
		4.2.7.1	Construction Impacts to Noise	4.2-38
		4.2.7.2	Operation Impacts to Noise	4.2-41
		4.2.7.3	Aquifer Restoration Impacts to Noise	4.2-42
		4.2.7.4	Decommissioning Impacts to Noise	4.2-43
	4.2.8	Historical	and Cultural Resources Impacts	4.2-43
		4.2.8.1	Construction Impacts to Historical and Cultural	
			Resources	4.2-44
		4.2.8.2	Operation Impacts to Historical and Cultural	
			Resources	4.2-45
		4.2.8.3	Aquifer Restoration Impacts to Historical and	
			Cultural Resources	4.2-45
		4.2.8.4	Decommissioning Impacts to Historical and Cultu	ıral
			Resources	4.2-45
	4.2.9	Visual/Sce	enic Resources Impacts	4.2-46
		4.2.9.1	Construction Impacts to Visual/	
			Scenic Resources	4.2-46
		4.2.9.2	Operation Impacts to Visual/Scenic Resources	4.2-47
		4.2.9.3	Aquifer Restoration Impacts to Visual/Scenic	
			Resources	4.2-48
		4.2.9.4	Decommissioning Impacts to Visual/Scenic	
			Resources	4.2-48
	4.2.10	Socioecor	nomic Impacts	4.2-49
		4.2.10.1	Construction Impacts to Socioeconomics	
		4.2.10.2	Operation Impacts to Socioeconomics	
		4.2.10.3	Aquifer Restoration Impacts to Socioeconomics	
		4.2.10.4	Decommissioning Impacts to Socioeconomics	
	4.2.11		d Occupational Health and Safety Impacts	
		4.2.11.1	Construction Impacts to Public and Occupational	
			Health and Safety	
		4.2.11.2	Operation Impacts to Public and Occupational He	
			and Safety	
			4.2.11.2.1 Radiological Impacts to Public and	00
			Occupational Health and Safety From	1
			Normal Operations	

Section					Page
			4.2.11.2.	2 Radiological Impacts to Public and Occupational Health and Safety	40.55
			4.2.11.2.	From Accidents Nonradiological Impacts on Public and Occupational Health and Safety From	I
			4.2.11.2.	Normal Operations4 Nonradiological Impacts to Public and	4.2-57
				Occupational Health and Safety From Accidents	1257
		4.2.11.3	•	estoration Impacts to Public and onal Health and Safety	
		4.2.11.4	Decomm	issioning Impacts to Public and onal Health and Safety	
	4.2.12	Waste Ma		mpacts	
				n Impacts to Waste Management	
				Impacts to Waste Management	
			•	storation Impacts to Waste Management	
		4.2.12.4		sioning Impacts to Waste Management	
4.0	4.2.13				
4.3	•			Region	
	4.3.1			Can large and to Large III.	
		4.3.1.1		tion Impacts to Land Use	
		4.3.1.2 4.3.1.3		n Impacts to Land Useestoration Impacts to Land Use	
		4.3.1.3	•	issioning Impacts to Land Use	
	4.3.2			ts	
	4.5.2	4.3.2.1		tion Impacts to Transportation	
		4.3.2.2		n Impacts to Transportation	
		4.3.2.3		estoration Impacts to Transportation	
		4.3.2.4		issioning Impacts to Transportation	
	4.3.3			pacts	
		4.3.3.1		tion Impacts to Geology and Soils	
		4.3.3.2		Impacts to Geology and Soils	
		4.3.3.3		estoration Impacts to Geology and Soils	
		4.3.3.4		issioning Impacts to Geology and Soils	
	4.3.4	Water Re	sources Imp	eacts	4.3-9
		4.3.4.1	Surface V	Vater Impacts	4.3-9
			4.3.4.1.1	Construction Impacts to	
				Surface Water	
			4.3.4.1.2	Operation Impacts to Surface Water	4.3-10
			4.3.4.1.3		
				Surface Water	4.3-10
			4.3.4.1.4	Decommissioning Impacts to	
			_	Surface Water	
		4.3.4.2		er Impacts	
				Construction Impacts to Groundwater	
			4.3.4.2.2	Operation Impacts to Groundwater	4.3-11

Section						Page
				4.3.4.2.2.1	Operation Impacts to Shallow (Near-Surface)	
				4.3.4.2.2.2	Aquifers Operation Impacts to Production and	4.3-11
				4.3.4.2.2.3	Surrounding Aquifers Operation Impacts to Deep	
					Aquifers Below the Production Aquifers	4.3-16
			4.3.4.2.3	•	storation Impacts to er	<i>1</i> 3 ₋ 17
			4.3.4.2.4	Decommiss	sioning Impacts	
	4.3.5	Ecologic	al Bacquree		ater	
	4.3.3	4.3.5.1				
		4.3.5.1			o Ecological Resources	
		4.3.5.2		estoration Im	Ecological Resources	4.3-21
						4.3-21
		4.3.5.4	•	ssioning Impa		
						4.3-22
	4.3.6	Air Qualit				
		4.3.6.1			o Air Quality	
		4.3.6.2			ir Quality	
		4.3.6.3			pacts to Air Quality	
		4.3.6.4	•		acts to Air Quality	
	4.3.7	Noise Im		• •	······	
		4.3.7.1			o Noise	
		4.3.7.2			loise	
		4.3.7.3			pacts to Noise	
		4.3.7.4			acts to Noise	
	4.3.8	Historica	and Cultur	al Resources	s Impacts	
		4.3.8.1			o Historical and	
			Cultural R	esources		4.3-26
		4.3.8.2	•	•	listorical and	
						4.3-26
		4.3.8.3			pacts to Historical and	
		4.3.8.4			acts to Historical and Cultura	
	4.3.9				S	
		4.3.9.1			o Visual/Scenic Resources	
		4.3.9.2	•	•	isual/Scenic Resources	4.3-28
		4.3.9.3			pacts to Visual/Scenic	
						4.3-28
		4.3.9.4			acts to Visual/	
	4.3.10					
		4.3.10.1	Construct	ion Impacts to	o Socioeconomics	4.3-29

Section					Page
				npacts to Socioeconomics	
				toration Impacts to Socioeconomics	
				ioning Impacts to Socioeconomics	
	4.3.11			al Health and Safety Impacts	4.3-33
		4.3.11.1		n Impacts to Public and Occupational	
			Health and	Safety	4.3-33
		4.3.11.2		npacts to Public and Occupational	
				Safety	4.3-33
			4.3.11.2.1	J 1	
				Occupational Health and Safety Fron	
				Normal Operations	4.3-33
			4.3.11.2.2	U 1	
				Occupational Health and Safety Fron	
				Accidents	
			4.3.11.2.3	Nonradiological Impacts to Public and	
				Occupational Health and Safety Fron	
				Normal Operations	
			4.3.11.2.4	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
				Occupational Health and Safety Fron	
				Accidents	
		4.3.11.3		toration Impacts to Public and Occupation	
				Safety	
		4.3.11.4		ioning Impacts to Public and Occupation	
				Safety	
	4.3.12			npacts	
				Impacts to Waste Management	
				npacts to Waste Management	
				toration Impacts to Waste Management	
	4040			ioning Impacts to Waste Management	
4.4	4.3.13				
4.4				ng Uranium Milling Region	
	4.4.1				
				Impacts to Land Use	
		4.4.1.2	•	npacts to Land Use	
		4.4.1.3		toration Impacts to Land Use	
	4.4.0	4.4.1.4		ioning Impacts to Land Use	
	4.4.2			3	
		4.4.2.1		n Impacts to Transportation	
		4.4.2.2		npacts to Transportation	
		4.4.2.3	•	toration Impacts to Transportation	
	4.4.0	4.4.2.4		ioning Impacts to Transportation	
	4.4.3	0,		pacts	
		4.4.3.1		n Impacts to Geology and Soils	
		4.4.3.2		npacts to Geology and Soils	
		4.4.3.3	•	toration Impacts to Geology and Soils	
	A A A	4.4.3.4		ioning Impacts to Geology and Soils	
	4.4.4			acts	
		4.4.4.1	Surface wa	ter Impacts	4.4-9

Section						Page
			4.4.4.1.1 4.4.4.1.2 4.4.4.1.3	Operation I Aquifer Res	on Impacts to Surface Williams Impacts to Surface Wate storation Impacts to Surf	er4.4-10 ace
			4.4.4.1.4	Decommiss	sioning Impacts to Surfa	ce
		4.4.4.2	Groundwa			
		1. 1. 1.2	4.4.4.2.1	•	on Impacts to Groundwat	
			4.4.4.2.2	Operation I	Impacts to Groundwater Operation Impacts to S	4.4-11
					(Near-Surface) Aquifer	s4.4.11
				4.4.4.2.2.2	Operation Impacts to Production and Surrou	•
				4.4.4.2.2.3	Aquifers Operation Impacts to D	
					Aquifers Below the	
			4 4 4 0 0	A culifor Do	Production Aquifers	4.4-17
			4.4.4.2.3		storation Impacts vater	1110
			4.4.4.2.4		sioning Impacts to	4.4-10
			7.7.7.2.7		er	4 4-20
	4.4.5	Ecologic	al Resources			
	1. 1.0	4.4.5.1			Ecological Resources	
		4.4.5.2		•	cological Resources	
		4.4.5.3	•	storation Imp	•	
			Ecological	Resources		4.4-22
		4.4.5.4		sioning Impa		
	4.4.6					
		4.4.6.1			Air Quality	
		4.4.6.2			ir Quality	
		4.4.6.3 4.4.6.4			eacts to Air Quality cts to Air Quality	
	4.4.7	_		• .		
	4.4.7	4.4.7.1	•		Noise	
		4.4.7.2			oise	
		4.4.7.3	•	•	acts to Noise	
		4.4.7.4			cts to Noise	
	4.4.8				Impacts	
		4.4.8.1			Historical and	
						4.4-26
		4.4.8.2	Operation	Impacts to Hi	istorical and	
						4.4-27
		4.4.8.3	•	•	acts to Historical and	
						4.4-28
		4.4.8.4			cts to Historical and	
			Cultural Re	esources		4.4-28

Section					Page
	4.4.9			es Impacts	
		4.4.9.1		Impacts to Visual/Scenic Resources	
		4.4.9.2 4.4.9.3		pacts to Visual/Scenic Resources oration Impacts to Visual/Scenic	4.4-29
					4.4-29
		4.4.9.4		oning Impacts to Visual/	
			Scenic Reso	urces	4.4-29
	4.4.10	Socioeco		S	
		4.4.10.1		Impacts to Socioeconomics	
		4.4.10.2	Operation Im	pacts to Socioeconomics	4.4-32
		4.4.10.3	Aquifer Rest	oration Impacts to Socioeconomics	4.4-33
		4.4.10.4	Decommission	oning Impacts to Socioeconomics	4.4-33
	4.4.11	Public an	d Occupation	al Health and Safety Impacts	4.4-34
				Impacts to Public and Occupational	
			Health and S	Safety	4.4-34
		4.4.11.2		pacts to Public and Occupational	
			Health and S	Safety	4.4-34
			4.4.11.2.1	Radiological Impacts to Public and	
				Occupational Health and Safety From	
				Normal Operation	4.4-34
			4.4.11.2.2	Radiological Impacts to Public and	
				Occupational Health and Safety From	
				Accidents	4.4-35
			4.4.11.2.3	Nonradiological Impacts to Public and	
				Occupational Health and Safety From	
				Normal Operations	4.4-35
			4.4.11.2.4	Nonradiological Impacts to Public and	
				Occupational Health and Safety From	
				Accidents	4.4-35
		4.4.11.3	Aguifer Rest	oration Impacts to Public and Occupatio	
				Safety	
		4.4.11.4		oning Impacts to Public and Occupation	
				Safety	
	4.4.12	Waste Ma		pacts	
		4.4.12.1	•	Impacts to Waste Management	
		4.4.12.2		pacts to Waste Management	
		4.4.12.3		oration Impacts to Waste Management.	
		4.4.12.4		oning Impacts to Waste Management	
	4.4.13				
4.5				um Milling Region	
	4.5.1				
		4.5.1.1		Impacts to Land Use	
		4.5.1.2		pacts to Land Use	
		4.5.1.3	•	oration Impacts to Land Use	
		4.5.1.4		oning Impacts to Land Use	
	4.5.2				
	· · · · · ·	4.5.2.1	•	Impacts to Transportation	
		4.5.2.2		pacts to Transportation	

Section						Page
		4.5.2.3			acts to Transportation	
		4.5.2.4			cts to Transportation	
	4.5.3	•••	and Soils In	npacts		4.5-4
		4.5.3.1			Geology and Soils	
		4.5.3.2			eology and Soils	
		4.5.3.3			acts to Geology and Soils	
		4.5.3.4			cts to Geology and Soils	
	4.5.4					
		4.5.4.1		•		
			4.5.4.1.1		on Impacts to Surface Wat	
			4.5.4.1.2		I Impacts to Surface Wate	r 4.5-10
			4.5.4.1.3	•	storation Impacts to	
					aters	4.5-10
			4.5.4.1.4		sioning Impacts to	
					ater	
		4.5.4.2	Groundwa			
			4.5.4.2.1	Construction	on Impacts to Groundwate	r4.5-11
			4.5.4.2.2	Operation I	mpacts to Groundwater	4.5-11
				4.5.4.2.2.1	Operation Impacts to	
					Shallow (Near-Surface)	
					Aquifers	4.5-11
				4.5.4.2.2.2	Operation Impacts to	
					Production and Surround	ding
					Aquifers	4.5-12
				4.5.4.2.2.3	Operation Impacts to De	ер
					Aquifers Below the	•
					Production Aquifers	4.5-16
			4.5.4.2.3	Aquifer Res	storation Impacts to	
				Groundwat	er	4.5-16
			4.5.4.2.4	Decommiss	sioning Impacts to	
					er	4.5-18
	4.5.5	Ecologic	al Resource	s Impacts		4.5-19
		4.5.5.1			Ecological Resources	
		4.5.5.2			cological Resources	
		4.5.5.3			acts to Ecological	
			•	•		4.5-23
		4.5.5.4	Decommis	sioning Impa	cts to Ecological	
					······	4.5-23
	4.5.6	Air Quali				
		4.5.6.1	•		Air Quality	
		4.5.6.2		•	r Quality	
		4.5.6.3			acts to Air Quality	
		4.5.6.4	•	•	cts to Air Quality	
	4.5.7					
		4.5.7.1			Noise	
		4.5.7.2			oise	
		4.5.7.3			acts to Noise	
		4.5.7.4	•	•	cts to Noise	

Section					Page
	4.5.8	Historical 4.5.8.1	Construction	Resources Impacts	
		4.5.8.2	Operation In	ourcesnpacts to Historical and	
		4.5.8.3	Cultural Res	ourcestoration Impacts to Historical and	
		4.5.8.4	Decommissi	ourcesoning Impacts to Historical and	
				ources	
	4.5.9			es Impacts	
		4.5.9.1		Impacts to Visual/Scenic Resources	
		4.5.9.2 4.5.9.3	Aquifer Res	npacts to Visual/Scenic Resources toration Impacts to Visual/	
				purces	4.5-30
		4.5.9.4		oning Impacts to Visual/	4 5 00
	4.5.40	· ·		ources	
	4.5.10			s	
				Impacts to Socioeconomics	
				npacts to Socioeconomics	
			•	toration Impacts to Socioeconomics	
	4 5 44			oning Impacts to Socioeconomics	
	4.5.11			al Health and Safety Impacts	4.5-35
		4.5.11.1		Impacts to Public and Occupational	4 5 05
		45440		Safety	4.5-35
		4.5.11.2		npacts to Public and Occupational	4 5 05
				Safety	4.5-35
			4.5.11.2.1	Radiological Impacts to Public and	
				Occupational Health and Safety From	4 5 05
			454400	Normal Operations	4.5-35
			4.5.11.2.2	Radiological Impacts to Public and Occupational Health and Safety From	4.5.05
			4.5.11.2.3	Accidents Nonradiological Impacts to Public and	4.5-35
			4.3.11.2.3	Occupational Health and Safety From	
				Normal Operations	15-35
			4.5.11.2.4	Nonradiological Impacts to Public and	4.5-55
			4.5.11.2.4	Occupational Health and Safety From Accidents	4 5-36
		45113	Aquifer Rest	toration Impacts to Public and	00
			•	al Health and Safety	4.5-36
		4.5.11.4		oning Impacts to Public and Occupation	
				Safety	
	4.5.12	Waste Ma		npacts	
		4.5.12.1		Impacts to Waste Management	
		4.5.12.2		npacts to Waste Management	
				toration Impacts to Waste Management.	
		4.5.12.4		oning Impacts to Waste Management	
	4.5.13			ormig impacto to videto intriagoriioni	4.5-38

Section	on		Page
5	CUMU	JLATIVE EFFECTS	5-1
	5.1	Introduction	5-1
	5.2	Other Past, Present, and Reasonably Foreseeable Future Actions in the	
		Four Regions	
		5.2.1 Uranium Recovery Sites	
		5.2.2 EISs as Indicators of Present and RFFAs	
	5.3	Concurrent Actions	
		5.3.1 Wyoming West Uranium Milling Region	
		5.3.2 Wyoming East Uranium Milling Region	
		5.3.3 Nebraska-South Dakota-Wyoming Uranium Milling Region	
	- 1	5.3.4 Northwestern New Mexico Uranium Milling Region	
	5.4	Approaches to Conducting a Site-Specific Cumulative Effects Analysis	
	5.5	References	5-27
6	ENVI	RONMENTAL JUSTICE	6-1
Ū	6.1	Environmental Justice Analysis	
		6.1.1 Background and Guidance	
		6.1.2 Identifying Minority and Low-Income Populations in the Four	
		Geographic Uranium Milling Regions Considered in This GEIS	6-5
	6.2	Wyoming West Uranium Milling Region	
	6.3	Wyoming East Uranium Milling Region	6-14
	6.4	Nebraska-South Dakota-Wyoming Uranium Milling Region	
	6.5	Northwestern New Mexico Uranium Milling Region	
	6.6	Summary	
	6.7	References	6-19
7	POTE	NTIAL BEST MANAGEMENT PRACTICES, MITIGATION	
,		SURES, AND MANAGEMENT ACTIONS TO MITIGATE ADVERSE	
		RONMENTAL IMPACTS	7-1
	7.1	Introduction	
	7.2	Best Management Practices	
	7.3	Management Actions	
	7.4	Potential Best Management Practices, Management Actions, and	
		Mitigation Measures	7-2
	7.5	References	
•	- N 11 / 11	CONNECTAL MONITORING ACTIVITIES	0.4
8		RONMENTAL MONITORING ACTIVITIESIntroduction	
	8.1 8.2	Radiological Monitoring	
	0.2	8.2.1 Airborne Radiation Monitoring Program	
		8.2.2 Direct Radiation Monitoring Program	
		8.2.3 Soils and Sediments Monitoring	
		8.2.4 Vegetation, Food, and Fish Monitoring	
		8.2.5 Surface Water Monitoring	
		8.2.6 Groundwater Monitoring	
		· · · · · · · · · · · · · · · · · · ·	

Sect	ion			Page
	8.3	Physiod 8.3.1	themical Monitoring	8-3
			8.3.1.2 Groundwater Quality Monitoring	
		8.3.2	Well Field and Pipeline Flow and Pressure Monitoring	
	8.4	Ecologic	cal Monitoring	
	8.5	Referen	nces	8-7
9	CONS	SULTATIO	DNS	9-1
10	SUMI	MARY OF	ENVIRONMENTAL CONSEQUENCES	10-1
11	LIST	OF PREP	PARERS	11-1
	11.1 11.2	Southwe	clear Regulatory Commission Contributorsest Research Institute [®] and Center for Nuclear Waste Regulato	ory
			es (CNWRA®) Contributors	
	11.3	CNWRA	A Subcontractors	11-3
12	GLOS	SSARY		12-1
۸۵۵	ENDIX A	50	OPING SUMMARY REPORT	Λ 1
	ENDIX F	R PO	TENTIALLY APPLICABLE FEDERAL STATUTES,	A-1
/ \l I		RE	GULATIONS, AND EXECUTIVE ORDERS	B–1
APP	ENDIX (SUI	MMARY OF CONVENTIONAL URANIUM MILLING	
		TE	CHNOLOGIES	C–1
APP	ENDIX [LTURAL AND HISTORICAL RESOURCE MANAGEMENT	
۸DD	ENDIX E		OCESSESZARDOUS CHEMICALS	
	ENDIX E		SCRIPTION OF PROCESSES FOR REVIEW OF	⊏−1
/ \l I			MULATIVE EFFECTS	F–1
APP	ENDIX (PU	BLIC COMMENTS ON THE DRAFT GENERIC ENVIRONMEN	1TAL
		IMF	PACT STATEMENT AND NRC RESPONSES	G–1

FIGURES

Figure	Page	Э
1.1-1	Four Geographic Regions Used as a Framework for the Analyses Presented in This GEIS1-2	2
1.1-2	Major Uranium Reserves Within the United States1-3	
1.4-1	Structure of This GEIS1-	
1.7-1	General Flow Diagram of the NRC Licensing Process for	,
1.7-1	10 CFR Part 40 Licenses1-19	a
	10 Of IX Fait 40 Licenses	ט
2.1-1	Simplified Cross Section of Sandstone Uranium Roll-Front Deposits Formed by Regional Groundwater Migration2-3	3
2.1-2	Schematic Diagram of the Types of Tabular Stratabound Uranium Deposits in the Grants Uranium District, New Mexico2-	3
2.1-3	Layout of the Crow Butte Uranium Project in Dawes County, Nebraska2-5	
2.1-4	Well Heads and a Header House at Smith Ranch, Converse County, Wyoming	
2.3-1	Schematic Diagram of a Well Field Showing Typical Injection/Production Well	
	Patterns, Monitoring Wells, Manifold Buildings, and Pipelines2-	
2.3-2	Manifold Inside Well Field Header House at an ISL Leach Facility2-10	
2.3-3	Computerized Meter for Monitoring Well Field Flow Rates2-1	1
2.3-4	Cross Section of a Typical Injection, Production, or Monitoring Well Completed Using the Underreamed Method2-13	2
2.4-1	Idealized Schematic Cross Section To Illustrate Ore-Zone Geology and Lixiviant	,
2.4-1		0
0.4.0	Migration From an Injection Well to a Production Well	
2.4-2	Flow Diagram of an ISL Uranium Recovery Process	
2.4-3	Typical Ion-Exchange Vessels in an ISL Facility2-22	
2.4-4	A Typical Thickener for an ISL Uranium Processing Facility2-24	4
2.4-5	Typical Vacuum Dryer for Uranium Yellowcake Processing at an ISL Uranium Processing Facility2-25	5
2.4-6	Labeled and Placarded 208-L [55-gal] Drum Used for Packaging and	
2.4 0	Shipping Yellowcake2-20	6
2.5-1	Schematic Diagram of Groundwater Sweep During Aquifer Restoration 2-28	
2.7-1	Downwind Distance Versus Relative Air Concentration (Which Is Proportional)
	to Dose)	5
3.1-1	Wyoming West Uranium Milling Region With Current and Potential	_
	ISL Milling Sites	2
3.1-2	Wyoming East Uranium Milling Region With Current and Potential	_
	ISL Milling Sites	3
3.1-3	Nebraska-South Dakota-Wyoming Uranium Milling Region With Current	
	and Potential ISL Milling Sites	4
3.1-4	Northwestern New Mexico Uranium Milling Region With Current and Potential ISL	_
	Milling Sites	5
3.1-5	Simplified Cross Section of Sandstone Uranium Roll-Front Deposits Formed	
	by Regional Groundwater Migration	7
3.2-1	Wyoming West Uranium Milling Region General Map With Current and Future	
	Uranium Milling Site Locations	2
3.2-2	Map Showing Outline of the Wyoming West Uranium Milling Region and	
	Locations of the Crooks Gap Uranium District in the Great Divide Basin	
	and the Gas Hills Uranium District in the Wind River Basin	3

Figure		Page
3.2-3 3.2-4	Wyoming West Uranium Milling Region Transportation Corridor	
3.2-5	Sweetwater Arch to the Great Divide Basin and the Wind River Basin Stratigraphic Section of Tertiary Age Formations in the Great Divide Basin	
	and Wind River Basin of Central Wyoming	
3.2-6	Watersheds in the Wyoming West Uranium Milling Region	
3.2-7	Ecoregions of the Wyoming West Uranium Milling Region	
3.2-8	Antelope Wintering Areas for the Wyoming West Uranium Milling Region	
3.2-9	Big Horn Wintering Areas for the Wyoming West Uranium Milling Region	
3.2-10	Elk Wintering Areas for the Wyoming West Uranium Milling Region	
3.2-11	Moose Wintering Areas for the Wyoming West Uranium Milling Region	
3.2-12 3.2-13	Mule Deer Wintering Areas for the Wyoming West Uranium Milling Region Sage-Grouse/Lek Nesting Areas for the Wyoming West Uranium	. 3.2-32
	Milling Region	. 3.2-33
3.2-14	White Tailed Deer Wintering Areas for the Wyoming West Uranium	
0 0 45	J - J -	-
3.2-15	Air Quality Attainment Status for Wyoming and Surrounding Areas	. 3.2-49
3.2-16	Prevention of Significant Deterioration Class I Areas in the Wyoming West	2 2 54
2 2 47	Uranium Milling Region and Surrounding Areas	
3.2-17	Comparison of Noise Levels Associated with Common Activities	
3.2-18	General Divisions of the American Plains	. 3.2-55
3.2-19	Regional Distribution of Native American Tribes in the Early 1800s in	2 2 60
2 2 20	Wyoming, South Dakota, and Nebraska	. 3.2-60
3.2-20	BLM Visual Resource Classifications for the Wyoming West Uranium	2 2 66
2 2 24	Milling Region	. 3.2-00
3.2-21	Wyoming West Uranium Milling Region With Population	
3.2-22	Average Annual Background Radiation in the United States	. 3.2-01
3.3-1	Wyoming East Uranium Milling Region General Map With Past, Current,	222
3.3-2	and Future Uranium Milling Site Locations	3.3-2
3.3-2	Map Showing Outline of the Wyoming East Region and Locations of the	ho
	Pumpkin Buttes and Monument Hill Districts in the Powder River Basin and t	
222	Shirley Basin Uranium District in the Shirley Basin	s.s-s
3.3-3	Wyoming East Uranium Milling Region General Map With Current and Future Uranium Milling Site Transportation Corridor	3.3-5
224	· · · · · · · · · · · · · · · · · · ·	s.s-s
3.3-4	Index and Structure Map of East-Central Wyoming Showing Relation of the Sweetwater Arch and Laramie Range to the Powder River Basin and	
	the Shirley Basin	220
225	Stratigraphic Section of Tertiary Age Formations in the Powder River Basin	s.s-9
3.3-5		2 2 40
226	and Shirley Basin of WyomingWatersheds Within the Wyoming East Uranium Milling Region	
3.3-6	, , ,	
3.3-7 3.3-8	Ecoregions of the Wyoming East Uranium Milling Region	
	Antelope Wintering Area for the Wyoming East Uranium Milling Region	
3.3-9	Big Horn Wintering Area for the Wyoming East Uranium Milling Region	
3.3-10	Elk Wintering Area for the Wyoming East Uranium Milling Region	
3.3-11	Moose Wintering Area for the Wyoming East Uranium Milling Region Mule Deer Wintering Area for the Wyoming East Uranium Milling Region	
3.3-12	initie deel viilieliing alea ioi the vyonning east oranium iviiliing region	. ა.ა-ა0

Figure	Pa	age
3.3-13	Sage-Grouse/Lek Nesting Areas for the Wyoming East Area Uranium	. 04
3.3-14	Milling Region	5-31
	Milling Region	3-32
3.3-15	Air Quality Attainment Status for Wyoming and Surrounding Areas3.3	-39
3.3-16	Prevention of Significant Deterioration Class I Areas in the Wyoming	
	East Uranium Milling Region and Surrounding Areas	-41
3.3-17	BLM Visual Resource Classifications for the Wyoming East Uranium	
0 0 40	Milling Region	i-45
3.3-18	Wyoming East Uranium Milling Region With Population	-49
3.4-1	Nebraska-South Dakota-Wyoming Uranium Milling Region General Map With Current (Crow Butte, Nebraska) and Potential Future Uranium	
	Milling Site Locations	.4-2
3.4-2	Map Showing the Nebraska-South Dakota-Wyoming Uranium Milling Region and	
	Uranium Milling Sites in the Black Hills Uranium Districts in South Dakota and	
	Wyoming and in the Crow Butte Uranium District in Nebraska	4-3
3.4-3	Nebraska-South Dakota-Wyoming Uranium Milling Region	
	Transportation Corridor	4-5
3.4-4	Outcrop Map of the Inyan Kara Group in the Black Hills of Western South	
	Dakota and Northeastern Wyoming Showing the Locations of Principal	4.0
3.4-5	Uranium Mining Districts	4-9
3.4-3	and Northeastern Wyoming3.4	1_10
3.4-6	Schematic Cross Section Through a Typical Inyan Kara Roll-Front Deposit	-10
0.10	Showing Differences in U ₃ O ₈ Concentration Between "Fresh"	
	(i.e., Unoxidized) and "Altered" Ground	-12
3.4-7	Bedrock Geology and Major Structural Features of the Crawford Basin 3.4	
3.4-8	Generalized Stratigraphic Units in the Crow Butte Area of	
	Northwestern Nebraska	-15
3.4-9	Location of Oxidation-Reduction Fronts Detected During Exploration Drilling	
0.4.40	Within the Chadron Sandstone in Northwestern Nebraska	-16
3.4-10	Watersheds Within the Nebraska-South Dakota-Wyoming Uranium	1 4 0
3.4-11	Milling Region	-18
3. 4- 11	Milling Region	-20
3.4-12	Antelope Wintering Areas for the Nebraska-South Dakota-Wyoming Uranium	-23
0.1 12	Milling Region	-32
3.4-13	Big Horn Wintering Areas for the Nebraska-South Dakota-Wyoming	-
	Uranium Milling Region	-33
3.4-14	Elk Wintering Areas for the Nebraska-South Dakota-Wyoming	
	Uranium Milling Region3.4	-34
3.4-15	Moose Wintering Areas for the Nebraska-South Dakota-Wyoming	
	Uranium Milling Region	-35
3.4-16	Mule Deer Wintering Areas for the Nebraska-South Dakota-Wyoming	
0 4 4 7	Uranium Milling Region	-36
3.4-17	Sage-Grouse/Lek Nesting Areas for the Nebraska-South Dakota-Wyoming Uranium Milling Region	ר פו
	01a111u111 WIIIIIIU NEUIUI	-31

Figure		Page
3.4-18	White Tailed Deer Wintering Areas for the Nebraska-South Dakota-Wyomir	
	Uranium Milling Region	3.4-38
3.4-19	Air Quality Attainment Status for Nebraska-South Dakota-Wyoming	0 4 5 4
0.4.00	Uranium Milling Region and Surrounding Areas	
3.4-20	Prevention of Significant Deterioration Class I Areas in the Nebraska-South	
3.4-21	Dakota-Wyoming Uranium Milling Region	
3.4-21 3.5-1	Nebraska-South Dakota-Wyoming Uranium Milling Region With Population Northwestern New Mexico Uranium Milling Region General Map with	3.4-77
3.5-1	Current and Future Uranium Milling Site Locations	3 5-2
3.5-2	Map Showing Outline of the Northwestern New Mexico Region and the Loc	
0.0 2	of the Grants Uranium District Along the Southern Margin of	ation
	the San Juan Basin	3.5-3
3.5-3	Northwestern New Mexico Uranium Milling Region Transportation	
	Corridor Locations	3.5-5
3.5-4	Index Map of the Grants Uranium District, San Juan Basin, New Mexico,	
	Showing Eight Subdistricts	3.5-8
3.5-5	Generalized Stratigraphic Section of Upper Jurassic and Cretaceous	
	Formations in the Grants Uranium District	3.5-9
3.5-6	Generalized Geologic Section Showing the Stratigraphic Relations of the	
	Morrison Formation Between the Ambrosia Lake and Laguna Areas	3.5-10
3.5-7	Schematic Diagram of the Different Types of Uranium Deposits in the	
	Morrison Formation, Grants Uranium District, New Mexico	
3.5-8	Watersheds in the Northwestern New Mexico Uranium Milling Region	
3.5-9	Ecoregions for the Northwestern New Mexico Uranium Milling Region	3.5-23
3.5-10	Elk Winter Habitat and Calving Areas for the Northwestern New Mexico	2 5 20
2 5 11	Uranium Milling Region	
3.5-11 3.5-12	Wind Rose for Gallup, New Mexico, Airport for 1991	ა.၁-ა၁
3.5-12	Milling Region and Surrounding Areas	3 5-37
3.5-13	Prevention of Significant Deterioration Class I Areas in the Northwestern	3.3-31
0.0-10	New Mexico Uranium Milling Region and Surrounding Areas	3 5-38
3.5-14	Distribution of Recorded Archaeological Sites in McKinley and	0.0 00
0.0 11	Cibola Counties, New Mexico	3.5-42
3.5-15	Paleoindian Sites	3.5-43
3.5-16	Distribution of Archaic-Period Sites	3.5-44
3.5-17	Distribution of Ancestral Puebloan Sites	3.5-46
3.5-18	Distribution of Historic Pueblo Sites	3.5-50
3.5-19	Distribution of Navajo Archaeological Sites	3.5-52
3.5-20	BLM Visual Resource Classifications for the Northwestern New Mexico	
	Uranium Milling Region	3.5-61
3.5-21	Northwestern New Mexico Uranium Milling Region With Population	3.5-64
4.2-1	Sound Levels Around a Typical Water Well Work Site	4.2-40
4.2-2	Community Surveys of Noise Annoyance	

Figure		Page
6.1-1	Environmental Justice Process Flow Chart	6-3
6.1-2	Affected Minority and Low-Income Population for the Wyoming West Uranium	
	Milling Region	6-7
6.1-3	Affected Minority and Low-Income Population for the Wyoming East Uranium	
	Milling Region	6-8
6.1-4	Affected Minority and Low-Income Population for the Nebraska-South	
	Dakota-Wyoming Uranium Milling Region	.6-10
6.1-5	Affected Minority and Low-Income Populations for the Northwestern New Mexic	
	Uranium Milling Region	. 6-11

TABLES

Table	Pa	age
1.7-1	New Source Review Permit Summary Information for Nebraska, New Mexico, South Dakota, and Wyoming	1-23
2.2-1	Typical Baseline Water Quality Parameters and Indicators	2-7
2.4-1	Typical Lixiviant Chemistry	
2.6-1	Estimated Decommissioning and Reclamation Waste Volumes (yd³) for	
	Offsite Disposal, Smith Ranch In-Situ Leach Facility	2-32
2.7-1	Combustion Engine Exhaust Sources for the Gas Hills In-Situ Leach Satellite	
	Facility During Construction, Operations, Reclamation, and Decommissioning	2-34
2.7-2	Estimated Particulate (Fugitive Dust) and Gaseous (Diesel Combustion	
	Products) Emissions for the Crownpoint, New Mexico, <i>In-Situ</i> Leach Facility	2 24
272	Construction Phase	2-34
2.7-3	Estimated Flow Rates and Constituents in Liquid Waste Steams for the Highland <i>In-Situ</i> Leach Facility	2 26
2.8-1	Estimated Annual Vehicle Trips for Phases of <i>In-Situ</i> Leach Facility Lifecycle	
2.11-1	Size of Permitted Areas for <i>In-Situ</i> Leach Facilities	
2.11-2	Common Bulk Chemicals Required at the Project Processing Sites	
2.11-3	Onsite Quantities of Process Chemicals at <i>In-Situ</i> Leach Facilities	
2.11-4	Irigaray Water Quality Summary for Designated Aquifer Restoration Wells	2-49
2.11-5	Baseline Groundwater Conditions, Aquifer Restoration Goals, and Actual Final	
	Restoration Values the U.S. Nuclear Regulatory Commission Approved for the	
	Q-Sand Pilot Well Field, Smith Ranch, Wyoming	2-50
3.2-1	Land Surface Ownership and General Use in the Wyoming West Uranium	
	Milling Region	3.2-4
3.2-2	Average Annual Daily Traffic Counts for Roads in the Wyoming West	
	Uranium Milling Region	3.2-6
3.2-3	Representative Transportation Routes for Yellowcake Shipments From the	
0.0.4	Wyoming West Uranium Milling Region	3.2-7
3.2-4	Primary Watersheds in the Wyoming West Uranium Milling Region Range of	2 42
3.2-5	Designated Uses of Water Bodies Within Each Watershed	
3.2-6	Information on Two Climate Stations in the Wyoming West Uranium	2-33
5.2-0	Milling Region	2-45
3.2-7	Climate Data for Stations in the Wyoming West Uranium Milling Region3.	
3.2-8	National Ambient Air Quality Standards	
3.2-9	Allowable Prevention of Significant Deterioration Class I and Class II Areas 3.	2-48
3.2-10	U.S. Environmental Protection Agency Class I Prevention of Significant	
	Deterioration Areas in Wyoming	2-50
3.2-11	National Register Listed Properties in Counties Included in the Wyoming West	
	Uranium Milling Region3.	2-58
3.2-12	List of Tribal Contacts for Tribes With Interests in Nebraska, Montana,	
0.0.45	South Dakota, and Wyoming	2-62
3.2-13	Summary of the Affected Environment Within the Wyoming West Uranium	0 0-
2 2 4 4	Milling Region	2-67
3.2-14	Wyoming West Uranium Milling Region	2_60
	vvyonning vvcst Oranium ivining incyloti	∠-09

Table		Page
3.215	U.S. Bureau of Census State Income Information for the Region Surrounding the Wyoming West Uranium Milling Region	
3.2-16	U.S. Bureau of Census County and Native American Income Information	0.2 72
0.2 .0	for the Wyoming West Uranium Milling Region	3.2-73
3.2-17	U. S. Bureau of Census Housing Information for Wyoming	
3.2-18	Employment Structure of the Wind River Indian Reservation Within the	
0.0.40	Affected Area	3.2-78
3.2-19	2007 State and Local Annual Sales and Use Tax Distribution of Affected	2 2 70
2 2 20	Counties Within the Wyoming West Uranium Milling Region	
3.2-20	Public Radiation Doses	3.2-01
3.3-1	Land Surface Ownership and General Use in the Wyoming East Uranium Milling Region	224
3.3-2	Average Annual Daily Traffic Counts for Roads in the Wyoming East	3.3 -4
3.3-2	Uranium Milling Region	226
3.3-3	Representative Transportation Routes for Yellowcake Shipments From the	3.3-0
3.3-3	Wyoming East Uranium Milling Region	3 3-7
3.3-4	Primary Watersheds in the Wyoming East Uranium Milling Region Range of	
J.J- T	Designated Uses of Water Bodies Within Each Watershed	
3.3-5	Information on Two Climate Stations in the Wyoming East Uranium	5.5-1-
0.0 0	Milling Region	3 3-36
3.3-6	Climate Data for Stations in the Wyoming East Uranium Milling Region	
3.3-7	U.S. Environmental Protection Agency Class I Prevention of Significant	0.0 00
0.0 1	Deterioration Areas in Wyoming	3 3-40
3.3-8	National Register Listed Properties in Counties Included in the	0.0 10
0.0 0	Wyoming East Uranium Milling Region	3.3-42
3.3-9	Summary of Affected Environment Within the Wyoming East Uranium	
	Milling Region	3.3-46
3.3-10	2000 U.S. Bureau of Census Population and Race Categories of the	
	Wyoming East Uranium Milling Region	3.3-48
3.3-11	U.S. Bureau of Census State Income Information for Wyoming East	
	Uranium Milling Region	3.3-50
3.3-12	U.S. Bureau of Census County Income Information for Wyoming East	
	Uranium Milling Region	3.3-51
3.3-13	U.S. Bureau of Census Housing Information for the Wyoming East	
	Uranium Milling Region	3.3-52
3.3-14	2007 State and Local Annual Sales and Use Tax Distribution of Affected	
	Counties Within the Wyoming East Uranium Milling Region	3.3-57
3.4-1	Land Surface Ownership and General Use in the Nebraska-South	
	Dakota-Wyoming Uranium Milling Region	3.4-4
3.4-2	Average Annual Daily Traffic Counts for Roads in the Nebraska-	
	South Dakota-Wyoming Uranium Milling Region	3.4-7
3.4-3	Representative Transportation Routes for Yellowcake Shipments From the	
	Nebraska-South Dakota-Wyoming Uranium Milling Region	3.4-7
3.4-4	Primary Watersheds in the Nebraska-South Dakota-Wyoming Uranium	
	District and Range of Generic Designated Uses of Water Bodies Within	
	Each Watershed	3.4-19
3.4-5	Fishes of the Angostura Reservoir, Cheyenne River Watershed	
3.4-6	Fishes of the Niobrara River Watershed	3.4-43

Table		Page
3.4-7	Information on Three Climate Stations in the Nebraska-South Dakota- Wyoming Uranium Milling Region	.3.4-51
3.4-8	Climate Data for Stations in the Nebraska-South Dakota-	
3.4-9	Wyoming Uranium Milling Region	.3.4-51
3.4-9	Deterioration Areas in Wyoming, South Dakota, Nebraska, and Montana	.3.4-55
3.4-10	National Register Listed Properties in Counties Included in the Nebraska-Sou	
	Dakota-Wyoming Uranium Milling Region	.3.4-63
3.4-11	Summary of Affected Areas Within the Nebraska-South Dakota- Wyoming Uranium Milling Region	2 4 72
3.4-12	2000 U.S. Bureau of Census Population and Race Categories of Nebraska	3.4-72 3.4-73
3.4-13	2000 U.S. Bureau of Census Population and Race Categories	
	of South Dakota	
3.4-14	2000 U.S. Bureau of Census Population and Race Categories of Northweste Wyoming	
3.4-15	U.S. Bureau of Census State Income Information for the Nebraska-South	3.4-76
0.1 10	Dakota-Wyoming Uranium Milling Region	.3.4-79
3.4-16	U.S. Bureau of Census County and Native American Income Information for	
0.4.47	the Nebraska-South Dakota-Wyoming Uranium Milling Region	
3.4-17	U.S. Bureau of Census Housing Information for Nebraska	
3.4-18	U.S. Bureau of Census Housing Information for South Dakota	
3.4-19	U.S. Bureau of Census Housing Information for Wyoming	.3.4-84
3.4-20	Employment Structure of the Pine Ridge Indian Reservation Within the Affected Area	3 4-86
3.4-21	2007 State and Local Annual Sales and Use Tax Distribution of Affected	
	Counties Within Wyoming	.3.4-91
3.5-1	Land Surface Ownership and General Use in the Northwestern New Mexico	0 = 4
	Uranium Milling Region	
3.5-2	Average Annual Daily Traffic Counts for Roads in the Northwestern New Mex Uranium Milling Region	
3.5-3	Representative Transportation Routes for Yellowcake Shipments From the	
3.0-3	Northwestern New Mexico Uranium Milling Region	
3.5-4	Named Uranium Deposits in New Mexico and Corresponding Watersheds	3 5-16
3.5-5	Primary Watersheds in New Mexico, Designated Uses, and	
	Known Impairments	.3.5-16
3.5-6	Native Fish Species Found in New Mexico	.3.5-27
3.5-7	Information on Two Climate Stations in the Northwestern New Mexico	
	Uranium Milling Region	. 3.5-33
3.5-8	Climate Data for Stations in the Northwestern New Mexico Uranium	0.5.00
0.5.0	Milling Region	3.5-33
3.5-9	U.S. Environmental Protection Agency Class I Prevention of Significant	0.5.04
0 5 40	Deterioration Areas in New Mexico and Arizona	
3.5-10	Number of Recorded Sites by Time Period and County	
3.5-11	Cebolleta Mesa and Pecos Chronologies	
3.5-12	National Register Listed Properties in Counties Included in the Northwestern	
	New Mexico Uranium Milling Region	. ა.၁-53

Table		Page
3.5-13 3.5-14	Known Culturally Significant Places in McKinley and Cibola Counties	
3.5-15	Cibola Counties, New Mexico Summary of Affected Environment Within the Northwestern New Mexico	.3.5-57
3.5-16	Uranium Milling Region	.3.5-62
3.5-17	Northwestern New Mexico Uranium Milling Region	.3.5-63
3.5-18	New Mexico Uranium Milling RegionU.S. Bureau of Census County Income Information for the Northwestern	.3.5-66
3.5-19	New Mexico Uranium Milling Region	.3.5-68
3.5-20	New Mexico Uranium Milling Region Employment Structure of Native American Communities Within the Affected	.3.5-69
	Environment of the Northwestern New Mexico Uranium Milling Region	
3.5-21 3.5-22	Net Taxable Values for Affected Counties Within New Mexico for 2006 Percent Change in Tax Values From 2005 to 2006 for the Affected Counties	
3.5-23	Within New Mexico Percent Distribution of New Mexico Property Tax Obligations Within Affected	
	Counties for 2006	.3.5-74
4.2-1	Average Noise Levels at 15 m [50 ft] From Representative Construction Heavy Equipment	4 2-38
4.2-2	Dose to Offsite Receptors From <i>In-Situ</i> Leach Facilities	
4.2-2	Pertinent Regulations for Chemicals Used at <i>In-Situ</i> Leach Facilities	
		.4.2-58
5.2-1	Past, Existing, and Potential Uranium Recovery Sites in Wyoming, South Dakota, Nebraska, and New Mexico	5-4
5.2-2	Draft and Final Environmental Impact Statements (EISs) Related to the Wyoming West Uranium Milling Region	
5.2-3	Draft and Final Environmental Impact Statements (EISs) Related to the Wyoming East Uranium Milling Region	
5.2-4	Draft and Final Programmatic or Large-Scale Environmental Impact Stateme (EISs) Related to One or Both of the Wyoming Regional Study Areas	nts
5.2-5	Draft and Final Environmental Impact Statements (EISs) Related to Nebraska-South Dakota-Wyoming Uranium Milling Region	
5.2-6	Draft and Final Environmental Impact Statements (EISs) Related to the Northwestern New Mexico Uranium Milling Region	
5.3-1	Other Actions Concurrent With Uranium Recovery in the Wyoming West Uranium Milling Region	
F 2 2		
5.3-2	Coal Mining Projects as Identified by the Wyoming Mining Association	5-17
5.3-3	Other Actions Related to or Conflicting With Uranium Recovery in the Wyoming East Uranium Milling Region	
5.3-4	Other Actions Concurrent With Uranium Recovery in Nebraska-South Dakota Wyoming Uranium Milling Region	
5.3-5	Other Actions Concurrent With Uranium Recovery in the Northwestern	5 20
5.5 0	New Mexico Uranium Milling Region	5-24

Table	P	age
6.1-1	Minority and Low-Income Populations in the Four Geographic Uranium Milling Regions Considered in This Generic Environmental Impact Statement	6-6
7.4-1	Summary of Potential Best Management Practices and Management Actions	7-3
8.2-1	Typical Baseline Water Quality Parameters and Indicators for Groundwater	8-4
10-1 10-2 10-3	9 - 9 -	
10-4	Summary of Impacts for the Northwestern New Mexico Uranium Milling Region	10-48

EXECUTIVE SUMMARY

BACKGROUND

The Atomic Energy Act of 1954 and the Uranium Mill Tailings Radiation Control Act of 1978 (UMTRCA) authorize the U.S. Nuclear Regulatory Commission (NRC) to issue licenses for the possession and use of source material and byproduct material. The statutes require NRC to license facilities that meet NRC regulatory requirements that were developed to protect public health and safety from radiological hazards. *In-situ* leach (ISL) uranium recovery facilities must meet NRC regulatory requirements in order to obtain this license to operate.

NRC designed the licensing process to assure the safe operation of ISL facilities. In addition to information for a safety evaluation review, license applicants must submit an environmental report as part of their license application. Under the NRC's environmental protection regulations in the Code of Federal Regulations, Title 10, Part 51 (10 CFR Part 51), which implement the National Environmental Policy Act (NEPA), issuance of a license to possess and use source material for uranium

Generic Environmental Impact Statement (GEIS)

A GEIS is an environmental impact statement that assesses the scope of the environmental effects that would be associated with an action (such as issuing a license for an ISL facility) at numerous sites. The Commission directed the NRC staff to prepare the GEIS to cover as many of the potential uranium recovery sites as possible.

Supplemental Environmental Impact Statement (SEIS)

A supplemental EIS updates or supplements an existing EIS (such as the GEIS). The Commission directed the NRC staff to issue site-specific supplements to the GEIS for each new license application.

milling requires an environmental impact statement (EIS) or a supplement to an EIS.

NRC prepared the Generic Environmental Impact Statement for *In-Situ* Leach Uranium Milling Facilities (GEIS) to help fulfill this requirement. The GEIS was prepared to assess the potential environmental impacts associated with the construction, operation, aquifer restoration, and decommissioning of an ISL facility in four specified geographic areas. The intent of the GEIS is to determine which impacts would be essentially the same for all ISL facilities and which ones would result in varying levels of impacts for different facilities, thus requiring further site-specific information to determine the potential impacts. As such, the GEIS provides a starting point for NRC's NEPA analyses on site-specific license applications for new ISL facilities, as well as for applications to amend or renew existing ISL licenses.

PURPOSE AND NEED

Commercial uranium recovery companies have approached NRC with plans to submit a number of license applications for new uranium recovery facilities and for the restart or expansion of existing facilities in the next several years. The large majority of these potential applications would involve use of the ISL process. The companies have indicated that these new, restarted, and expanded ISL facilities would be located in Wyoming, South Dakota, Nebraska, and New Mexico.

NRC is the regulatory authority responsible for issuing a source material license for an ISL facility in those four states. 10 CFR Part 51 regulations require evaluating the environmental impacts of the ISL facility as part of the licensing process. Recognizing that the technology for ISL uranium milling is relatively standardized, that the applications may be submitted over a relatively short period of time, and that the potential ISL facilities would be located in relatively

EXECUTIVE SUMMARY (continued)

discrete regions in the western United States, NRC decided to prepare a GEIS to avoid unnecessary duplicative efforts and to identify environmental issues of concern to focus on in site-specific environmental reviews. In this way, NRC could increase the efficiency and consistency in its site-specific environmental review of license applications for ISL facilities and so provide an option for applicants to use and licensees to continue to use the ISL process for uranium recovery.

THE PROPOSED FEDERAL ACTION AND ALTERNATIVES

In states where NRC is the regulatory authority over the licensing of uranium milling (including the ISL process), NRC has a statutory obligation to assess each site-specific license application to ensure it complies with NRC regulations before issuing a license. The proposed federal action is to grant an application to obtain, renew, or amend a source material license for an ISL facility.

Under NRC's environmental protection regulations at 10 CFR 51.20(b)(8), issuing a license to possess and use source material to a uranium milling facility is identified as a major federal action that requires the preparation of an EIS or a supplement to an EIS. NRC will prepare a SEIS for new ISL facility license applications. NRC will prepare an EA, SEIS or EIS for applications to amend or renew an existing ISL facility license.

The Proposed Federal Action

To grant applications to obtain, renew, or amend source material licenses for an ISL facility.

Purpose for the Proposed Federal Action

To provide an option for an applicant to use or a licensee to continue to use ISL technology for uranium recovery

The environmental review requirements for a material license are in 10 CFR Part 51. NRC's public health and safety requirements for ISL facilities are found in 10 CFR Parts 20 and 40. Parts 20, 40, and 51 require applicants to provide NRC with sufficient information to evaluate the impacts to public health and safety and the environment during the life-cycle of the ISL facility. NRC then prepares safety and environmental reviews that are used by NRC officials to decide whether to grant the source material license.

In reviewing an ISL license application, NRC will use the GEIS as starting point for its site-specific environmental reviews. NRC will evaluate site-specific data and information to determine whether the applicant's proposed activities and the site characteristics are consistent with those evaluated in the GEIS. NRC will then determine which sections of the GEIS can be incorporated by reference and which impact conclusions can be adopted in the site-specific environmental review, and whether additional data or analysis is needed to determine the environmental impacts to a specific resource area. Additionally, the GEIS provides guidance in the evaluation for certain impact analyses (e.g., cumulative impacts, environmental justice) for which the GEIS did not make impact conclusions. No decision on whether to license an ISL facility will be made based on the GEIS alone. The licensing decision will be based, in part, on a site-specific environmental analysis that makes use of the GEIS.

Uranium milling techniques are designed to recover the uranium from uranium-bearing ores. Various physical and chemical processes may be used, and selection of the uranium milling technique depends on the physical and chemical characteristics of the ore deposit and the attendant cost considerations. Generally, the ISL process is used to recover uranium from low-grade ores or deeper deposits that are not economically recoverable by conventional mining and milling techniques. In the ISL process, a leaching agent, such as oxygen with sodium carbonate, is added to native groundwater and injected through wells into the subsurface ore body to mobilize the uranium. The leach solution containing the mobilized uranium is pumped from there to the surface processing plant, and then ion exchange separates the uranium from the solution. After additional purification and drying, the resultant product, a mixture of uranium oxides also known as "yellowcake," is placed in 55-gallon drums prior to shipment offsite for further processing.

A range of alternatives was evaluated for inclusion in the GEIS. As defined in the GEIS, the proposed federal action is NRC's determination to grant an application to obtain, renew, or amend a source material license for an ISL facility. Under the no-action alternative, NRC would deny the applicant's or licensee's request. As a result, the new license applicant may choose to resubmit the application to use an alternate uranium recovery method or decide to obtain the yellowcake from other sources. A licensee whose renewal application is denied would have to commence shutting down operations in a timely manner. Denials of license amendments would require the licensee to continue operating under its previously approved license conditions.

Alternative methods for milling uranium were considered as possible alternatives to the ISL process. As stated previously, not all uranium deposits are suitable for ISL extraction. For example, if the uranium mineralization is above the saturated zone (i.e., all of the pore spaces in the ore-bearing rock are not filled with water), ISL techniques may not be appropriate. Likewise, if the ore is not located in a porous and permeable rock unit, it will not be accessible to the leach solution used in the ISL process. Because ISL techniques may not be appropriate in these circumstances, conventional mining (underground or open-pit/surface mining) and milling techniques (conventional milling and heap leaching) are viable alternative technologies.

Inasmuch as the suitability and practicality of using alternative milling methodologies depends on site-specific conditions, a generic discussion of alternative milling methodologies is not appropriate. Accordingly, this GEIS does not contain a detailed analysis of alternative milling methodologies. A detailed analysis of alternative milling methodologies that can be applied at a specific site will be addressed in NRC's site-specific environmental review for individual ISL license applications.

ANALYTICAL APPROACH

The GEIS serves to increase efficiency and eliminate repetitive discussions in NRC's environmental review process by identifying and evaluating environmental impacts that are generic and common to ISL uranium recovery facilities. Information from the GEIS can be summarized and incorporated by reference into the subsequent site-specific environmental review documents. The GEIS also identifies resource areas that need site-specific information to more fully determine the environmental impact to particular resource areas. The site-specific environmental impact analysis also will include any new or significant information necessary to evaluate the ISL facility license application.

For the GEIS, NRC identified the potential environmental impacts associated with the ISL process and the resource areas that could be affected. The general methodology for doing so was to (1) describe the ISL process activity or activities that could affect the resource, (2) identify the resource(s) that can be affected, (3) evaluate past licensing actions and associated environmental review documents and other available information, (4) assess the nature and magnitude of the potential environmental impacts to the resource(s), (5) characterize the significance of the potential impacts, and (6) identify site conditions and mitigation measures that may affect the significance. For some types of impacts analyses (e.g., cumulative impacts, environmental justice evaluations), NRC recognized the difficulty in making determinations in the GEIS, given the location-specific nature of these analyses. For these categories, NRC collected information and conducted initial evaluations, which are documented in the GEIS. The purpose of this information gathering and initial evaluation is intended to provide background data and guidance for the site-specific analyses for these types of impact evaluations.

NRC developed this GEIS based on its experience in licensing and regulating ISL facilities gained during the past 30 years. In the GEIS, NRC does not consider specific facilities, but rather provides an assessment of potential environmental impacts associated with ISL facilities that might be located

in four regions of the western United States. These regions are used as a framework for discussions in this GEIS and were identified based on several considerations, including

- Past and existing uranium milling sites are located within States where NRC has regulatory authority over uranium recovery;
- Potential new sites are identified based on NRC's understanding of where the uranium recovery industry has plans to develop uranium deposits using ISL technology; and
- Locations of previously identified uranium deposits within portions of Wyoming, Nebraska, South Dakota, and New Mexico.

Using these criteria, four geographic regions were identified (Figure ES-1). For the purpose of this GEIS, these regions are

- Wyoming West Uranium Milling Region
- Wyoming East Uranium Milling Region
- Nebraska-South Dakota-Wyoming Uranium Milling Region
- Northwestern New Mexico Uranium Milling Region

The foundation of the environmental impact assessment in the GEIS is based on (1) the historical operations of NRC-licensed ISL facilities and (2) the affected environment in each of the four regions. The structure of the GEIS is presented in Figure ES–2.

Chapter 2 of the GEIS provides a description of the ISL process, addressing construction, operation, aquifer restoration, and decommissioning of an ISL facility. This section also discusses financial assurance, whereby the licensee or applicant establishes a bond or other financial mechanism prior to operations to ensure that sufficient funds are available to complete aquifer restoration, decommissioning, and reclamation activities.

Chapter 3 of the GEIS describes the affected environment in each uranium milling region using the environmental resource areas and topics identified through public scoping comments on the GEIS and from NRC guidance to its staff in NUREG–1748, "Environmental Review Guidance for Licensing Actions Associated With NMSS Programs," issued in 2003.

Chapter 4 of the GEIS provides an evaluation of the potential environmental impacts of constructing, operating, aquifer restoration, and decommissioning at an ISL facility in each of the four uranium milling regions. In essence, this involves placing an ISL facility with the characteristics described in Chapter 2 of the GEIS within each of the four regional areas described in Chapter 3 and describing and evaluating the potential impacts in each region separately. The potential environmental impacts are evaluated for the different stages in the ISL process: construction, operation, aquifer restoration, and decommissioning. Impacts are examined for the resource areas identified in the description of the affected environment. These resource areas are

- Land use
- Transportation
- Geology and soils
- Water resources
- Ecology
- Air quality

- Noise
- Historical and cultural resources
- Visual and scenic resources
- Socioeconomic
- Public and occupational health




Figure ES-1. Location of Four Geographic Regions Used as a Framework for the Analyses Presented in This GEIS




Figure ES-2. Structure of This GEIS

NRC identified a number of other issues that helped in the evaluation of the potential environmental impacts of an ISL facility. These issues include

- Applicable Statutes, Regulations, and Agencies. Various statutes, regulations, and implementing agencies at the federal, state, tribal, and local levels that have a role in regulating ISL facilities are identified and discussed.
- **Waste Management**. Potential impacts from the generation, handling, treatment, and final disposal of chemical, radiological, and municipal wastes are addressed.
- Accidents. Potential accident conditions are assessed in the GEIS. These include consideration of a range of possible accidents and estimation of their consequences, including well field leaks and spills, excursions, processing chemical spills, and ion-exchange resin and yellowcake transportation accidents.
- Environmental Justice. Although not required for a GEIS, to facilitate subsequent
 site-specific analyses, this GEIS provides a first order definition of minority and low
 income populations. Early consultations will be initiated with some of these populations,
 and the potential for disproportionately high and adverse impacts from future ISL
 licensing in the uranium milling regions will be evaluated in the event ISL license
 applications are submitted.
- Cumulative Impacts. The GEIS addresses cumulative impacts from proposed ISL facility construction, operation, groundwater restoration, and decommissioning on all aspects of the affected environment, by identifying past, present, and reasonably foreseeable future actions in the uranium milling regions.
- Monitoring. The GEIS discusses various monitoring methodologies and techniques used to detect and mitigate the spread of radiological and nonradiological contaminants beyond ISL facility boundaries.

SIGNIFICANCE LEVELS

In the GEIS, NRC has categorized the potential environmental impacts using significance levels. According to the Council on Environmental Quality, the significance of impacts is determined by examining both context and intensity (40 CFR 1508.27). Context is related to the affected region, the affected interests, and the locality, while intensity refers to the severity of the impact, which is based on a number of considerations. In this GEIS, the NRC used the significance levels identified in NUREG–1748:

- SMALL Impact: The environmental effects are not detectable or are so minor that
 they will neither destabilize nor noticeably alter any important attribute of the
 resource considered.
- MODERATE Impact: The environmental effects are sufficient to alter noticeably, but not destabilize, important attributes of the resource considered.
- LARGE Impact: The environmental effects are clearly noticeable and are sufficient to destabilize important attributes of the resource considered.

SUMMARY OF IMPACTS

Chapter 4 of the GEIS provides NRC's evaluation of the potential environmental impacts of the construction, operation, aquifer restoration, and decommissioning at an ISL facility in each of the four uranium milling regions. A summary of this evaluation by environmental resource area and phase of the ISL facility lifecycle is provided next.

Land Use Impacts

CONSTRUCTION—Land use impacts could occur from land disturbances (including alterations of ecological cultural or historic resources) and access restrictions (including limitations on other mineral extraction activities, grazing activities, or recreational activities). The potential for land use conflicts could increase in areas with higher percentages of private land ownership and Native American land ownership or in areas with a complex patchwork of land ownership. Land disturbances during construction would be temporary and limited to small areas within permitted boundaries. Well sites, staging areas, and trenches would be reseeded and restored. Unpaved access roads would remain in use until decommissioning. Competing access to mineral rights could be either delayed for the duration of the ISL project or be intermixed with ISL operations (e.g., oil and gas exploration). Changes to land use access including grazing restrictions and impacts on recreational activities would be limited due to the small size of restricted areas, temporary nature of restrictions, and availability of other land for these activities. Ecological, historical, and cultural resources could be affected, but would be protected by careful planning and surveying to help identify resources and avoid or mitigate impacts. For all land use aspects except ecological, historical, and cultural resources, the potential impacts would be SMALL. Due to the potential for unidentified resources to be altered or destroyed during excavation. drilling, and grading, the potential impacts to ecological, historical, or cultural resources would be SMALL to LARGE, depending on local conditions.

OPERATION—The types of land use impacts for operational activities would be similar to construction impacts regarding access restrictions because the infrastructure would be in place. Additional land disturbances would not occur from conducting operational activities. Because access restriction and land disturbance related impacts would be similar to, or less than, those for construction, the overall potential impacts to land use from operational activities would be SMALL.

AQUIFER RESTORATION—Due to the use of the same infrastructure, land use impacts would be similar to operations during aquifer restoration, although some operational activities would diminish—SMALL.

DECOMMISSIONING—Land use impacts would be similar to those described for construction with a temporary increase in land-disturbing activities for dismantling, removing, and disposing of facilities, equipment, and excavated contaminated soils. Reclamation of land to preexisting conditions and uses would help mitigate potential impacts—SMALL to MODERATE during decommissioning, and SMALL once decommissioning is completed.

Transportation Impacts

CONSTRUCTION—Low magnitude traffic generated by ISL construction relative to local traffic counts would not significantly increase traffic or accidents on many of the roads in the region. Existing low traffic roads could be moderately impacted by the additional worker commuting traffic during periods of peak employment. This impact would be expected to be more pronounced in areas with relatively lower traffic counts. Moderate dust, noise, and incidental

wildlife or livestock kill impacts would be possible on, or near, site access roads (dust in particular for unpaved access roads)—SMALL to MODERATE.

OPERATION—Low magnitude traffic relative to local traffic counts on most roads would not significantly increase traffic or accidents. Existing low traffic roads could be moderately impacted by commuting traffic during periods of peak employment including dust, noise, and possible incidental wildlife or livestock kill impacts on or near site access roads. High consequences would be possible for a severe accident involving transportation of hazardous chemicals in a populated area. However, the probability of such accidents occurring would be low owing to the small number of shipments, comprehensive regulatory controls, and use of best management practices. For radioactive material shipments (yellowcake product, ion-exchange resins, waste materials), compliance with transportation regulations would limit radiological risk for normal operations. Low radiological risk is estimated for accident conditions. Emergency response protocols would help mitigate long-term consequences of severe accidents involving release of uranium—SMALL to MODERATE.

AQUIFER RESTORATION—The magnitude of transportation activities would be lower than for construction and operations, with the exception of workforce commuting, which could have moderate impacts on, or in the vicinity of, existing low traffic roads—SMALL to MODERATE.

DECOMMISSIONING—The types of transportation activities, and therefore the types of impacts, would be similar to those discussed for construction and operations, except the magnitude of transportation activities (e.g., number and types of waste and supply shipments, no yellowcake shipments) from decommissioning could be lower than for operations. Accident risks would be bounded by the operations yellowcake transportation risk estimates—SMALL.

Geology and Soils Impacts

CONSTRUCTION—Disturbance to soil would occur from construction (clearing, excavation, drilling, trenching, road construction); however, such disturbances would be expected to be temporary, disturbed areas would be small (approximately 15 percent of the total site area), and potential impacts would be mitigated by using best management practices. A large portion of the well fields, trenches, and access roads would be restored and reseeded after construction. Excavated soils would be stockpiled, seeded, and stored onsite until needed for reclamation fill. No impacts to subsurface geological strata would be likely—SMALL.

OPERATION—Temporary contamination or alteration of soils would be likely from operational leaks and spills and possible from transportation, use of evaporation ponds, or land application of treated waste water. However, detection and response to leaks and spills (e.g., soil cleanup), monitoring of treated waste water, and eventual survey and decommissioning of all potentially impacted soils would limit the magnitude of overall impacts to soils—SMALL.

AQUIFER RESTORATION—Impacts to geology and soils from aquifer restoration activities would be similar to impacts from operations due to use of the same infrastructure and similar activities conducted (e.g., well field operation, transfer activities, liquid effluent treatment and disposal)—SMALL.

DECOMMISSIONING—Impacts to geology and soils from decommissioning would be similar to impacts from construction. Activities to clean up, recontour, and reclaim disturbed lands during decommissioning would mitigate long-term impacts to soils—SMALL.

Surface Water Impacts

CONSTRUCTION—Impacts to surface waters and related habitats from construction (road crossings, filling, erosion, runoff, spills or leaks of fuels and lubricants for construction equipment) would be mitigated through proper planning, design, construction methods, and best management practices. Some impacts directly related to the construction activities would be temporary and limited to the duration of the construction period. U.S. Army Corps of Engineers permits may be required when filling and crossing of wetlands. Temporary changes to spring and stream flow from grading and changes in topography and natural drainage patterns could be mitigated or restored after the construction phase. Impacts from incidental spills of drilling fluids into local streams could occur, but would be temporary due to the use of mitigation measures. Impacts from roads, parking areas, and buildings on recharge to shallow aquifers would be SMALL, owing to the limited area of impervious surfaces proposed. Impacts from infiltration of drilling fluids into the local aquifer would be localized, small, and temporary—SMALL to MODERATE depending on site-specific characteristics.

OPERATION—Through permitting processes, federal and state agencies regulate the discharge of storm water runoff and the discharge of process water. Impacts from these discharges would be mitigated as licensees would operate within the conditions of their permits. Expansion of facilities or pipelines during operations would generate impacts similar to construction—SMALL to MODERATE depending on site-specific characteristics.

AQUIFER RESTORATION—Impacts from aquifer restoration would be similar to impacts from operations due to use of the same (in-place) infrastructure and similar activities conducted (e.g., well field operation, transfer of fluids, water treatment, storm water runoff)—SMALL to MODERATE depending on site-specific characteristics.

DECOMMISSIONING—Impacts from decommissioning would be similar to impacts from construction. Activities to clean up, recontour, and reclaim disturbed lands during decommissioning would mitigate long-term impacts to surface waters—SMALL to MODERATE depending on site-specific characteristics.

Groundwater Impacts

CONSTRUCTION—Water use impacts would be limited by the small volumes of groundwater used for routine activities such as dust suppression, mixing cements, and drilling support over short and intermittent periods. Contamination of groundwater from construction activities would be mitigated by best management practices—SMALL.

OPERATION—Potential impacts to shallow aquifers can occur from leaks or spills from surface facilities and equipment. Shallow aquifers are important sources of drinking water in some areas of the four uranium milling regions. Potential impacts to the ore-bearing and surrounding aquifers include consumptive water use and degradation of water quality (from normal production activities, off-normal excursion events, and deep well injection disposal practices). Consumptive use impacts from withdrawal of groundwater would occur because approximately 1 to 3 percent of pumped groundwater is not returned to the aquifer (e.g., process bleed). That amount of water lost could be reduced substantially by available treatment methods (e.g., reverse osmosis, brine concentration). Effects of water withdrawal on groundwater would be expected to be SMALL as the ore zone normally occurs in a confined aquifer. Estimated drawdown effects vary depending on site conditions and water treatment technology applied. Excursions of lixiviant and mobilized chemical constituents could occur from failure of well seals or other operational conditions that result in incomplete recovery of lixiviant. Well-seal-related

excursions would be detected by the groundwater monitoring system, and periodic well mechanical integrity testing, and impacts would be expected to be mitigated during operation or aquifer restoration. Other excursions could result in plumes of mobilized uranium and heavy metals extending beyond the mineralization zone. The magnitude of potential impacts from vertical excursions would vary depending on site-specific conditions. To reduce the likelihood and consequences of potential excursions at ISL facilities, NRC requires licensees to take preventative measures prior to starting operations, including well tests, monitoring, and development of procedures that include excursion response measures and reporting requirements. Impacts from the alterations of ore body aquifer chemistry would be SMALL, because the aquifer would (1) be confined, (2) not be a potential drinking water source, and (3) be expected to be restored during the restoration period. Potential environmental impacts to confined deep aquifers below the production aquifers from deep well injection of processing wastes would be addressed by the underground injection permitting process regulated by the states and NRC's approval process—SMALL to LARGE, depending on site-specific conditions.

AQUIFER RESTORATION—Potential impacts would be from consumptive use and potential deep disposal of brine slurries after reverse osmosis, if applicable. The volume of water removed from the aquifer and related impacts would be dependent on site-specific conditions and the type of water treatment technology the facility uses. In some cases, groundwater consumptive use for the aquifer restoration has been reported to be less than groundwater use during the ISL operation, and drawdowns due to aquifer restorations have been smaller than drawdown caused by ISL operations. Potential environmental impacts associated with water consumption during aquifer restorations are determined by (1) the restoration techniques chosen, (2) the volume of water to be used, (3) the severity and extent of the contamination, and (4) the current and future use of the production and surrounding aquifers near the ISL facility or at the regional scale—SMALL to MODERATE, depending on site-specific conditions.

DECOMMISSIONING—Potential impacts from decommissioning would be similar to construction (water use, spills) with an additional potential to mobilize contaminants during demolition and cleanup activities. Contamination of groundwater from decommissioning activities would be mitigated by implementation of an NRC-approved decommissioning plan and use of best management practices—SMALL.

Terrestrial Ecology Impacts

CONSTRUCTION—Potential terrestrial ecology impacts would include the removal of vegetation from the well fields and the milling site, the modification of existing vegetative communities, the loss of sensitive plants and habitats from clearing and grading, and the potential spread of invasive species and noxious weed populations. These impacts would be expected to be temporary because restoration and reseeding occur rapidly after the end of construction. Introduction of invasive species and noxious weeds would be mitigated by restoration and reseeding after construction. Shrub and tree removal and loss would take longer to restore. Construction noise could affect reproductive success of sage-grouse leks by interfering with mating calls. Temporary displacement of some animal species would also occur. Critical wintering and year-long ranges are important to survival of both big game and sage-grouse. Raptors breeding onsite may be impacted by construction activities or milling operations, depending on the time of year construction occurs. Wildlife habitat fragmentation, temporary displacement of animal species, and direct or indirect mortalities would be possible. Implementation of wildlife surveys and mitigation measures following established guidelines would limit impacts. The magnitude of impacts depends on whether a new facility is being licensed or an existing facility is being extended—SMALL to MODERATE, depending on site-specific habitat conditions.

OPERATION—Habitats could be altered by operations (fencing, traffic, noise), and individual takes could occur due to conflicts between species habitat and operations. Access to crucial wintering habitat and water could be limited by fencing. However, the State of Wyoming Game and Fish Department specifies fencing construction techniques to minimize impediments to big game movement. Migratory birds could be affected by exposure to constituents in evaporation ponds, but perimeter fencing and netting would limit impacts. Temporary contamination or alteration of soils would be likely from operational leaks and spills and possible from transportation or land application of treated waste water. However, detection and response to leaks and spills (e.g., soil cleanup) and eventual survey and decommissioning of all potentially impacted soil limit the magnitude of overall impacts to terrestrial ecology. Mitigation measures such as perimeter fencing, netting, alternative sites, and periodic wildlife surveys would reduce overall impacts—SMALL.

AQUIFER RESTORATION—Impacts include habitat disruption, but existing (in-place) infrastructure would be used during aquifer restoration, with little additional ground disturbance. Migratory birds could be affected by exposure to constituents in evaporation ponds, but perimeter fencing and netting would limit impacts. Contamination of soils could result from leaks and spills and land application of treated waste water. However, detection and response techniques, and eventual survey and decommissioning of all potentially impacted soils, would limit the magnitude of overall impacts to terrestrial ecology. Mitigation measures such as perimeter fencing, netting, and alternative sites would reduce overall impacts—SMALL.

DECOMMISSIONING—During decommissioning and reclamation, there would be a temporary disturbance to land (e.g., excavated soils, buried piping, removal of structures). However, revegetation and recontouring would restore habitat altered during construction and operations. Wildlife would be temporarily displaced, but are expected to return after decommissioning and reclamation are completed and vegetation and habitat are reestablished—SMALL to MODERATE, depending on site-specific conditions.

Aquatic Ecology Impacts

CONSTRUCTION—Clearing and grading activities associated with construction could result in a temporary increase in sediment load in local streams, but aquatic species would recover quickly as sediment load decreases. Clearing of riparian vegetation could affect light and thus the temperature of water. Construction impacts to wetlands would be identified and managed through U.S. Army Corps of Engineers permits, as appropriate. Construction impacts to surface waters and aquatic species would be temporary and mitigated by best management practices—SMALL.

OPERATION—Impacts could result from spills or releases into surface water. Impacts would be minimized by spill prevention, identification, and response programs, and National Pollutant Discharge Elimination System (NPDES) permit requirements—SMALL.

AQUIFER RESTORATION—Activities would use existing (in-place) infrastructure, and impacts could result from spills or releases of untreated groundwater. Impacts would be minimized by spill prevention, identification, and response programs, and NPDES permit requirements—SMALL.

DECOMMISSIONING—Decommissioning and reclamation activities could result in temporary increases in sediment load in local streams, but aquatic species would recover quickly as

sediment load decreases. With completion of decommissioning, revegetation, and recontouring, habitat would be reestablished and impacts would, therefore, be limited—SMALL.

Threatened and Endangered Species Impacts

CONSTRUCTION—Numerous threatened and endangered species and state species of concern are located in the four uranium milling regions. Small fragmentation of habitats would occur, but most species readapt quickly. The magnitude of impact would depend on the size of a new facility or extension to an existing facility and the amount of land disturbance. Inventory of threatened or endangered species would be developed during site-specific reviews to identify unique or special habitats, and Endangered Species Act consultations conducted with the U.S. Fish and Wildlife Service would assist in reducing impacts—SMALL to LARGE—depending on site-specific habitat and presence of threatened or endangered species.

OPERATION—Impacts could result from individual takes due to conflicts with operations. Small fragmentation of habitats would occur, but most species readapt quickly. The magnitude of impact would depend on the size of a new facility or extension to an existing facility and the amount of land disturbance. Impacts could potentially result from spills or permitted effluents, but would be minimized through the use of spill prevention measures, identification and response programs, and NPDES permit requirements. Inventory of threatened or endangered species developed during site-specific reviews would identify unique or special habitats, and Endangered Species Act consultations conducted with the U.S. Fish and Wildlife Service would assist in reducing impacts—SMALL to LARGE—depending on site-specific habitat and presence of threatened or endangered species.

AQUIFER RESTORATION—Impacts could result from individual takes due to conflicts with aquifer restoration activities (equipment, traffic). Existing (in-place) infrastructure would be used during aquifer restoration, so additional land-disturbing activities and habitat fragmentation would not be anticipated. Impacts may result from spills or releases of treated or untreated groundwater, but impacts would be minimized through the use of spill prevention measures, identification and response programs, and NPDES permit requirements. Inventory of threatened or endangered species would be developed during site-specific reviews to identify unique or special habitats, and Endangered Species Act consultations with the U.S. Fish and Wildlife Service would assist in reducing impacts—SMALL.

DECOMMISSIONING—Impacts resulting from individual takes would occur due to conflicts with decommissioning activities (equipment, traffic). Temporary land disturbance would occur as structures are demolished and removed and the ground surface is recontoured. Inventory of threatened or endangered species developed during site-specific environmental review of the decommissioning plan would identify unique or special habitats, and Endangered Species Act consultations with the U.S. Fish and Wildlife Service would assist in reducing impacts. With completion of decommissioning, re-vegetation, and re-contouring, habitat would be reestablished and impacts would, therefore, be limited—SMALL to LARGE.

Air Quality Impacts

CONSTRUCTION—Fugitive dust and combustion (vehicle and diesel equipment) emissions during land-disturbing activities associated with construction would be small, short-term, and reduced through best management practices (e.g., dust suppression). For example, estimated fugitive dust emissions during ISL construction are less than 2 percent of the National Ambient Air Quality Standards (NAAQS) for PM_{2.5} and less than 1 percent for PM₁₀. For NAAQS attainment areas, nonradiological air quality impacts would be SMALL. A Prevention of

Significant Deterioration Class I area exists in only one of the four regions (Wind Cave National Park in the Nebraska-South Dakota-Wyoming Region). More stringent air quality standards would apply to a facility that impacts the air quality of that area. If impacts were initially assessed at a higher significance level, permit requirements would impose conditions or mitigation measures to reduce impacts—SMALL.

OPERATION—Radiological impacts can result from dust releases from drying of lixiviant pipeline spills, radon releases from well system relief valves, resin transfer or elution, and gaseous/particulate emissions from yellowcake dryers. Only small amounts of low dose materials would be expected to be released based on operational controls and rapid response to spills. Required spill prevention, control, and response procedures would be used to minimize impacts from spills. HEPA filters and vacuum dryer designs reduce particulate emissions from operations, and ventilation reduces radon buildup during operations. Compliance with the NRC-required radiation monitoring program would ensure releases are within regulatory limits. Other potential nonradiological emissions during operations include fugitive dust and fuel from equipment, maintenance, transport trucks, and other vehicles. For NAAQS attainment areas, nonradiological air quality impacts would be SMALL. A Prevention of Significant Deterioration Class I area is located in the Nebraska-South Dakota-Wyoming Region (Wind Cave National Park). More stringent air quality standards would apply to a facility that impacts the air quality of that area. If impacts were initially assessed at a higher significance level, permit requirements would impose conditions or mitigation measures to reduce impacts—SMALL.

AQUIFER RESTORATION—Because the same infrastructure is used, air quality impacts are expected to be similar to, or less than, those during operations. For NAAQS attainment areas, nonradiological air quality impacts would be SMALL. Where a Prevention of Significant Deterioration Class I area exists, such as the Wind Cave National Park in the Nebraska-South Dakota-Wyoming Region, more stringent air quality standards would apply to a facility that impacts the air quality. If impacts were initially assessed at a higher significance level, permit requirements would impose conditions or mitigation measures to reduce impacts—SMALL.

DECOMMISSIONING—Fugitive dust, vehicle, and diesel emissions during land-disturbing activities associated with decommissioning would be similar to, or less than, those associated with construction, would be short-term, and would be reduced through best management practices (e.g., dust suppression). Potential impacts would decrease as decommissioning and reclamation of disturbed areas are completed. For NAAQS attainment areas, nonradiological air quality impacts would be SMALL. However, where a Prevention of Significant Deterioration Class I area exists (Wind Cave National Park in the Nebraska-South Dakota-Wyoming Region), more stringent air quality standards would apply to a facility that impacts the air quality of that area. If impacts were initially assessed at a higher significance level, permit requirements would impose conditions or mitigation measures to reduce impacts—SMALL.

Noise Impacts

CONSTRUCTION—Noise generated during construction would be noticeable in proximity to operating equipment, but would be temporary (typically daytime only). Administrative and engineering controls would be used to maintain noise levels in work areas below Occupational Health and Safety Administration (OSHA) regulatory limits and mitigated by use of personal hearing protection. Traffic noise during construction (commuting workers, truck shipments to and from the facility, and construction equipment such as trucks, bulldozers, and compressors) would be localized, and limited to highways in the vicinity of the site, access roads within the site, and roads in the well fields. Relative increases in traffic levels would be SMALL for the

larger roads, but may be MODERATE for lightly traveled rural roads through smaller communities. Noise may also adversely affect wildlife habitat and reproductive success in the immediate vicinity of construction activities. Noise levels decrease with distance, and at distances more than about 300 m [1,000 ft], ambient noise levels would return to background. Wildlife avoid construction areas because of noise and human activity. Generally, the uranium districts are located more than 300 m [1,000 ft] from the closest community. As a result, noise impacts would be SMALL to MODERATE.

OPERATION—Noise-generating activities in the central uranium processing facility would be indoors, reducing offsite sound levels. Well field equipment (e.g., pumps, compressors) would be contained within structures (e.g., header houses, satellite facilities), also reducing sound levels to offsite receptors. Administrative and engineering controls would be used to maintain noise levels in work areas below OSHA regulatory limits and mitigated by use of personal hearing protection. Traffic noise from commuting workers, truck shipments to and from the facility, and facility equipment would be expected to be localized, limited to highways in the vicinity of the site, access roads within the site, and roads in well fields. Relative increases in traffic levels would be SMALL for the larger roads, but may be MODERATE for lightly traveled rural roads through smaller communities. Most noise would be generated indoors and mitigated by regulatory compliance and best management practices. Noise from trucks and other vehicles is typically of short duration. Also, noise usually is not discernable to offsite receptors at distances of more than 300 m [1,000 ft.] Generally, the uranium districts are located more than 300 m [1,000 ft.] from the closest community—SMALL to MODERATE.

AQUIFER RESTORATION—Noise generation is expected to be less than during construction and operations. Pumps and other well field equipment contained in buildings reduce sound levels to offsite receptors. Existing operational infrastructure would be used, and traffic levels would be expected to be less than those during construction and operations. There are additional sensitive areas that should be considered within some of the regions, but because of decreasing noise levels with distance, aquifer restoration activities would have only SMALL and temporary noise impacts for residences, communities, or sensitive areas, especially those located more than about 300 m [1,000 ft] from specific noise-generating activities. Noise usually is not discernable to offsite receptors at distances more than 300 m [1,000 ft]. Generally, the uranium districts are located more than 300 m [1,000 ft] from the closest community—SMALL to MODERATE.

DECOMMISSIONING—Noise generated during decommissioning would be noticeable only in proximity to equipment and temporary (typically daytime only). Administrative and engineering controls would be used to maintain noise levels in work areas below OSHA regulatory limits and mitigated by use of personal hearing protection. Noise levels during decommissioning would be less than during construction and would diminish as less and less equipment is used and truck traffic is reduced. Noise usually is not discernable to offsite receptors at distances more than 300 m [1,000 ft]. Generally, the uranium districts are located more than 300 m [1,000 ft] from the closest community—SMALL to MODERATE.

Historical and Cultural Resources Impacts

CONSTRUCTION—Potential impacts during ISL facility construction could include loss of, or damage and temporary restrictions on access to, historical, cultural, and archaeological resources. The eligibility evaluation of cultural resources for listing in the National Register of Historic Places (NRHP) under criteria in 36 CFR 60.4(a)–(d) and/or as Traditional Cultural Properties (TCP) would be conducted as part of the site-specific review and NRC licensing procedures undertaken during the NEPA review process. The evaluation of impacts to any

historic properties designated as TCPs and tribal consultations regarding cultural resources and TCPs also occurs during the site-specific licensing application and review process. To determine whether significant cultural resources would be avoided or mitigated, consultations with State Historic Preservation Offices (SHPO), other government agencies (e.g., U.S. Fish and Wildlife Service and State Environmental Departments), and Native American Tribes (the THPO) occur as part of the site-specific review. Additionally, as needed, the NRC license applicant would be required, under conditions in its NRC license, to adhere to procedures regarding the discovery of previously undocumented cultural resources during initial construction. These procedures typically require the licensee to stop work and to notify the appropriate federal, tribal, and state agencies with regard to mitigation measures—SMALL or MODERATE to LARGE depending on site-specific conditions.

OPERATION—Because less land disturbance occurs during the operations phase, potential impacts to historical, cultural, and archaeological resources would be less than during construction. Conditions in the NRC license requiring adherence to procedures regarding the discovery of previously undocumented cultural resources would apply during operation. These procedures typically require the licensee to stop work and to notify the appropriate federal, tribal, and state agencies with regard to mitigation measures—SMALL, depending on site-specific conditions.

AQUIFER RESTORATION—Because less land disturbance occurs during the aquifer restoration phase, potential impacts to historical, cultural, and archaeological resources would be less than those during construction. Conditions in the NRC license requiring adherence to procedures regarding the discovery of previously undocumented cultural resources would apply during aquifer restoration. These procedures typically require the licensee to stop work and to notify the appropriate federal, tribal, and state agencies with regard to mitigation measures—SMALL, depending on site-specific conditions.

DECOMMISSIONING—Because less land disturbance occurs during the decommissioning phase and because decommissioning and reclamation activities would be focused on previously disturbed areas, potential impacts to historical, cultural, and archaeological resources would be less than during construction. Conditions in the NRC license requiring adherence to procedures regarding the discovery of previously undocumented cultural resources would apply during decommissioning and reclamation. These procedures typically require the licensee to stop work and to notify the appropriate federal, tribal, and state agencies with regard to mitigation measures—SMALL, depending on site-specific conditions.

Visual and Scenic Impacts

CONSTRUCTION—Visual impacts result from equipment (drill rig masts, cranes), dust/diesel emissions from construction equipment, and hillside and roadside cuts. Most of the four uranium milling regions are classified as Visual Resource Management (VRM) Class II through IV by the U.S. Bureau of Land Management. A number of VRM Class II areas surround national monuments (EI Morro and EI Malpais), the Chaco Culture National Historic Park, and sensitive areas managed within the Mount Taylor district in the Northwestern New Mexico Uranium Milling District and would have the greatest potential for impacts to visual resources. Most of these areas, however, are located away from potential ISL facilities at distances greater than 16 km [10 mi]. Most potential facilities are located in VRM Class III and IV areas. The general visual and scenic impacts associated with ISL facility construction would be temporary and SMALL, but from a Native American perspective, any construction activities would likely result in adverse impacts to the landscape, particularly for facilities located in areas within view of tribal lands and areas of special significance such as Mount Taylor. As previously discussed,

a Prevention of Significant Deterioration Class I area (Wind Cave National Park) is located in the Nebraska-South Dakota-Wyoming Uranium Milling Region. Prevention of Significant Deterioration Class I areas require more stringent air quality standards that can affect visual impacts. Nevertheless, most potential visual impacts during construction would be temporary as equipment is moved and would be mitigated by best management practices (e.g., dust suppression). Because these sites are in sparsely populated areas and there is generally rolling topography of the region, most visual impacts during construction would not be visible from more than about 1 km [0.6 mi]. The visual impacts associated with ISL construction would be consistent with the predominant VRM Class III and IV—SMALL.

OPERATION—Visual impacts during operations would be less than those associated with construction. Most of the well field surface infrastructure has a low profile, and most piping and cables would be buried. The tallest structures include the central uranium processing facility {10 m [30 ft]} and power lines {6 m [20 ft]}. Because these sites are in sparsely populated areas and there is generally rolling topography of the regions, most visual impacts during operations would not be visible from more than about 1 km [0.6 mi]. Irregular layout of well field surface structures such as wellhead protection and header houses would further reduce visual contrast. Best management practices, and design (e.g., painting buildings) and landscaping techniques would be used to mitigate potential visual impact. The uranium districts in the four regions are all located more than 16 km [10 mi] from the closest VRM Class II region, and the visual impacts associated with ISL construction would be consistent with the predominant VRM Class III and IV—SMALL.

AQUIFER RESTORATION—Aquifer restoration activities would use in-place infrastructure. As a result, potential visual impacts would be the same as, or less than, those during operations—SMALL.

DECOMMISSIONING—Because similar equipment would be used and activities conducted, potential visual impacts during decommissioning would be the same as, or less than, those during construction. Most potential visual impacts during decommissioning would be temporary as equipment is moved and would be mitigated by best management practices (e.g., dust suppression). Visual impacts would be low, because these sites are in sparsely populated areas, and impacts would diminish as decommissioning activities decrease. An approved site reclamation plan is required prior to license termination, with the goal of returning the landscape to preconstruction conditions (predominantly VRM Class III and IV). Some roadside cuts and hill slope modifications, however, may persist beyond decommissioning and reclamation—SMALL.

Socioeconomic Impacts

CONSTRUCTION—Potential impacts to socioeconomics would result predominantly from employment at an ISL facility and demands on the existing public and social services, tourism/recreation, housing, infrastructure (schools, utilities), and the local work force. Total peak employment would be about 200 people, including company employees and local contractors, depending on timing of construction with other stages of the ISL lifecycle. During construction of surface facilities and well fields, the general practice would be to use local contractors (drillers, construction), as available. A local multiplier of 0.7 (U.S. Bureau of the Census) is used to indicate how many ancillary jobs could be created (in this case about 140). For example, local building materials and building supplies would be used to the extent practical. Most employees would live in larger communities with access to more services. Some construction employees, however, would commute from outside the county to the ISL facility, and skilled employees (e.g., engineers, accountants, managers) would come from outside the

local work force. Some of these employees would temporarily relocate to the project area and contribute to the local economy through purchasing goods and services and taxes. Because of the small relative size of the ISL workforce, net impacts would be SMALL to MODERATE.

OPERATION—Employment levels for ISL facility operations would be less than those for construction, with total peak employment depending on timing and overlap with other stages of the ISL lifecycle. Use of local contract workers and local building materials would diminish, because drilling and facility construction would diminish. Revenues would be generated from federal, state, and local taxes on the facility and the uranium produced. Employment types would be similar to construction, but the socioeconomic impacts would be less due to fewer employees—SMALL to MODERATE.

AQUIFER RESTORATION—In-place infrastructure would be used for aquifer restoration, and employment levels would be similar to those for operations—SMALL to MODERATE.

DECOMMISSIONING—A skill set similar to the construction workforce would be involved in dismantling surface structures, removing pumps, plugging and abandoning wells, and reclaiming/recontouring the ground surface. Employment levels and use of local contractor support during decommissioning would be similar to those required for construction. Employment would be temporary, however, as decommissioning activities are short in duration. Because of similar employment levels, other socioeconomic impacts would be similar to construction—SMALL to MODERATE.

Public and Occupational Health and Safety Impacts

CONSTRUCTION—Worker safety would be addressed by standard construction safety practices. Fugitive dust would result from construction activities and vehicle traffic, but would likely be of short duration and would not result in a radiological dose. Diesel emissions would also be of short duration and readily dispersed into the atmosphere—SMALL to MODERATE.

OPERATION—Potential occupational radiological impacts from normal operations would result from (1) exposure to radon gas from the well field, (2) ion-exchange resin transfer operations, and (3) venting during processing activities. Workers would also be exposed to airborne uranium particulates from dryer operations and maintenance activities. Potential public exposures to radiation could occur from the same radon releases and uranium particulate releases (i.e., from facilities without vacuum dryer technology). Both worker and public radiological exposures are addressed in NRC regulations at 10 CFR Part 20, which require licensees to implement an NRC-approved radiation protection program. (Measured and calculated doses for workers and the public are commonly only a fraction of regulated limits.) Nonradiological worker safety matters are addressed through commonly applied occupational health and safety regulations and practices. Radiological accident risks could involve processing equipment failures leading to yellowcake slurry spills, or radon gas or uranium particulate releases. Consequences of accidents to workers and the public are generally low, with the exception of a dryer explosion which could result in worker dose above NRC limits. The likelihood of such an accident would be low, and therefore the risk would also be low. Potential nonradiological accidents impacts include high consequence chemical release events (e.g., ammonia) for both workers and nearby populations. The likelihood, however, of such release events would be low based on historical operating experience at NRC-licensed facilities. primarily due to operators following commonly applied chemical safety and handling protocols— SMALL to MODERATE.

AQUIFER RESTORATION—Activities during aquifer restoration overlap with similar activities during operations (e.g., operation of well fields, waste water treatment and disposal). The resultant impacts on public and occupational health and safety would be bound by operational impacts. The reduction of some operational activities (e.g., yellowcake production and drying, remote ion exchange) will limit the relative magnitude of potential worker and public health and safety hazards—SMALL.

DECOMMISSIONING—Worker and public health and safety would be addressed in a NRC-required decommissioning plan. This plan details how a 10 CFR Part 20 compliant radiation safety program would be implemented during decommissioning, how ensuring the safety of workers and the public would be maintained, and how applicable safety regulations would be complied with—SMALL.

Waste Management Impacts

CONSTRUCTION—Relatively small-scale construction activities (Section 2.3) and incremental well field development at ISL facilities would generate low volumes of construction waste—SMALL.

OPERATION—Operational wastes primarily result from liquid waste streams including process bleed, flushing of depleted eluant to limit impurities, resin transfer wash, filter washing, uranium precipitation process wastes (brine), and plant wash down water. State permit actions, NRC license conditions, and NRC inspections ensure the proper practices would be used to comply with safety requirements to protect workers and the public. Waste treatments such as reverse osmosis and radium settling would be used to segregate wastes and minimize disposal volumes. Potential impacts from surface discharge and deep well injection would be limited by the conditions specified in the applicable state permit. NRC regulations address constructing, operating, and monitoring for leakage of evaporation ponds used to store and reduce volumes of liquid wastes. Potential impacts from land application of treated wastewater would be addressed by NRC review of site-specific conditions prior to approval and routine monitoring in decommissioning surveys. Offsite waste disposal impacts would be SMALL for radioactive wastes as a result of required preoperational disposal agreements. Impacts for hazardous and municipal waste would also be SMALL due to the volume of wastes generated. For remote areas with limited available disposal capacity, such wastes may need to be shipped greater distances to facilities that have capacity; however, the volume of wastes generated and magnitude of such shipments are estimated to be low—SMALL.

AQUIFER RESTORATION—Waste management activities during aquifer restoration would use the same treatment and disposal options implemented for operations. Therefore, impacts associated with aquifer restoration would be similar to operational impacts. While the amount of wastewater generated during aquifer restoration would be dependent on site-specific conditions, the potential exists for additional wastewater volume and associated treatment wastes during the restoration period. However, this would be offset to some degree by the reduction in production capacity from the removal of a well field. NRC review of future ISL facility applications would verify that sufficient water treatment and disposal capacity (and the associated agreement for disposal of byproduct material) are addressed. As a result, waste management impacts from aquifer restoration would be SMALL.

DECOMMISSIONING—Radioactive wastes from decommissioning ISL facilities (including contaminated excavated soil, evaporation pond bottoms, process equipment) would be disposed of as byproduct material at an NRC-licensed facility. A preoperational agreement with a licensed disposal facility to accept radioactive wastes ensures sufficient disposal capacity

would be available for byproduct wastes generated by decommissioning activities. Safe handling, storage, and disposal of decommissioning wastes would be addressed in a required decommissioning plan for NRC review prior to starting decommissioning activities. Such a plan would detail how a 10 CFR Part 20 compliant radiation safety program would be implemented during decommissioning to ensure the safety of workers and the public and compliance with applicable safety regulations. Overall, volumes of decommissioning radioactive, chemical, and solid wastes would be SMALL.

ABBREVIATIONS/ACRONYMS

BLM U.S. Bureau of Land Management CBSA Core-Based Statistical Area

CEA Cumulative Effects Assessment

CERCLA Comprehensive Environmental Response, Compensation, and Liability Act

of 1980

CEQ Council on Environmental Quality
EIS Environmental Impact Statement
EPA U.S. Environmental Protection Agency

FONSI Finding of No Significant Impact

GEIS Generic Environmental Impact Statement

ISL In-situ Leach

MIT Mechanical Integrity Testing

NAAQS National Ambient Air Quality Standards

NAGPRA Native American Graves Protection and Repatriation Act

NDEQ Nebraska Department of Environmental Quality

NEPA National Environmental Policy Act NHPA National Historic Preservation Act

NPDES National Pollutant Discharge Elimination System

NRC U.S. Nuclear Regulatory Commission NRCS Natural Resources Conservation Service NRHP National Register of Historic Places

PVC Polyvinyl Chloride

RFFA Reasonably Foreseeable Future Action SHPO State Historic Preservation Officer

TDS Total Dissolved Solids

THPO Tribal Historic Preservation Officer

UCL Upper Control Limit

UIC Underground Injection Control

UMTRCA Uranium Mill Tailings Radiation Control Act

USACE U.S. Army Corps of Engineers USDA U.S. Department of Agriculture

USFS U.S. Forest Service

VRM Visual Resource Management

WDEQ Wyoming Department of Environmental Quality

SI* (MODERN METRIC) CONVERSION FACTORS

Approximate Conversions From SI Units					
Symbol	When You Know	Multiply By	To Find	Symbol	
		Length			
mm	millimeters	0.039	inches	in	
m	meters	3.28	feet	ft	
m	meters	1.09	yards	yd	
km	kilometers	0.621	miles	mi	
		Area			
mm ²	square millimeters	0.0016	square inches	in ²	
m²	square meters	10.764	square feet	ft ²	
m²	square meters	1.195	square yards	yd ²	
ha	hectares	2.47	acres	ac	
km²	square kilometers	0.386	square miles	mi ²	
	Volume				
mL	milliliters	0.034	fluid ounces	fl oz	
L	liters	0.264	gallons	gal	
m ³	cubic meters	35.314	cubic feet	ft ³	
m ³	cubic meters	1.307	cubic yards	yd ³	
m ³	cubic meters	0.0008107	acre-feet	acre-feet	
		Mass			
g	grams	0.035	ounces	OZ	
kg	kilograms	2.202	pounds	lb	
Mg (or "t")	megagrams (or "metric ton")	1.103	short tons (2000 lb)	Т	
	Temperatu	ure (Exact Degre	es)		
°C	Celsius	1.8 °C + 32	Fahrenheit	°F	

^{*}SI is the symbol for the International System of Units. Appropriate rounding should be performed to comply with Section 4 of ASTM E380 (ASTM International. "Standard for Metric Practice Guide." West Conshohocken, Pennsylvania: ASTM International. Revised 2003.).

5 CUMULATIVE EFFECTS

5.1 Introduction

The Council on Environmental Quality (CEQ) National Environmental Policy Act (NEPA) regulations, as amended (40 CFR Parts 1500–1508) define cumulative effects as "... the impact on the environment that results from the incremental impact of the action when added to other past, present, and reasonably foreseeable future actions regardless of what agency (Federal or non-Federal) or person undertakes such other actions. Cumulative impacts can result from individually minor but collectively significant actions taking place over a period of time." This chapter describes past, present, and reasonably foreseeable future actions (RFFAs) in the uranium milling regions and evaluates which resource areas would be potentially impacted by both *in-situ* leaching (ISL) facilities and the types of RFFAs identified in the regions. Due to the complex and site-specific nature of a cumulative impact assessment, this chapter provides useful information for understanding the potential for cumulative impacts when licensing future ISL facilities in the milling regions, but does not make conclusions regarding cumulative impacts that could be applied to specific sites.

A National Research Council study on hardrock mining on federal lands recognized the cumulative effects could become a concern due to past, current, and future activities in the vicinity of the mine under consideration. Specifically, cumulative impacts were defined as the collective impacts of several operations involving human activities, including mining, grazing, farming, timbering, water diversion or discharge, and industrial processing; they also include future impacts not immediately observable (Committee on Hardrock Mining on Federal Lands, 1999, p. 242). While this definition does not precisely match the definition in the CEQ's NEPA regulations, it does include the concept that a variety of other past, present, and future actions in the vicinity of the proposed project could cumulatively contribute to the effects on specific resources resulting from the proposed project subjected to NEPA analyses.

The study also noted that there were many uncertainties related to the cumulative effects of mineral production, including technologies such as the *in-situ* leaching (ISL) process for uranium recovery. As a result, several research needs were articulated. Examples include the need for methodologies (or models) for predicting cumulative effects from mineral recovery activities under different environmental circumstances, the need for collaborative approaches for resolving multiple and conflicting demands on common resources, and the need for the design of a long-term monitoring program and strategies that can be used to identify impact contributions from various actions, as well as the resource sustainability (Committee on Hardrock Mining on Federal Lands, 1999).

When the many activities potentially associated with an ISL project (e.g., several satellite well fields, solution-water injection wells, and associated extraction wells are drilled; extracted fluids are processed at remote locations; pipelines are built to transport liquid from these locations to a central processing plant; selected wastewaters are disposed of using deep wells; and yellowcake is shipped by truck) are considered, they could cause impacts to specific local and regional resources. In addition, ISL projects could involve relicensing or expanding existing facilities and operations, possibly with the use of new designs for new well fields or modifications in existing designs. These new or relicensed projects could be located within or near geographical areas that have been subject to uranium recovery via conventional mining and milling, oil and gas exploration and production, and other energy developments such as

coal-bed methane projects. For all of these reasons, cumulative effects assessment is an important part of the licensing process for ISL projects.

Establishing the appropriate "scope" of the cumulative effects portion of an impact study is a fundamental feature of planning and conducting such a study for an ISL project. The CEQ NEPA regulations in 40 CFR Parts 1500–1508 indicate that "scope consists of the range of actions ..." to be considered in a NEPA compliance document. CEQ regulations in 40 CFR 1508.25 identify the following three types of actions for consideration, which all pertain to ISL projects:

- Connected actions are closely related and should be discussed in the same environmental impact statement (EIS), supplemental EIS, or environmental assessment.
 The multiple activities of an ISL project illustrate connected actions. Such actions are interdependent parts of a larger action (the overall ISL project) and depend on the larger action for their justification.
- Cumulative actions, when viewed with other proposed actions, have cumulatively significant impacts and should therefore be discussed in the same NEPA compliance document. Cumulative actions could include future planned expansion of the proposed ISL facility, proposals for other new ISL projects in the same geographic areas, and relicensing of nearby existing ISL projects.
- Similar actions, when viewed with other reasonably foreseeable or proposed agency
 actions, have similarities that provide a way to evaluate their environmental
 consequences together, such as common timing, or geography or impacts on common
 resources. Similar actions could include other local or regional energy or industrial
 development projects, or land usage activities, which could impact the same resources
 the proposed ISL project hopes to change.

In 1997, the CEQ published guidance on an approach to consider cumulative effects within the NEPA compliance process (CEQ, 1997) as described in Appendix F. This guidance contains an 11-step process, integrated within the traditional NEPA (or environmental impact assessment) process. Steps 1–4 relate to scoping (including the establishment of the scope), Steps 5–7 to describe the affected environment, and Steps 8–11 to determine the environmental consequences. These 11 steps can be applied at a general study planning level and at a detailed level for specific resources, ecosystems, and human communities, which are impacted by the original proposed action. For uranium recovery, the original action could be associated with a license application for a new ISL facility or with a relicensing action for an existing facility.

The resource areas addressed in this generic EIS (GEIS) include land use, transportation, geology and soils, surface water, groundwater, wetlands, terrestrial ecology, aquatic ecology, threatened or endangered species, air quality, noise, historical and cultural resources, visual and scenic resources, socioeconomic conditions, public health and safety, occupational health and safety, waste management, and environmental justice.

Cumulative impacts (effects) was one of the topical areas addressed in three public scoping meetings related to this GEIS (see Appendix A). In addition, impacts from ISL facilities on groundwater and surface water, ecology, historic and cultural resources, and environmental justice were also noted. Such impacts could occur from direct and indirect effects from ISL facilities, as well as cumulative effects from these facilities and other past, present, and RFFAs within the four defined geographic uranium milling regions.

5.2 Other Past, Present, and Reasonably Foreseeable Future Actions in the Four Regions

This section includes summary information on historical, current, and anticipated uranium recovery sites. In addition, other current and potential projects in the regions are illustrated by current draft and final EISs within the regions. Information sources for the regions are then included. Finally, "actions matrices" for each of the regions are included.

5.2.1 Uranium Recovery Sites

Table 5.2-1 includes tabulations of the history and also the short-term future of uranium recovery sites in the states of Wyoming, South Dakota, Nebraska, and New Mexico based on indications from industry to NRC (NRC, 2009). A total of 62 sites are included, with the sites subdivided into three types (ISL facilities, conventional uranium milling, and heap leach facilities). A total of nine ISI research and development sites are listed. Additionally, several other ISL research and development sites were associated with basic information gathering on the ISL process for a particular site that was later used to support approval for a license for commercial production.

Twenty-four of the sites involve conventional milling. Many of these sites are either in active decommissioning or have already been decommissioned. The U.S. Department of Energy (DOE) is the long-term custodian for the decommissioned sites under a general license issued by NRC. It should be noted that in the table, under the Wyoming West Uranium Milling Region, there are two entries for the Sweetwater site name. One entry is for an existing conventional uranium mill site, while the other lists both a potential conventional mine and a potential ISL facility.

In addition, there are abandoned convention uranium mining sites from the past that exist in the four uranium milling regions. For example, from 1944 to 1986, nearly 4 million tons of uranium ore was extracted in New Mexico under lease agreements with the Navajo National (EPA, 2008). This has resulted in over 500 abandoned uranium mines and associated environmental contamination in that area alone (EPA, 2008). Evaluating the potential impacts from past mining activities on new ISL proposals is a site-specific analysis that, if applicable to a proposed site, would be evaluated by applicants during site characterization and by the NRC staff when a site-specific licensing review is conducted.

A total of 31 past, present, and potential future sites are in Wyoming and associated with the ISL process (including the Sweetwater site, which lists both the ISL process and a conventional mine). Out of these 31 ISL sites, 21 sites are in the Wyoming East Uranium Milling Region, 9 sites are in the Wyoming West Uranium Milling Region, and 1 site is in the Nebraska-South Dakota-Wyoming Uranium Milling Region. Five additional ISL sites are or potentially may be located in the Nebraska-South Dakota-Wyoming Uranium Milling Region, and one research and development site and one licensed ISL site are in the Northwestern New Mexico Uranium Milling Region. The table also shows four potential conventional milling sites (three in the Northwestern New Mexico Uranium Milling Region and the Sweetwater site in the Wyoming West Uranium Milling Region) and one potential heap leach site (in the Northwestern New Mexico Uranium Milling Region).

To reflect present actions and RFFAs related to uranium recovery in the four uranium milling regions analyzed in the GEIS, certain of the sites are identified as "potential sites" under Status

Table 5.2-1. Past, Existing, and Potential Uranium Recovery Sites in Wyoming, South Dakota, Nebraska, and New Mexico*				
Site Name	Company/Owner	Type††	County, State	Status‡
	Wyoming V	Vest Urani	um Milling Region	
Sky	Strathmore Minerals Corp.	ISL	Fremont, WY	Potential site
Jab & Antelope	Uranium One	ISL ^{2,3}	Fremont, WY	Potential site license application under review by the U.S. Nuclear Regulatory Commission (NRC)
Lucky Mc	Pathfinder Mines Corp.	Conv.	Fremont, WY	Decommissioning
Split Rock	Western Nuclear, Inc.	Conv.	Fremont, WY	Decommissioning
Bison Basin	Ogle Petroleum	ISL ³	Fremont, WY	License terminated
Riverton	U.S. Department of Energy (DOE)	Conv.	Fremont, WY	UMTRCA Title I processing site
Gas Hills	Power Resources Inc.	ISL ²	Natrona & Fremont, WY	Licensed—on standby
Gas Hills	Strathmore Minerals Corp.	Conv.	Natrona & Fremont, WY	Potential site
Gas Hills	Umetco Minerals Corp.	Conv.	Natrona & Fremont, WY	Decommissioning
ANC	American Nuclear Corp.	Conv.	Natrona, WY	Decommissioning
Nine Mile Lake	Rocky Mountain Energy Co.	ISL ¹	Natrona, WY	License terminated
Lost Soldier	UR-Energy Corp.	ISL	Sweetwater, WY	Potential site
Sweetwater	Wildhorse Energy	ISL & Conv.	Sweetwater, WY	Potential site
West Alkali Creek	Wildhorse Energy	ISL	Sweetwater, WY	Potential site
Lost Creek	UR-Energy Corp.	ISL ³	Sweetwater, WY	Potential site—license application under review by NRC
Sweetwater	Kennecott Uranium Co.	Conv.	Sweetwater, WY	Licensed—on standby
	Wyoming E		um Milling Region	
Reno Creek 2	International Uranium Corp.	ISL ³	Campbell, WY	Not licensed—applicant withdraws
Ruby Ranch	Conoco	ISL ¹	Campbell, WY	Not licensed—applicant withdraws
Ruby Ranch	Power Resources Inc.	ISL	Campbell, WY	Potential site
Reno Creek	Strathmore Minerals Corp.	ISL	Campbell, WY	Potential site
Nichols Ranch & Hank	Uranerz Energy Corp.	ISL ^{2,3}	Campbell & Johnson, WY	Potential site—license application under review by NRC

Table 5.2-1. Past, Existing, and Potential Uranium Recovery Sites in Wyoming, South Dakota, Nebraska, and New Mexico* (continued)				
Site Name	Company/Owner	Type††	County, State	Status‡
	Wyoming East U	ranium Mil	ling Region (conti	nued)
Moore Ranch	Uranium One	ISL ³	Campbell, WY	Potential site—license application under review by NRC
North Butte & Ruth	Power Resources Inc.	ISL ^{2,3}	Campbell, WY	Licensed—on standby
Reno Creek 1	Rocky Mountain Energy Co.	ISL ¹	Campbell, WY	License terminated
Collins Draw	Cleveland Cliffs Iron Co.	ISL ¹	Campbell, WY	License terminated
Shirley Basin South	DOE	Conv.	Carbon, WY	UMTRCA Title II disposal site
Peterson Ranch	Arizona Public Service Co. Malapai Resources	ISL ¹	Converse, WY	Not pursued
Ludeman	Uranium One	ISL	Converse, WY	Potential site
Highland 1	Exxon Minerals	ISL ³	Converse, WY	Licensed but not pursued
Reynolds Ranch	Power Resources Inc.	ISL ²	Converse, WY	Licensed but not operational
Highland 2	Everest Minerals	ISL ³	Converse, WY	Licensed—later combined with Smith Ranch facility license
Smith Ranch - Highland	Power Resources Inc.	ISL ³	Converse, WY	Operating
Bear Creek	Bear Creek Uranium Co.	Conv.	Converse, WY	Decommissioning
Highlands	Exxon Mobil Corp.	Conv.	Converse, WY	Decommissioning
Leuenberger	Teton Exploration Drilling	ISL ^{1,3}	Converse, WY	License terminated
South Powder River Basin	Kerr-McGee	ISL ¹	Converse, WY	License terminated with approval of Smith Ranch license
Spook	Department of Energy	Conv.	Converse, WY	UMTRCA Title I disposal site
Allemand- Ross	Uranium One	ISL	Johnson, WY	Potential site
Irigaray/ Christensen Ranch	Cogema Malapai Resources	ISL ^{2,3}	Johnson, WY	Licensed for operations
Willow Creek	J&P Corp. Western Nuclear	ISL ¹	Johnson, WY	License terminated with approval of Irigaray license
Shirley Basin	Pathfinder Mines Corp.	Conv.	Natrona, WY	Decommissioning

Table 5.2-1. Past, Existing, and Potential Uranium Recovery Sites in Wyoming, South Dakota, Nebraska, and New Mexico* (continued)				
Site Name	Company/Owner	Type††	County, State	Status‡
	Wyoming East U	ranium Mil	ling Region (conti	nued)
North Platte	Uranium Resources	ISL ¹	Platte, WY	License terminated
	Nebraska-South Dak	ota-Wyom	ing Uranium Millin	g Region
Marsland	Cameco (Crow Butte Resources)	ISL ²	Dawes, NE	Potential site
Three Crow	Cameco (Crow Butte Resources)	ISL ²	Dawes, NE	Potential site
North Trend	Cameco (Crow Butte Resources)	ISL ²	Dawes, NE	Potential site—license application under review by NRC
Crow Butte	Cameco (Crow Butte Resources)	ISL ³	Dawes, NE	Operating
Dewey Burdock	Powertech Uranium Corp.	ISL ³	Fall River, SD	Potential site—license application submitted to NRC
Edgemont	DOE	Conv.	Fall River, SD	UMTRCA Title II disposal site
Dewey	Powertech Uranium	ISL ²	Niobrara, WY	Potential site
Terrace	Corp.			
0 1 5:1	Northwestern New Mexico Uranium Milling Region			
Grants Ridge	Uranium Energy Corp.	Heap Leach	Cibola, NM	Potential site
Homestake	Homestake Mining Co.	Conv.	Cibola, NM	Decommissioning
Bluewater	DOE	Conv.	Cibola, NM	UMTRCA Title II disposal site
L-Bar	DOE	Conv.	Cibola, NM	UMTRCA Title II disposal site
Marquez	Neutron Energy	Conv.	McKinley, NM	Potential site
Mt. Taylor	Rio Grande Resources	Conv.	McKinley, NM	Potential site
Roca Honda	Strathmore Minerals Corp.	Conv.	McKinley, NM	Potential site
Crownpoint	Hydro Resources, Inc.	ISL ³	McKinley, NM	Licensed but not operational
Ambrosia Lake	Rio Algom	Conv.	McKinley, NM	Decommissioning
Churchrock	United Nuclear Corp.	Conv.	McKinley, NM	Decommissioning
Section 9 Ambrosia	Mobil Corp. DOE	ISL ¹ Conv.	McKinley, NM McKinley, NM	License terminated UMTRCA Title I disposal
Lake				site

Table 5.2-1. Past, Existing, and Potential Uranium Recovery Sites in Wyoming, South Dakota, Nebraska, and New Mexico* (continued)				
Site Name	Company/Owner Lynett County State Statust			
	Northwestern New Mexico Uranium Milling Region (continued)			
Shiprock	DOE	Conv.	San Juan, NM	UMTRCA Title I disposal site

*Information on potential future uranium recovery applications is based on indication from industry summarized in NRC. "Expected New Uranium Recovery Facility Applications/Restarts/Expansions: Updated 3/11/2009" http://www.nrc.gov/info-finder/materials/uranium/2008-ur-projects-list-public.pdf (07 April 2009).

- 1 = Research and Development/Pilot
- 2 = Satellite
- 3 = Commercial scale

Conv. = Conventional uranium mill

‡Status: Uranium Mill Tailings Radiation Control Act (UMTRCA) Title I and Title II sites are uranium mill processing or tailings sites that have been decommissioned. The U.S. Department of Energy is the long-term custodian of these sites.

column, consistent with either license applications received by NRC or formal letters of intent to submit license applications sent to NRC by the identified company/owner (NRC, 2008).

5.2.2 EISs as Indicators of Present and RFFAs

One indicator of present and RFFAs in the four uranium milling regions is the number of draft and final EISs prepared by federal agencies within a recent time period. The informational database which was queried is the U.S. Environmental Protection Agency (EPA) EIS Database at http://yosemite.epa.gov/oeca/ webeis.nsf/viEIS01?OpenView>. The time period selected for the review was the 38-month period from January 7, 2005, through February 22, 2008. A total of 10 draft and 22 final EISs were identified for specific projects and counties within the four regional areas. In addition, three draft programmatic and seven final programmatic EISs were identified for large-scale actions primarily related to several states, including Wyoming, Nebraska, and South Dakota. Tables 5.2-2 through 5.2-6 include lists of the specific project-related EISs for the four regional areas. The EISs can be obtained via Internet searching and utilized in site-specific cumulative effects assessments for proposed ISL facilities.

For the Wyoming West Uranium Milling Region, Table 5.2-2 includes four draft EISs and seven final EISs. Four projects are related to gas developments, two are associated with natural gas pipelines, and one involves coal mining. These seven projects could contribute to both local and regional cumulative impacts on air quality, land usage, terrestrial plants and animals, and groundwater and surface water resources. The extent of such contributions depends on the locations of these projects in relation to other past actions and RFFAs, including ISL facilities for uranium recovery. The remaining three projects listed in Table 5.5-2 involve resource management actions which are focused on reducing historical impacts from grazing practices, improving resource conditions by planning and management, and/or minimizing continuing practices with adverse impacts.

Table 5.2-2. Draft and Final Environmental Impact Statements (EISs) Related to the Wyoming West Uranium Milling Region (in Chronological Order From January 2005 to February 2008)		
Date	Statement	
February 4, 2005	U.S. Forest Service, Final EIS, Upper Green River Area Rangeland Project, Proposed Site-Specific Grazing Management Practices, Bridger-Teton Forest, Sublette, Teton and Fremont Counties, WY (resource management)	
July 8, 2005	Federal Energy Regulatory Commission, Final EIS, Entrega Pipeline Project, Construction and Operation New Interstate Natural Gas Pipeline System, Right-of-Way Grant Issue by BLM, Meeker Hub and Cheyenne Hub, Rio Blanco and Weld Counties, CO, and Sweetwater County, WY (gas pipeline)	
August 19, 2005	Federal Energy Regulatory Commission, Final EIS, Piceance Basin Expansion Project, Construction and Operation of a New Interstate Natural Gas Pipeline System, Wamsutter Compressor Station to Interconnections and Greasewood Compressor Station, Rio Blanco County, CO, and Sweetwater County, WY (gas pipeline)	
December 2, 2005	Seminoe Road Natural Gas Development Project, Proposed Coal Bed Natural Gas Development and Operation, Carbon County, WY (gas development)	
November 17, 2006	U.S. Bureau of Land Management (BLM), Final EIS, Pit 14 Coal Lease- by-Application Project, Black Butte Coal Mine, Surface Mining Operations, Federal Coal Lease Application WYW160394, Sweetwater County, WY (coal mining)	
December 1, 2006	BLM, Final EIS, Atlantic Rim Natural Gas Field Development Project, Proposed Natural Gas Development to 2000 Wells, 1800 to Coal Beds and 200 to Other Formations, Carbon County, WY (gas development)	
June 8, 2007	BLM, Final EIS, Casper Field Office Planning Area Resource Management Plan, Implementation, Natrona, Converse, Goshen, and Platte Counties, WY (resource management)	
October 12, 2007	BLM, Draft EIS, Moxa Arch Area Infill Gas Development Project, Drill, Extract, Remove, and Market Natural Gas Under Valid Existing Oil and Gas Leases, Approval, Right-of-Way Grants and U.S. Army COE Section 404 Permit(s), Lincoln, Uinta, and Sweetwater Counties, WY (gas development)	
November 1, 2007	Bureau of Indian Affairs, Draft EIS, Riverton Dome Coal Bed Natural Gas and Conventional Gas Development Project, Construction of Well Pads, Roads, Pipelines, and Production Facilities, Wind River Indian Reservation, Fremont County, WY (gas development)	
January 14, 2008	BLM, Final EIS, Rawlins Field Office Planning Area Resource Management Plan, Addresses the Comprehensive Analysis of Alternatives for the Planning and Management of Public Land and Resources Administered by BLM, Albany, Carbon, Laramie, and Sweetwater Counties, WY (resource management)	

	Table 5.2-3 Draft and Final Environmental Impact Statements (EISs) Related to the Wyoming East Uranium Milling Region (in Chronological Order From January 2005 to February 2008)		
Date	Statement		
February 4, 2005	U.S. Forest Service (USFS), Final EIS, Tongue Allotment Management Plan, Proposal To Continue Livestock Grazing on All or Portions of the 22 Allotments, Bighorn National Forest, Tongue and Medicine Wheel/Paintrock Ranger Districts, Johnson, Sheridan, and Bighorn Counties, WY (resource management-grazing)		
April 13, 2007	U.S. Bureau of Land Management (BLM), Final EIS, Maysdorf Coal Lease by Application (LBA) Tract, Federal Coal Application WYW154432, Implementation, Campbell County, WY (coal mining)		
August 17, 2007	USFS, Final EIS, Thunder Basin Analysis Area Vegetation Management, To Implement Best Management Grazing Practices and Activities, Douglas Ranger District, Medicine Bow-Routt National Forests and Thunder Basin National Grassland, Campbell, Converse, and Weston Counties, WY (resource management-grazing)		
August 31, 2007	BLM, Final EIS, Eagle Butte West Coal Lease Application, Issuance of Lease for a Tract of Federal Coal, Wyoming Powder River Basin, Campbell County, WY (coal mining)		
August 31, 2007	Rural Utilities Service, Draft EIS, Dry Fork Station and Hughes Transmission Line, Construct Electric Generating Facilities, Campbell and Sheridan Counties, WY; withdrawn (power plant and transmission line)		
December 21, 2007	USFS, Draft EIS, Thunder Basin National Grassland Prairie Dog Management Strategy, Land and Resource Management Plan Amendment #3, Proposes To Implement a Site-Specific Strategy To Manage Black-Tailed Prairie Dog, Douglas Ranger District, Medicine Bow-Routt National Forest and Thunder Basin National Grassland, Campbell, Converse, Niobrara, and Weston Counties, WY (species management)		
February 2, 2008	BLM, Draft EIS, West Antelope Coal Lease Application Federal Coal Lease Application WYW163340, Implementation, Converse and Campbell Counties, WY (coal mining)		
October 24, 2008	South Gillette Area Coal Lease Applications. Draft EIS, Proposal to Lease Four Tracts of Federal Coal Reserves, Belle Ayr, Coal Creek, Caballo, and Cordero Rojo Mines, Wyoming Power River Basin, Campbell County, WY		

Table 5.2-4. Draft and Final Programmatic or Large-Scale Environmental Impact Statements (EISs) Related to One or Both of the Wyoming Regional Study Areas (in Chronological Order From January 2005 to February 2007)		
Date	Statement	
March 30, 2006	U.S. Bureau of Land Management (BLM), Revised Final EIS, Programmatic—Proposed Revision to Grazing Regulations for the Public Lands, 42 CFR Part 4100, in the Western Portion of the United	
	States (resource management-grazing)	

Statements (EISs)	Table 5.2-4. Draft and Final Programmatic or Large-Scale Environmental Impact Statements (EISs) Related to One or Both of the Wyoming Regional Study Areas (in Chronological Order From January 2005 to February 2007) (continued)		
Date	Statement		
May 26, 2006	Bureau of Reclamation, Final EIS, Programmatic—Platte River Recovery Implementation Program, Assessing Alternatives for the Implementation of a Basinwide, Cooperative, Endangered Species Recovery Program, Four Target Species: Whooping Crane, Interior Least Tern, Piping Plover, and Pallid Sturgeon, NE, WY, and CO (resource management-endangered species recovery)		
August 17, 2006	Federal Railroad Administration, Final EIS, Powder River Basin Expansion Project, Construction of New Rail Facilities, Finance Docket No. 33407 Dakota, Minnesota and Eastern Railroad, SD, WY, and MN (railroad)		
March 22, 2007	Federal Energy Regulatory Commission, Final EIS, Rockies Express Western Phase Project, Construction and Operation for the Natural Gas Pipeline Facilities: Rockies Express (CP06–354–000), TransColorado (CP06–401–000), and Overthrust (CP06–423–000), CO, WY, NE, KS, MO, and NM (gas pipeline)		
June 15, 2007	U.S. Forest Service, Final EIS, Northern Rockies Lynx Management Direction, Selected Alternative F, Conservation and Promote Recovery of the Canada Lynx, NFS and BLM to Amend Land Resource Management Plans for 18 National Forests (NF), MT, WY, UT, and ID (resource management-Canada lynx)		
June 29, 2007	BLM, Final EIS, Programmatic—Vegetation Treatments Using Herbicides on BLM Public Lands in 17 Western States, including Alaska (resource management-herbicides)		
August 24, 2007	BLM, Final EIS, Overland Pass Natural Gas Liquids Pipeline Project (OPP), Construction and Operation of 760-mile Natural Gas Liquids Pipeline, Right-of-Way Grant, KS, WY, and CO (gas pipeline)		
November 16, 2007	U.S. Department of Energy, Draft EIS, PROGRAMMATIC—Designation of Energy Corridors in 11 Western States, Preferred Location of Future Oil, Gas, and Hydrogen Pipelines and Electricity Transmission and Distribution Facilities on Federal Land, AZ, CA, CO, ID, MT, NV, NM, UT, WA, and WY (energy corridors)		
November 30, 2007	Federal Energy Regulatory Commission, Draft EIS, Rockies Express Pipeline Project, (REX-East) Construction and Operation of Natural Gas Pipeline Facilities, WY, NE, MO, IL, IN, and OH (gas pipeline)		
December 21, 2007	BLM, Draft EIS, Programmatic EIS—Oil Shale and Tar Sands Resource Management Plan (RMP) Amendments To Address Land Use Allocations in Colorado, Utah, and Wyoming (oil shale and tar sands)		

Table 5.2-5. Draft and Final Environmental Impact Statements (EISs) Related to the Nebraska-South Dakota-Wyoming Uranium Milling Region (in Chronological Order From January 2005 to February 2007)			
Date	Statement		
June 3, 2005	U.S. Forest Service (USFS), Final EIS, Dean Project Area, Proposes To Implement Multiple Resource Management Actions, Black Hills National Forest, Bearlodge Ranger District, Sundance, Crook County, WY (resource management)		
August 12, 2005	USFS, Final EIS, Black-Tailed Prairie Dog Conservation and Management on the Nebraska National Forest and Associated Units, Implementation, Dawes, Sioux, Blaine, Cherry, Thomas Counties, NE, and Custer, Fall River, Jackson, Pennington, Jones, Lyman, Stanley Counties, SD (resource management-prairie dog)		
October 28, 2005	National Park Service, Draft EIS, Badlands National Park/North Unit General Management Plan, Implementation, Jackson, Pennington, and Shananon Counties, SD (resource management)		
November 20, 2005	USFS, Final EIS, Deerfield Project Area, Proposes To Implement Multiple Resource Management Actions, Mystic Ranger District, Black Hills National Forest, Pennington County, SD (resource management)		
November 25, 2005	USFS, Final EIS, Bugtown Gulch Mountain Pine Beetle and Fuels Projects, To Implement Multiple Resource Management Actions, Black Hills National Forest, Hell Canyon Ranger District, Custer County, SD (resource management)		
January 13, 2006	USFS, Final EIS, Black Hills, National Forest Land and Resource Management Plan Phase II Amendment, Proposal To Amend the 1997 Land and Resource Management Plan, Custer, Fall River, Lawrence, Meade, and Pennington Counties, SD, and Crook and Weston Counties, WY (resource management)		
February 3, 2006	USFS, Final EIS, Black-Tailed Prairie Dog Conservation and Management on the Nebraska National Forest and Associated Units, Implementation, Dawes, Sioux, Blaine, Cherry, Thomas Counties, NE, and Custer, Fall River, Jackson, Pennington, Jones, Lyman, Stanley Counties, SD (resource management-prairie dog)		
May 12, 2006	USFS, Final Supplemental EIS, Dean Project Area, Proposes To Implement Multiple Resource Management Actions, New Information to Disclose Direct, Indirect, and Cumulative Environmental Impacts, Black Hills National Forest, Bearlodge Ranger District, Sundance, Crook County, WY (resource management)		
June 1, 2007	USFS, Final EIS, Norwood Project, Proposes To Implement Multiple Resources Management Actions, Black Hills National Forest, Hell Canyon Ranger District, Pennington County, SD, and Weston and Crook Counties, WY (resource management)		
June 8, 2007	USFS, Draft EIS, Nebraska and South Dakota Black-Tailed Prairie Dog Management, To Manage Prairie Dog Colonies in an Adaptive Fashion, Nebraska National Forest and Associated Units, Including Land and Resource Management Plan Amendment 3, Dawes, Sioux, Blaine Counties, NE, and Custer, Fall River, Jackson, Pennington, Jones, Lyman, Stanley Counties, SD (resource management-prairie dog)		

Table 5.2-5. Draft and Final Environmental Impact Statements (EISs) Related to the Nebraska-South Dakota-Wyoming Uranium Milling Region (in Chronological Order From			
January 2005 to February 2007) (continued)			
Date	Statement		
June 29, 2007	USFS, Final EIS, Mitchell Project Area, To Implement Multiple		
	Resource Management Actions, Mystic Ranger District, Black Hills		
	National Forest, Pennington County, SD (resource management)		
September 14, 2007	USFS, Final EIS, Citadel Project Area, Proposes To Implement		
	Multiple Resource Management Actions, Northern Hills Ranger		
	District, Black Hills National Forest, Lawrence County, SD (resource		
	management)		
February 22, 2008	USFS, Draft EIS, Upper Spring Creek Project, Proposes To Implement		
	Multiple Resource Management Actions, Mystic Ranger District, Black		
	Hills National Forest, Pennington County, SD (resource management)		

Table 5.2-6. Draft and Final Environmental Impact Statements (EISs) Related to the Northwestern New Mexico Uranium Milling Region (in Chronological Order From January 2005 to February 2007)		
Date	Statement	
February 2, 2005	Bureau of Indian Affairs, Final Supplemental EIS, Programmatic— Navajo Nation 10-Year Forest Management Plan, Selected Preferred Alternative Four, Chuska Mountain and Defiance Plateau Area, AZ and NM (forest management)	
April 20, 2007	U.S. BLM, Draft EIS, Socorro Resource Management Plan Revision, Implementation, Socorro and Catron Counties, NM (resource management)	

For the Wyoming East Uranium Milling Region, Table 5.2-3 includes four draft EISs and four final EISs. Three of the projects are related to leases for coal extractions (mining), and one to the development of a power plant and transmission line. However, the draft EIS on the power plant and transmission line was withdrawn. Nonetheless, it was included in Table 5.2-3 because it could be reactivated at a future date. Coal extraction projects can contribute to local and regional cumulative impacts on air quality, land usage, terrestrial plants and animals, and surface and groundwater hydrology and quality. Further, impacts on wetlands, threatened and endangered species, and cultural resources could also occur as a result of specific project locations.

As noted for the Wyoming West Uranium Milling Region, the extent of contributions of these projects to cumulative effects depends on their locations in relation to other past and present actions and RFFAs, including future ISL facilities. Two of the three remaining projects involve better management of grazing practices, while the final one is focused on the management of black-tailed prairie dogs. These latter three projects should result in environmental improvements. Table 5.2-4 includes five listed "programmatic" EISs (two draft EISs and three final EISs) and five regional EISs (one draft EIS and four final EISs). These 10 EISs are characterized by either management actions encompassing large geographical areas or proposed projects extending over large areas. For purposes of this GEIS, all 10 EISs will be considered as programmatic documents, whether or not they are labeled as such. Six of the EISs are related, either directly or indirectly, to energy development projects. Three of the six involve natural gas pipelines encompassing several states (two related to the Rockies Express and one to the Overland Pass project). Of interest herein are segments of the projects related

to Wyoming (the Wyoming West and Wyoming East Uranium Milling Regions) and Nebraska (the Nebraska-South Dakota-Wyoming Uranium Milling Region). The U.S. Department of Energy draft EIS addresses energy corridors involving future oil, gas, and hydrogen pipelines and electricity transmission lines on federal lands in 11 western states, including Wyoming. In general, pipeline projects can have impacts on terrestrial resources within their specified corridors, and on aquatic resources near pipeline crossings of surface streams and rivers. The fifth energy-related project in Table 5.2-4 involves rail facilities associated with the Powder River Basin in Wyoming and South Dakota; regional coal transport could be enhanced by this project. The final energy-related project is associated with land use allocations for oil shale and tar sands development activities. Each of these six programmatic projects should be considered for inclusion, as appropriate, within any cumulative effects analyses of proposed ISL facilities in the Wyoming West and Wyoming East, Uranium Milling Regions. Further, the four resource management actions listed in Table 5.2-4 (grazing regulations, endangered species recovery programs for four listed species, lynx management, and herbicide usage) should also be considered within any cumulative effects studies of proposed ISL facilities in the three regions.

For the Nebraska-South Dakota-Wyoming Uranium Milling Region, a total of three draft EISs and 10 final EISs are identified in Table 5.2-5. All 13 EISs are related to resource management actions in the Black Hills National Forest or associated management units. Multiple actions related general resources management are addressed in 10 of the EISs. The remaining three actions are specifically associated with black-tailed prairie dog conservation and management. The actions in all 13 EISs are focused on improving natural resources conditions and reducing adverse impacts from various man-related activities.

For the Northwestern New Mexico Uranium Milling Region, Table 5.2-6 includes only one draft EIS and one final EIS issued over the study period. Both EISs are related to resource management; hence they are focused on improving natural resources conditions and reducing adverse impacts from various man-related activities.

5.3 Concurrent Actions

5.3.1 Wyoming West Uranium Milling Region

Table 5.3-1 contains a listing of six categories of actions in the State of Wyoming that could impact the resources and topics addressed in Chapters 3 and 4 (see Sections 3.2 and 4.2). The six categories (traditional land uses; wildlife/fisheries/forest management; recreation; government lands and land management; mineral extraction/energy development; and cultural resources preservation) include specific actions which illustrate the respective categories. Step 4 of the CEQ's 11-step cumulative effect process (see Appendix F) indicates that other past, present, and RFFAs that could contribute to cumulative effects on specific resources and topics should be identified. The listed actions in Table 5.3-1 are reflective of both past and continuing actions; further, the majority of the actions are expected to continue into the future. Locational information (by county) is included for several of the listed actions. Where county information is not available, it is assumed that the actions are statewide and applicable in both the Wyoming West and Wyoming East Uranium Milling Regions.

Table 5.3-1 also includes a series of codes to reflect that each listed action can impact certain resources and topics that are known to be impacted the ISL process for uranium recovery. The 12 resources and topics and their designator codes are defined in the footnotes to the table.

Table 5.3-1. Other Actions Concurrent With Uranium Recovery in the Wyoming West Uranium Milling Region*				
Categories of Actions	Impacts on Resource and Topics†			
Traditional Land Uses				
Livestock grazing	LU, WR, E, HC, S			
Agricultural activities	LU, WR, E, HC, S			
Protection of significant alluvial farmland	LU, WR, S			
Irrigation	GS, WR, S			
Development of new or expanded communities	LU, T, GS, WR, E, HC, S, WM			
Roads and highways	LU, T, WR, E, HC, S			
Indian Reservations	LU, WR, E, HC, VS			
Wind River [Northern Arapaho and Eastern Shoshone (Fremont)]				
Wildlife/Fisheries/Forest Management				
Timber harvests (see National Forests)	LU, T, GS, WR, E, N, S			
Wild horse management (Carbon, Sweetwater, Fremont)	LU, E			
Protection of T/E species – critical habitat identification	LU, E			
Riparian habitat preservation/enhancement	LU, WR, E			
Recreation (See Information on National Forests and State Parks for Specific Location of Activities)				
Hunting, fishing, hiking	E			
Camping	LU, E			
Overland vehicle use (OHVs)	LU, GS, WR, E			
Trail riding	LU, GS			
Recreation management plans (Natrona, Converse)	LU, WR, E, HC, VS			
Government Lands and Land Management				
State Parks				
Sinks Canyon and Boysen State Park and Reservoir (Fremont)	LU, WR, E			
Endess K. Wilkins State Park and Independence Rock State Historical Site (Natrona)	LU, E, HC			
Seminoe SP & Reservoir (Carbon)	LU, WR, E			
National Forest/Grasslands • Shoshone National Forest (Fremont)	LU, WR, E, HC, VS			
National Wildlife Areas • Pathfinder National Wildlife Refuge	LU, E, HC, VS			
(Natrona/Carbon)Seedskadee National Wildlife Refuge (Sweetwater)	LU, E, HC, VS			

Table 5.3-1. Other Actions Concurrent With Uranium Recovery in the Wyoming West Uranium Milling Region* (continued)			
Categories of Actions	Impacts on Resource and Topics†		
Mineral Extraction/Energy Development			
Transmission lines/substations (Fremont)	LU, E		
Coal-related actions (Weston, Campbell, Converse, Carbon, Sheridan, Sweetwater) • Power plants • Railroad development for hauling coal; past and present action, throughout coal regions • Coal mines	WR, E, AQ, N, HC, VS, S, WM LU, T, WR, E, N, S LU, GS, WR, E, AQ, N, HC, VS, S, WM		
Mine reclamation (Carbon, Converse, Campbell)	GS, WR, E, AQ		
Natural gas and oil	LIL CS MID E AO NI HO VS S MAA		
 Conventional oil development (Natrona, Sweetwater) Natural gas field development 	LU, GS, WR, E, AQ, N, HC, VS, S, WM LU, GS, WR, E, AQ, HC, S		
(Carbon, Sweetwater)Overland natural gas pipelines and compressor stations (Carbon,	LU, T, WR, E, N, HC, S		
Sweetwater, Natrona, Fremont) Oil shale and tar sands energy development (Fremont,	LU, GS, WR, E, AQ, N, HC, VS, S, WM		
Sweetwater) • CO ₂ -enhanced oil recovery (Natrona, Sweetwater)	LU, GS, WR, E, AQ, N, HC, VS, S, WM		
 Coal bed natural gas/methane development (Campbell, Carbon, Converse, Fremont, Johnson, Sweetwater, Sheridan) 	LU, GS, WR. E. AQ, N. HC, VS, S		
Uranium activities • Permitting of new or inactive ISL	LU, T, GS, WR, E, AQ, N, HC, VS, S, PO, WM		
facilities (Johnson, Campbell, Fremont, Sweetwater) Conventional mining and milling	LU, T, GS, WR, E, AQ, N, HC, VS, S, PO, WM		
 Reclaimed open pit mines (Converse, Carbon, Fremont) 	LU, T, GS, WR, E, AQ, N, HC, VS, S, PO, WM		
Mining of other mineralsTrona (Sweetwater)Sand and gravel	LU, T, GS, WR, E, AQ, N, HC, VS, S, WM		

Table 5.3-1. Other Actions Concurrent With Uranium Milling Re	, , , , , , , , , , , , , , , , , , , ,		
Categories of Actions	Impacts on Resource and Topics†		
Cultural Resources Preservation			
Fort Robinson—Nebraska	LU, HC		
Historic trails—crisscrossing state of Wyoming	LU, HC		
Ghost towns (Fremont)	LU HC		

^{*} The Wyoming West Uranium Milling Region includes the western parts of Natrona and Carbon Counties, the northeastern portion of Sweetwater County, and the eastern portion of Fremont County.

LU = land use

T = transportation

GS = geology and soils

WR = water resources (wetlands, surface water, and groundwater)

E = ecology (terrestrial, aquatic, and threatened/endangered species)

AQ = air quality (non-radiological)

N = noise

HC = historical and cultural resources

VS = visual and scenic resources

S = socioeconomics

PO = public and occupational health and safety

WM = waste management

Further, these resources and topics provide the basic structure used in this GEIS for describing the affected environment (Chapter 3) and addressing the impacts of the four phases of an ISL project (Chapters 4 and 10). When a designator code (e.g., LU for land use) is listed for a specific action within a category, this denotes that the action would be anticipated to cause an impact on the resource or topic.

Table 5.3-2 contains a list of 21 coal mines in Wyoming. This listing and status information was obtained from http://www.wma-minelife.com/coal/coalfrm/coaldat.htm. A total of four surface mines and one underground mine are located in the Wyoming West Uranium Milling Region, with three in Carbon County and two in Sweetwater County. The 2006 production from these mines in the Hanna Coal Field and the Green River Coal Region ranged from about 25,580 to 4,912,960 metric tons [28,200 to 5,414,423 short tons]. Surface mining of coal can cause adverse impacts on land use, geology and soils, water resources, ecology, air quality, noise, historical and cultural resources, visual and scenic resources, socioeconomics, and waste management. The impacts of additional coal-related actions are included in Table 5.3-3.

5.3.2 Wyoming East Uranium Milling Region

Table 5.3-3 contains a listing of six categories of actions in the State of Wyoming that could impact the 12 resources and topics addressed in Chapters 3 and 4 for the Wyoming East Uranium Milling Region (see Section 3.3 and 4.3). The structure of Table 5.3-3 is the same as that for the Wyoming West Uranium Milling Region (Table 5.3-1). Where county information is not available, it is assumed that the actions are statewide and applicable in both the Wyoming West and Wyoming East Uranium Milling Regions. The listed actions in Table 5.3-3 are reflective of both past and continuing actions; further, the majority of the actions are expected to continue into the future.

[†]The resources and topics codes include

Table 5.3-2. Coal Mining Projects as Identified by the Wyoming Mining Association (Data Through 2006)*				
Mine Name	Owner/Operator (If Different)	Location	Mine Type	Production in 2006 (Tons)
Powder River Basin Coal				
Buckskin	Buckskin Mining Co.	Campbell Co.	Surface	22,768,303
Rawhide	Powder River Coal	Campbell Co.	Surface	17,092,993
Dry Fork	Western Fuels of WY	Campbell Co.	Surface	5,860,998
Eagle Butte	Foundation Coal West	Campbell Co.	Surface	25,355,158
KFx	KFx Fuel Partners	Campbell Co.	Surface	87,863 (just recently back in production)
Wyodak	Wyodak Resources Development	Campbell Co.	Surface	4,698,473
Caballo	Powder River	Campbell Co.	Surface	32,700,000
Belle Ayr	Foundation Coal West	Campbell Co.	Surface	24,593,035
Cordero/Rojo	Rio Tinto Energy America	Campbell Co.	Surface	39,747,620
Coal Creek		Campbell Co.		3,097,584 (No production 2000-2005)
Jacobs Ranch	Rio Tinto Energy America	Campbell Co.	Surface	40,000,376
Black Thunder	Thunder Basin Coal	Campbell Co.	Surface	92,517,728
North Antelope/ Rochelle	Powder River Coal	Campbell Co. Converse Co.	Surface	88,527,969
Antelope	Rio Tinto Energy America	Campbell Co. Converse Co.	Surface	33,984,178
Dave Johnston	Glenrock Coal	Converse Co.	Surface	Reclaimed—no production since 2000
Seminoe #2	Arch Coal, Inc.	Carbon Co.	Surface	Final reclamation in 2006
Medicine Bow	Arch Coal, Inc.	Carbon Co. 1 River Coal Regi	Surface	28,212, but 0 in 2005; relatively small operation
Jim Bridger	Bridger Coal	Sweetwater Co.	Surface	5,414,423
Black Butte	Black Butte Coal		Surface	3,410,309
*Wyoming Mining A	Association. "Wyoming Coal In> (16 November 2008).	Sweetwater Co. Data." 2008. 2008. <a <="" href="http://w" td=""><td></td><td></td>		

Table 5.3-3. Other Actions Related to or Conflicting With Uranium Recovery in the
Wyoming East Uranium Milling Region*

Categories of Actions Impacts on Resource and Topics†

Traditional Land Uses

Livestock grazing LU, WR, E, HC, S

Agricultural activities LU, WR, E, HC, S

Protection of significant alluvial farmland LU, WR, S

Irrigation GS, WR, S

Categories of Actions	Milling Region* (continued) Impacts on Resource and Topics†	
_	d Uses (continued)	
Development of new or expanded communities	LU, T, GS, WR, E, HC, S, WM	
Roads and highways	LU, T, WR, E, HC, S	
• •	/Forest Management	
Timber harvests (see National Forests)	LU, T, GS, WR, E, N, S	
Wild horse management (Carbon, Sweetwater,	LU, E	
Fremont)	,	
Protection of T/E species – critical habitat	LU, E	
identification		
Riparian habitat preservation/enhancement	LU, WR, E	
Prairie dog management (Campbell, Converse,	LU, E	
Weston)		
•	Forests and State Parks for Specific Location	
	ctivities)	
Hunting, fishing, hiking	E	
Camping	LU, E	
Overland vehicle use (OHVs)	LU, GS, WR, E	
Trail riding	LU, GS	
Recreation management plans (Natrona, Convers		
	and Land Management	
State Parks	III E HC	
Endess K. Wilkins State Park and Independence Book State Historical Site Independence Book State Historical Site	LU, E, HC	
Independence Rock State Historical Site (Natrona)		
Seminoe SP & Reservoir (Carbon)	LU, WR, E	
National Forest/Grasslands	20, 1111, 2	
Thunder Basin National Grasslands	LU, WR, E, HC, VS	
(Weston, Campbell, Converse)	20, 1111, 2, 110, 10	
 Medicine Bow National Forest (Converse, 	LU, WR, E, HC, VS	
Natrona, Carbon)		
Bighorn National Forest (Johnson)	LU, WR, E, HC, VS	
National Wildlife Areas		
 Pathfinder NWA (Natrona/Carbon) 	LU, E, HC, VS	
Mineral Extraction	/Energy Development	
Transmission lines/substations (Fremont)	LU, E	
Coal-related actions (Weston, Campbell,		
Converse, Carbon, Sheridan, Sweetwater)		
 Power plants 	WR, E, AQ, N, HC, VS, S, WM	
 Railroad development for hauling coal; 	LU, T, WR, E, N, S	
past and present action, throughout coal		
regions	III GS WA E VO VI HO VS S WW	
Coal mines Mine and leasting (Coathon Coanness)	LU, GS, WR, E, AQ, N, HC, VS, S, WM GS, WR, E, AQ	
 Mine reclamation (Carbon, Converse, 	GS, WK, E, AQ	
Campball)		
Campbell) Coal leasing (Campbell, Converse)	LU, S	

Table 5.3-3. Other Actions Related to or Conflicting With Uranium Recovery in the Wyoming East Uranium Milling Region* (continued)			
Categories of Actions Impacts on Resource and Topics†			
	gy Development (continued)		
Natural gas and oil	Jy Development (continued)		
Conventional oil development (Natrona, Sweetwater)	LU, GS, WR, E, AQ, N, HC, VS, S, WM		
Natural gas field development (Carbon, Sweetwater)	LU, GS, WR, E, AQ, HC, S		
Overland natural gas pipelines and compressor stations (Carbon,	LU, T, WR, E, N, HC, S		
Sweetwater, Natrona, Fremont) Oil shale and tar sands energy development (Fremont, Sweetwater)	LU, GS, WR, E, AQ, N, HC, VS, S, WM		
 CO₂-enhanced oil recovery (Natrona, Sweetwater) 	LU, GS, WR, E, AQ, N, HC, VS, S, WM		
 Coal Bed natural gas/methane development (Campbell, Carbon, Converse, Fremont, Johnson, 	LU, GS, WR, E, AQ, N, HC, VS, S		
Sweetwater, Sheridan)			
 Uranium activities Permitting of new or inactive ISL facilities (Johnson, Campbell, Fremont, 	LU, T, GS, WR, E, AQ, N, HC, VS, S, PO, WM		
Sweetwater) Continued operation of ISL facilities	LU, T, GS, WR, E, AQ, N, HC, VS, S, PO, WM		
(Converse) Conventional mining and milling	LU, T, GS, WR, E, AQ, N, HC, VS, S, PO, WM		
Reclaimed open pit mines (Converse, Carbon, Fremont) Mining of other minerals	LU, T, GS, WR, E, AQ, N, HC, VS, S, PO, WM		
 Mining of other minerals Bentonite (Weston, Johnson, Natrona) Sand and Gravel Scoria 	LU, T, GS, WR, E, AQ, N, HC, VS, S, WM		
Cultural Resources Preservation			
Historic trails – crisscrossing state of Wyoming	LU, HC		
Historic mines and other pioneer sites (Converse	e, LU, HC		
County, the southeastern portion of Johnson County, a Nebraska-South Dakota-Wyoming Milling Region inclu			
AQ = air quality (non-radiological) N = noise HC = historical and cultural resources VS = visual and scenic resources S = socioeconomics PO = public and occupational health and safety WM = waste management			

As noted previously, Table 5.3-2 contains a list of coal mines in Wyoming. This listing and status information was obtained from the following Wyoming website at http://www.wma-minelife.com/coal/coalfrm/coaldat.htm. The Wyoming East Uranium Milling Region includes 15 surface mines in the Powder River Basin, with 13 in Campbell County and two in Converse County. The 2006 coal production levels indicated that 14 mines were in operation in the Wyoming East Uranium Milling Region, with annual production levels ranging from 79,700 to about 83,916,000 metric tons [87,900 to 92,500,000 short tons]. Surface mining of coal can cause adverse impacts on land use, geology and soils, water resources, ecology, air quality, noise, historical and cultural resources, visual and scenic resources, socioeconomics, and waste management. The impacts of additional coal-related actions are included in Table 5.3-3.

5.3.3 Nebraska-South Dakota-Wyoming Uranium Milling Region

Table 5.3-4 is structured similarly to Table 5.3-1, with a listing of six categories of actions in the states of Nebraska and South Dakota that could impact the resources and topics addressed in Chapters 3 and 4 (see Sections 3.4 and 4.4). Concurrent actions in Wyoming are described in Tables 5.3-1 and 5.3-3. When the county is not identified for the action, it is assumed that the actions are statewide and applicable in the South Dakota and Nebraska portions of the Nebraska-South Dakota-Wyoming Uranium Milling Region. There are no coal mines identified in the affected counties in this uranium milling region. The listed actions in Table 5.3-4 are

Table 5.3-4. Other Actions Concurrent With Uranium Recovery in the			
Nebraska-South Dakota-Wyoming Uranium Milling Region*			
Categories of Actions	Impacts on Resource and Topics†		
Traditional Land Uses			
Livestock grazing	LU, WR, E, HC, S		
Agricultural activities	LU, WR, E, HC, S		
Protection of significant alluvial farmland	LU, WR, S		
Irrigation	GS, WR, S		
Development of new or expanded	LU, T, GS, WR, E, HC, S, WM		
communities			
Roads and highways	LU, T, WR, E, HC, S		
Indian Reservations			
 Pine Ridge (Oglala Sioux) 	LU, WR, E, HC, VS		
Wildlife/Fisheries/Forest Management			
Timber harvests (see National Forests)	LU, T, GS, WR, E, N, S		
Wild horse management	LU, E		
Protection of T/E species; critical habitat	LU, E		
identification			
Riparian habitat preservation/enhancement	LU, WR, E		
Prairie dog management (Weston, Sioux,	LU, E		
Dawes)			
Wildland fires (Black Hills National Forest; all	LU, T, WR, E, AQ, HC, VS, S		
four counties)			

Table 5.3-4. Other Actions Concur			
Nebraska-South Dakota-Wyoming U			
Categories of Actions	Impacts on Resource and Topics†		
Recreation (See Information on National Forests and State Parks for Specific Location			
of Acti			
Hunting, fishing, hiking	E		
Camping	LU, E		
Overland vehicle use (OHVs)	LU, GS, WR, E		
Trail riding	LU, GS		
Recreation management plans	LU, WR, E, HC, VS		
Scenic byways (Custer, Lawrence, Pennington)	LU, T, WR, E, HC, VS, S		
Black Hills major tourist center (all four	LU, T, WR, E, HC, VS, S		
counties in South Dakota)	20, 1, WK, 2, 110, VO, 0		
Government Lands ar	nd Land Management		
National Forest/Grasslands (Wyoming)	, and the second		
Thunder Basin National Grasslands	LU, WR, E, HC, VS		
(Weston, Campbell, Converse)			
National Parks/Monuments (Wyoming)			
 Devils Tower, Wyoming (Weston) 	LU, WR, E, HC, VS		
State Parks (South Dakota)			
Custer State Park (Custer)	LU, WR, E		
 Angostura State Recreation Area (Fall River) 	LU, WR, E		
National Forest/Grasslands (South Dakota)			
 Black Hills National Forest (Fall River, 	LU, WR, E, HC, VS		
Custer, Pennington, Lawrence)	LIL WD E LIC VC		
Buffalo Gap National Grassland (Fall Biver Guster Bennington)	LU, WR, E, HC, VS		
River, Custer, Pennington)			
National Parks/Monuments (South Dakota) • Mt. Rushmore National Memorial	LU, WR, E, HC, VS		
(western Pennington)	LO, WK, E, FIG, V3		
Jewel Cave National Monument	LU, WR, E, HC, VS		
(Custer)			
Wind Cave National Park (Custer)	LU, WR, E, HC, VS		
State Parks/Recreation Areas (Nebraska)			
Chadron SP (Dawes); within the	LU, WR, E, HC, VS		
Nebraska National Forest			
 Ft. Robinson SP (Sioux, Dawes) 	LU, WR, E, HC, VS		
Box Butte Reservoir State Recreation	LU, WR, E, HC, VS		
Area (Dawes)			

Table 5.3-4. Other Actions Concurrent With Uranium Recovery in the Nebraska-South Dakota-Wyoming Uranium Milling Region* (continued)			
Categories of Actions	Impacts on Resource and Topics†		
Government Lands and Lar	nd Management (continued)		
National Forests/Grasslands			
 Oglala National Grasslands (Sioux, Dawes) 	LU, WR, E, HC, VS		
 Toadstool Geologic Park (Sioux); operated by US Forest Service 	LU, WR, E, HC, VS		
 Nebraska National Forest (Sioux, Dawes) 	LU, WR, E, HC, VS		
 Within the Forest is Soldier Creek Wilderness (Sioux) 	LU, WR, E, HC, VS		
 Wilderness (Glodx) Within the Forest is Pine Ridge National Recreation Area (Dawes) 	LU, WR, E, HC, VS		
National Parks/Monuments			
Agate Fossil Beds National Monument (Sioux)	LU, WR, E, HC, VS		
	nergy Development		
Transmission lines/substations	LU, E		
Coal-related actions			
Power plants	WR, E, AQ, N, HC, VS, S, WM		
 Railroad development for hauling 	LU, T, WR, E, N, S		
coal; past and present action,			
throughout coal regions			
Coal mines	GS, WR, E, AQ		
 Mine reclamation 	LU, GS, WR, E, AQ, N, HC, VS, S, WM		
 Coal leasing 	LU, GS, WR, E, AQ, N, HC, VS, S		
Natural gas and oil			
 Oil and gas leasing (Custer National Forest) 	LU, GS		
 Conventional oil development (Fall River) 	LU, GS, WR, E, AQ, N, HC, VS, S, WM		
 Natural gas field development 	LU, GS, WR, E, AQ, N, HC, S		
 Overland natural gas pipelines and compressor stations 	LU, T, WR, É, Ń, HĆ, Ś		
Uranium activities			
 Permitting of new or inactive ISL facilities (Fall River, Custer, Dawes) 	LU, T, GS, WR, E, AQ, N, HC, VS, S, PO, WM		
Continued operation of ISL facilitiesConventional mining and milling	LU, T, GS, WR, E, AQ, N, HC, VS, S, PO,		
	WM LU, T, GS, WR, E, AQ, N, HC, VS, S, PO, WM		
Other			
Energy corridors‡	LU, T, WR, E, N, HC, S		
 Limestone conveyor system (Custer)§ 	LU, T, E, AQ, N, HC, VS, S		

Table 5.3-4. Other Actions Concurrent With Uranium Recovery in the Nebraska-South Dakota-Wyoming Uranium Milling Region* (continued)			
Categories of Actions Impacts on Resource and Topic			
Cultural Resources Preservation			
Big Thunder historic gold mine (Pennington)	LU, HC		
Several pioneer homesteads in Black Hills	LU, HC		
Museum of the Fur Trade (Dawes)	LU, HC		

*The Nebraska-South Dakota-Wyoming Uranium Milling Region includes all or portions of three Wyoming counties; specifically, this region includes Crook County, the eastern half of Weston County, and the northeastern portion of Niobrara County. In addition, the South Dakota portion of the region includes Fall River, Custer, and Lawrence Counties and the western half of Pennington County. The Nebraska portion of the region includes Sioux, Box Butte, and Dawes Counties in the far northwestern portion of the state.

†The resources and topics codes include

LU = land use

T = transportation

GS = geology and soils

WR = water resources (wetlands, surface water, and groundwater)

E = ecology (terrestrial, aquatic, and threatened/endangered species)

AQ = air quality (non-radiological)

N = noise

HC = historical and cultural resources

VS = visual and scenic resources

S = socioeconomics

PO = public and occupational health and safety

WM = waste management

‡Federal Departments of Agriculture, Commerce, Defense, Energy, and the Interior are proposing to designate corridors on Federal land for locating future oil, natural gas, and hydrogen pipelines and electricity transmission and distribution infrastructure in the West. These corridors would be the agency-preferred locations where pipelines and transmission lines may be sited and built in the future. Such corridors could be proposed for South Dakota.

§This is a proposed 11-km [7-mi] enclosed, aboveground conveyor belt to transfer limestone in Custer County, South Dakota. The project will cross national forest lands, BLM lands, and private lands. The BLM is preparing an EIS on this project.

reflective of both past and continuing actions; further, the majority of the actions are expected to continue into the future.

5.3.4 Northwestern New Mexico Uranium Milling Region

Table 5.3-5 is structured similarly to Table 5.3-1, with a listing of six categories of actions in the State of New Mexico that could impact the resources and topics addressed in Chapters 3 and 4 (see Sections 3.5 and 4.5). The six categories (traditional land uses; wildlife/fisheries/forest management; recreation; government lands and land management; mineral extraction/energy development; and cultural resources preservation) include specific actions which illustrate the respective categories. The listed actions in Table 5.3-5 are reflective of both past and continuing actions; further, the majority of the actions are expected to continue into the future.

5.4 Approaches to Conducting a Site-Specific Cumulative Effects Analysis

Each of the four uranium milling regions analyzed in this GEIS includes existing and previous uranium recovery facilities (Table 5.2-1), as well as anticipated new, modified, or planned restarts of uranium ISL facilities (NRC, 2009). In addition, each region includes a number of

Table 5.3-5. Other Actions Concurrent With Uranium Recovery in the Northwestern New Mexico Uranium Milling Region*			
Categories of Actions	Impacts on Resource and Topics†		
Traditional Land Uses			
Livestock grazing	LU, WR, E, HC, S		
Agricultural activities	LU, WR, E, HC, S		
Protection of significant alluvial farmland	LU, WR, S		
Irrigation	GS, WR, S		
Development of new or expanded	LU, T, GS, WR, E, HC, S, WM		
communities			
Roads and highways	LU, T, WR, E, HC, S		
Indian reservations			
 Navajo (McKinley) 	LU, WR, E, HC, VS		
 Zuni (McKinley, Cibola) 	LU, WR, E, HC, VS		
 Ramah Navajo (Cibola) 	LU, WR, E, HC, VS		
Acoma (Cibola)	LU, WR, E, HC, VS		
Lacuna (Cibola)	LU, WR, E, HC, VS		
Canonito (Cibola)	LU, WR, E, HC, VS		
 Alamo Bend Navajo (Socorro) 	LU, WR, E, HC, VS		
Wildlife/Fisheries/Fo	rest Management		
Timber harvests (see National Forests)	LU, T, GS, WR, E, N, S		
Wild horse management	LU, E		
Protection of T/E species; critical habitat	LU, E		
identification	,		
Riparian habitat preservation/enhancement	LU, WR, E		
Endangered species reintroduction (Aplomado	LU, E		
falcon) (Socorro)			
Recreation (See Information on National	Forests and State Parks for Specific		
Location of A	Activities)		
Hunting, fishing, hiking	E		
Camping	LU, E		
Overland vehicle use (OHVs) (Catron,	LU, GS, WR, E		
Socorro)			
Trail riding	LU, GS		
Recreation management plans	LU, WR, E, HC, VS		
Government Lands and	Land Management		
State Parks			
 Bluewater SP (Cibola) 	LU, WR, E		
 Red Rock SP (McKinley) 	LU, WR, E		
National Forest/Grasslands			
 Cibola National Forest (all four counties) 	LU, WR, E, HC, VS		
 Apache-Sitgreaves National Forest (Catron) 	LU, WR, E, HC, VS		
Gila National Forest (Catron)	LU, WR, E, HC, VS		

Table 5.3-5. Other Actions Concurrent With Uranium Recovery in the Northwestern New Mexico Uranium Milling Region* (continued)			
Categories of Actions	Impacts on Resource and Topics†		
Government Lands and L	and Management (continued)		
National Monuments/Recreation			
areas/Wildlife refuges/Conservation areas			
Gila Cliff Dwelling National	LU, E, HC, VS		
Monument (Catron)			
El Morro National Monument	LU, E, HC, VS		
(Cibola)	LU, E, HC, VS		
 Chain of Craters Wilderness Study Area (Cibola) 	LU, E, HC, VS		
 El Malpais National Conservation 			
Area (surrounds El Malpais National			
Monument, but does not include it;	LU, E, HC, VS		
Cibola)	LO, E, HO, VS		
El Malpais National Monument; lava beds (Cibola)	LU, E, HC, VS		
 Salinas Pueblo Mission National Monument (Socorro) 	LU, E, HC, VS		
 Datil Well NRA (Catron; within the Cibola National Forest) 	LU, E, HC, VS		
Bosque del Apache NWR (Socorro)			
Ft. Wingate Military Reservation (McKinley)	LU, E, HC		
Mineral Extraction	/Energy Development		
Transmission lines/substations	LU, E		
Coal-related actions			
 Power plants (McKinley) 	WR, E, AQ, N, HC, VS, S, WM		
 Coal mines (McKinley, Cibola) 	GS, WR, E, AQ		
Coal leasing	LU, GS, WR, E, AQ, N, HC, VS, S		
Natural gas and oil			
Conventional oil development	LU, GS, WR, E, AQ, N, HC, VS, S, WM		
 Natural gas field development (McKinley) 	LU, GS, WR, E, AQ, HC, S		
Overland natural gas pipelines and	LU, T, WR, E, N, HC, S		
compressor stations			
Uranium activities			
Permitting of new or inactive ISL facilities	LU, T, GS, WR, E, AQ, N, HC, VS, S, PO, WM		
Continued operation of ISL facilities	LU, T, GS, WR, E, AQ, N, HC, VS, S, PO, WM		
Conventional mining and milling	LU, T, GS, WR, E, AQ, N, HC, VS, S, PO, WM		
Reclaimed open pit mines	LU, T, GS, WR, E, AQ, N, HC, VS, S, PO, WM		
Mining of other minerals			
Perlite (Socorro)	LU, T, GS, WR, E, AQ, N, HC, VS, S, WM		
Humate (McKinley)	LU, T, GS, WR, E, AQ, N, HC, VS, S, WM		
Travertine (Cibola)	LU, T, GS, WR, E, AQ, N, HC, VS, S, WM		

Table 5.3-5. Other Actions Concurrent With Uranium Recovery in the Northwestern)
New Mexico Uranium Milling Region* (continued)	

Categories of Actions

Impacts on Resource and Topics†

Cultural Resources Preservation

LU, HC

Numerous Native American sacred sites

*The Northwestern New Mexico Uranium Milling Region includes McKinley County and the northern portions of Cibola, Catron, and Socorro Counties.

†The resources and topics codes include

LU = land use

T = transportation

GS = geology and soils

WR = water resources (wetlands, surface water, and groundwater)

E = ecology (terrestrial, aquatic, and threatened/endangered species)

AQ = air quality (non-radiological)

N = noise

HC = historical and cultural resources

VS = visual and scenic resources

S = socioeconomics

PO = public and occupational health and safety

WM = waste management

individual and programmatic present and RFFAs as reflected by recent EISs (Tables 5.2-2 through 5.2-6).

As described in Chapter 4, construction, operations, aquifer restoration, and decommissioning/reclamation activities associated with uranium ISL facilities can affect different resource areas within each of the uranium milling regions. In conducting a site-specific cumulative effects analysis, an approach such as the CEQ (1997) 11-step process described in Appendix F can be tailored, depending on the current conditions of the affected environment and the level of impacts (SMALL, MODERATE, or LARGE), to a specific resource area.

If a proposed ISL facility (or an expansion/restart) is in compliance with applicable federal and state laws and policies (e.g., the Endangered Species Act) and if the expected impacts to a specific resource area are small, then a Level 1 site-specific cumulative effects analysis would be appropriate. Based on the CEQ (1997) 11-step process described in Appendix F, a Level 1 analysis is based on consideration of the four scoping steps (Steps 1–4) along with two of the three environmental description steps (Steps 6 and 7). Further, brief consideration should be given to the types, sizes, and locations of other present and RFFAs in the uranium milling region (including other uranium ISL facilities) and their contribution to effects on each resource area.

If concerns are identified during the site-specific analysis with respect to the sustainability or quality of a given resource area in the uranium milling region, then a Level 2 cumulative effects analysis would be appropriate. Based on the CEQ (1997) 11-step process (see Appendix F), a Level 2 analysis is based on the same considerations as a Level 1 analysis, with a more detailed evaluation of the types, sizes, and locations of present and RFFAs and their relative contributions to effects on each resource area (Step 8). The effects of each of the other actions (for example, activities included in the EISs identified in Tables 5.2-3 through 5.2-6) would be tabulated and discussed with respect to the timing of different stages (construction, operation, aquifer restoration, and decommissioning/reclamation) of the ISL facility life cycle.

If the site-specific analysis identifies that a specific resource area reflects stresses that exceed regulatory or policy limits, has diminished usage due to quality degradation, or there are

concerns regarding noncompliance with respect to statutory or policy requirements as reflected by moderate or large impacts, then a Level 3 cumulative effects analysis would be appropriate. In undertaking a site-specific Level 3 analysis, each of the CEQ (1997) 11 steps would be applied, including scoping (Steps 1 through 4), environmental description (Steps 5–7) and environmental consequences (Steps 8 through 11). Detailed descriptions and analysis would be used to fully characterize the cumulative effects of the ISL facility and other past, present, and RFFAs on the status of a resource area, such as land use or groundwater, within the affected environment.

A systematic resource-by-resource review of the conditions of the affected environment within each geographic region; the levels of impacts of ISL facilities for all four stages of the ISL lifecycle (construction, operations, aquifer restoration, and decommissioning); and the identification of other past, present, and RFFAs in each designated region, was used to determine the potential level of cumulative effects analysis. The results of this analysis revealed that a Level 1 or Level 2 site-specific cumulative effects analysis would be expected to be sufficient for nine resources in each of the four regions. The nine resources included land use, transportation, geology and soils, air quality, noise, visual and scenic resources, socioeconomics, public and occupational health and safety, and waste management. Another result of this review was that for the four other resources, a Level 1, 2, or 3 analysis might be required. The Level 3 analysis would be highly dependent on local site-specific conditions. The four resources that could potentially be analyzed at this level included surface water resources (primarily wetlands), groundwater resources, terrestrial and aquatic ecology (primarily threatened or endangered species), and historical and cultural resources.

5.5 References

CEQ. "Considering Cumulative Effects Under the National Environmental Policy Act." Washington, DC: Executive Office of the President. 1997.

Committee on Hard Rock Mining on Federal Lands. "Hardrock Mining on Federal Lands." Washington, DC: National Research Council, National Academics Press. 1999.

EPA. "Health and Environmental Impacts of Uranium Contamination in the Navajo Nation: Five Year Plan; as Requested by the House Committee on Oversight and Government Reform." 2008. http://www.epa.gov/region09/superfund/Navajo-nation/pdf/NN-5-Year-Plan-12.pdf (25 February 2009).

NRC. "Expected Uranium Recovery Facility Applications / Restarts / Expansions: Updated 04/24/2009." 2009. http://www.nrc.gov/info-finder/materials/uranium/2008-ur-projects-list-public.pdf (28 April 2009).

6 ENVIRONMENTAL JUSTICE

Environmental justice means that people of all races, cultures, and incomes are treated fairly with regard to the development and implementation (or lack thereof) of environmental laws, regulations, and policies (Executive Order 12898). On February 11, 1994, the President signed Executive Order 12898, "Federal Actions to Address Environmental Justice in Minority Populations and Low-Income Populations," which directs each federal agency to "... make achieving environmental justice part of its mission by identifying and addressing, as appropriate, disproportionately high and adverse human health or environmental effects of its programs, policies, and activities on minority populations and low income populations" (Office of the President, 1994). Executive Order 12898 makes it clear that environmental justice matters also apply to programs involving Native Americans (CEQ, 1997).

On December 10, 1997, the Council on Environmental Quality (CEQ) issued, "Environmental Justice Guidance Under the National Environmental Policy Act." The Council developed this guidance to "... further assist Federal agencies with their National Environmental Policy Act (NEPA) procedures." As an independent agency, the Council's guidance is not binding on the U.S. Nuclear Regulatory Commission (NRC). However, the NRC considered the Council's guidance on environmental justice in developing its own environmental justice analysis procedures.

In August 2004, NRC published a final policy statement in the Federal Register to provide a "... comprehensive statement of the Commission's policy on the treatment of environmental justice matters in NRC regulatory and licensing actions" (NRC, 2004). The NRC Environmental Justice Policy is to use its normal and traditional NEPA review process to meet the goals articulated in Executive Order 12898. "NRC believes that an analysis of disproportionately high and adverse impacts needs to be done as part of the agency's NEPA obligations to accurately identify and disclose all significant environmental impacts associated with a proposed action." In drafting the policy statement, NRC received comments on its draft Environmental Justice Policy on whether environmental justice should be considered in a programmatic or generic environmental impact statement (GEIS). In clarifying its position, NRC noted that for a non-site-specific assessment of potential environmental impacts such as that presented in a GEIS, it is "... difficult to foresee or predict many circumstances, if any, in which a meaningful environmental justice analysis could be completed." However, the final policy statement does not preclude the possibility of an environmental justice analysis in a GEIS if "... a meaningful review can be completed."

NRC has concluded that it can use the GEIS to help conduct a meaningful environmental justice analysis by using population information available through the U.S. Census Bureau, the regional and sub-regional information discussed in Chapter 3, and the potential environmental impacts evaluated in Chapters 4 and 5. The GEIS lists regional resource areas where there is no information indicating that the impacts described in Chapters 4 and 5 would be any different for the identified minority or low-income population than the general population. The GEIS also lists regional resource areas where further site-specific information should be gathered to evaluate whether there is a disproportionately high and adverse environmental or health impact on the minority or low-income populations in the area.

It should be noted, under NEPA, the identification of a disproportionately high and adverse human health or environmental effect on a minority or low-income population does not preclude a proposed agency action from going forward, nor does it necessarily result in a conclusion that

a proposed action is environmentally unsatisfactory. Rather, the identification of such an effect should heighten agency attention to alternatives (including alternative sites), mitigation strategies, monitoring needs, and preferences expressed by the affected community or population (CEQ 1997).

The following sections in this chapter discuss NRC's procedure to conduct an environmental justice analysis and then apply the procedure to the regional areas under consideration in this GEIS.

6.1 Environmental Justice Analysis

6.1.1 Background and Guidance

NRC addresses environmental justice in environmental impact statements (EISs) and, as appropriate, in supplemental EISs (NRC, 2004; 2003, Appendix C). As discussed in Section 1.8, NRC will use the GEIS to prepare a supplemental EIS for the issuance of a new ISL license. Additionally, NRC may use the GEIS to prepare a site-specific environmental assessment or EIS for applications to renew or amend existing ISL licenses. NRC environmental justice guidance (NRC, 2004) discusses the procedure to evaluate potential disproportionately high and adverse impacts associated with physical, socioeconomic, health, and cultural resources to low-income and minority populations. The environmental justice process is shown in Figure 6.1-1.

The first step in the process is to gather demographic and socioeconomic data for the immediate site and surrounding communities to identify minority or low-income populations. The guidance document describes the radius of influence to consider when evaluating potential environmental justice concerns for licensing a uranium recovery facility, as an ISL mill. That radius is normally 1 km [0.6 mi] from the center of the proposed site in urban areas and 6.4 km [4 mi] if the facility is located in a rural area.

Most potential ISL facilities are expected to be located in rural areas, indicating that the 6.4-km [4-mi] radius would generally be appropriate. The NRC final policy statement (NRC, 2004) notes, however, that the distances are intended as guidelines, not requirements. The geographic scale

<u>Components of an Environmental Justice</u> <u>Analysis (CEQ, 1997; NRC, 2004)</u>

Minority population is identified as consisting of individual(s) who are American Indian or Alaskan Native, Asian or Pacific Islander, Black (not of Hispanic origin), or Hispanic.

Low-income population is identified in comparison to statistical poverty thresholds identified in U.S. Census Bureau information.

Disproportionately high and adverse effects include potential effects on both human health and the environment. Disproportionately high and adverse effects are evaluated by determining whether there are one or more attributes that could lead to impacts that would be expected to significantly and adversely affect a minority or low-income population more than the general population as a whole.

considered in a site-specific environmental justice analysis should be appropriate for the potential impact area. Because ISL facilities may employ both local and outside workers {i.e., workers willing to commute more than 48 km [30 mi] (Section 3.2.10.4)}, NRC has decided to evaluate demographic and socioeconomic data within at least an 80-km [50-mi] radius of the existing or potential facilities. This analysis ensures consideration of an adequate sample of the surrounding population, because the goal of environmental justice analysis is to evaluate the communities, neighborhoods, or areas that may be disproportionately impacted (NRC, 2003, Appendix C).




Figure 6.1-1. Environmental Justice Process Flow Chart

NRC guidance recommends using the U.S. Census Bureau "census block group" as the geographic area for evaluating demographic and income data. NRC used this data source and examined delineations of tribal lands and resources for this GEIS. NRC can also use other site-specific information to identify minority or low-income populations not identified through this

demographic data to determine whether further environmental justice analysis is needed in an environmental review for an individual license application.

The next step is to compare the census block group percentage of minority populations and economically stressed households in the area for assessment to the state and county percentages. As general guidance, NRC (2003, Appendix C) notes that differences greater than 20 percentage points may be considered significant and would merit further analysis. Additionally, based on U.S., county, or state date, if either the minority or low-income population percentage in the radius of influence exceeds 50 percent, environmental justice should be considered in greater detail. Depending on a specific facility's location, it is possible that the radius of influence could cross county and state lines—a fact that should be considered when making comparisons. If no minority or low-income populations are identified in the potentially affected area or environmental impact area, then the conclusion should be documented and the environmental justice review is complete.

After minority or low-income populations are identified, the next step is to determine whether there is a "disproportionately high and adverse" impact (human health or environmental effect) to these populations.

NRC guidance recommends determining the impacts of the proposed action in the usual manner, including cumulative and multiple impacts, where appropriate. Environmental impacts and cumulative impacts for facilities using ISL technology are discussed in Chapters 4 and 5 of the GEIS. These impacts have been evaluated to determine whether they would disproportionately affect minority or low-income populations by considering whether there are unique pathways of exposure to these populations compared to the general population. Where a proposed action would not cause adverse environmental impacts, and therefore not cause any high and adverse health or environmental impacts, specific demographic analysis may not be warranted (CEQ, 1997).

The next step is to determine whether the impacts disproportionately impact the minority or low-income populations. In general, populations located next to a site would likely have a disproportionate impact compared to other populations located farther from the site. For example, potential exposure to effluents may be greater to those living closest to the facility, noise and traffic may disrupt nearby residents to a greater extent than those living far from the site, and the potential risk due to accidents may be greater for nearby residents. Additionally, cultural differential patterns of consumption of natural resources may change the impact to the identified population (NRC, 2003, Appendix C). For this GEIS, a subsistence consumption analysis can be used to evaluate whether there are cultural factors that change the estimated "dose" for the sections discussing impacts on public and occupational health and safety. If there are no disproportionate impacts, no further analysis would be needed and the reviewer would document this finding in the environmental justice section (NRC, 2003, Appendix C).

If there are disproportionate impacts to minority or low-income populations, the next step in the analysis would be to evaluate the significance of the impacts to determine whether they are "high and adverse." Impacts that are significant, unacceptable, or above generally accepted levels (such as regulatory limits or state and local statutes and ordinances) may be considered high and adverse. Each impact, and where appropriate, the cumulative and multiple effect of the impacts, should be reviewed for significance. If it can be stated that no combination of the impacts is significant, then they are not disproportionately adverse or high on the minority or low-income populations, and this finding should be documented in the environmental justice section of the environmental review (NRC, 2003, Appendix C).

If there are significant impacts to minority or low-income populations, it is then necessary to look at mitigative measures and benefits. Any mitigation measures that could be taken to reduce the impact should be considered. To the extent practicable, mitigation measures should also reflect the needs and preferences of the affected minority or low-income populations. The environmental review should also discuss benefits of the project to the surrounding communities, including economic benefits (NRC, 2003, Appendix C).

The resulting environmental justice review should indicate whether there is a disproportionately high and adverse human health or environmental impact that is likely to result from the proposed action and if there are any alternatives. It should also indicate any mitigation measures that could be used to reduce this impact and any benefits of the project to the surrounding community. In this way, the final decision makers can weigh all aspects when making the agency decision (NRC, 2003, Appendix C).

6.1.2 Identifying Minority and Low-Income Populations in the Four Geographic Uranium Milling Regions Considered in This GEIS

Demographic and socioeconomic information from the 2000 Census is presented in detail in Sections 3.2.10 (Wyoming West), 3.3.10 (Wyoming East), 3.4.10 (Nebraska-South Dakota-Wyoming), and 3.5.10 (Northwestern New Mexico) for the four geographic regions considered in this GEIS. Minority and low-income populations within the regions were identified using the criteria in NRC guidance (NRC, 2004, 2003) by comparing community demographics to the state level (Table 6.1-1). The distances provided in Table 6.1-1 are given from the border of an identified population (e.g., a reservation boundary) to the nearest existing or potential ISL facility as well as to the farthest ISL facility, based on current information (NRC, 2009).

In the Wyoming West Uranium Milling Region, the only sensitive population identified using the criterion from NRC (2004, 2003) is the Wind River Indian Reservation (Figure 6.1-2). The boundary of the Wind River Indian Reservation is 16 km [10 mi] from the closest potential ISL facility and about 107 km [65 mi] from the farthest potential facility. The reservation has a Native American population of about 35 percent (Eastern Shoshone and Northern Arapaho). This compares to the Wyoming state level of 2.3 percent. The towns of Arapahoe, Ethete, and Fort Washakie are located within the reservation and have both minority (80 percent or more Native American) and low-income populations. The closest potential ISL facility to one of these communities would be about 24 km [15 mi] to the southeast of Arapahoe at Sand Draw.

In the Wyoming East Uranium Milling Region, no minority populations were identified using 2000 Census data and the criteria from NRC (2004, 2003), but Albany County was identified as a low-income population (Figure 6.1-3). Albany County is about 8 km [5 mi] from the nearest location of past, present, or future uranium milling activity in the Wyoming East Uranium Milling Region. Northern Albany County is predominantly rural (see Section 3.3.1), with no population centers or towns identified by the U.S. Census Bureau within the portion of the county that lies within the Wyoming East Uranium Milling Region.

In the Nebraska-South Dakota-Wyoming Uranium Milling Region, the closest sensitive population identified using criteria from NRC (2004, 2003) is the Pine Ridge Indian Reservation,

Table 6.1-1. Minority and Low-Income Populations* in the Four Geographic Uranium Milling Regions Considered in This Generic Environmental Impact Statement					
Uranium Milling	Affected Area Within	Distance (Range) of Project Locations to	Minority	Low-Income	
Region	Region of Influence	Affected Area	Population	Population?	
West Wyoming	Wind River Indian Reservation (Towns of Arapahoe, Ethete, and Fort Washakie)	16–105 km (10–65 mi)	Native American (Eastern Shoshone and Northern Arapaho Tribes)	Yes	
East Wyoming	Albany County	8–161 km (5–100 mi)	None	Yes	
Nebraska- South Dakota- Wyoming	Pine Ridge Indian Reservation (Towns of Oglala and Pine Ridge)	32–161 km (20–100 mi)	Native American (Oglala Sioux Tribe)	Yes	
Northwestern New Mexico	Cibola County	0–43 km (0–27 mi)	Native American and Hispanic Origin	Yes	
	McKinley County	0–5 km (0–3 mi)	Native American	Yes	
	City of Gallup	29–101 km (18–63 mi)	Native American and Hispanic Origin	Yes	
	Town of Grants	16–85 km (10–53 mi)	Some Other Race and Hispanic Origin	Yes	
	Acoma Pueblo (Cibola County)	21–92 km (13–57 mi)	Native American (Acoma)	Yes	
	Laguna Pueblo (Bernanillo, Cibola, Sandoval, Valencia Counties)	27–97 km (17–60 mi)	Native American (Laguna)	Yes	
	Navajo Nation (Cibola and McKinley Counties)	2–74 km (1–46 mi)	Native American (Navajo)	Yes	
	Ramah Navajo Indian Reservation (Cibola and McKinley Counties)	37–64 km (23–40 mi)	Native American (Ramah Navajo)	Yes	
	Tohajiilee Indian Reservation (Cibola and Sandoval Counties)	45–129 km (28–80 mi)	Native American (Tohajiilee)	Yes	
	Zuni Indian Reservation (Cibola and McKinley Counties) Census Bureau. "American FactF	37–80 km (23–50 mi)	Native American (Zuni)	Yes	

*Based on U.S. Census Bureau. "American FactFinder." 2000. http://factfinder.census.gov/home/saff/main.html?_lang=en (18 October 2007 and 25 February 2008).




Figure 6.1-2. Affected Minority and Low-Income Population for the Wyoming West Uranium Milling Region




Figure 6.1-3. Affected Minority and Low-Income Population for the Wyoming East Uranium Milling Region (No Minority Populations Were Identified)

adjacent to the southeastern boundary of the region (Figure 6.1-4). The Pine Ridge Indian Reservation is 48 km [30 mi] from the closest existing and potential ISL facilities at Crow Butte in Dawes County, Nebraska, and about 160 km [100 mi] from the farthest potential facility in Crook County, Wyoming. Communities within the Pine Ridge Indian Reservation include the towns of Oglala and Pine Ridge. Based on U.S. Census Bureau information, these towns have both minority (greater than 90 percent Native American) and low-income populations. They are a little over 75 km [47 mi] from the nearest existing ISL facility at Crow Butte.

In the Northwestern New Mexico Uranium Milling Region (Figure 6.1-5), the potential sensitive minority and low-income populations include the following:

Acoma Indian Reservation

The Acoma Indian Reservation is 21 km [13 mi] from the nearest potential ISL facility and approximately 92 km [57 mi] from the farthest potential known facility. A portion of the Acoma Indian Reservation lies within eastern Cibola County.

Tohajiilee Indian Reservation

The Tohajiilee Indian Reservation is about 45 km [28 mi] from the closest potential ISL facility and approximately 129 km [80 mi] from the farthest potential ISL facility.

Laguna Indian Reservation

The Laguna Indian Reservation is 27 km [17 mi] from the closet potential ISL facility and 97 km [60 mi] from the farthest ISL facility. The majority of the Tohajiilee and Laguna Indian Reservations lie within eastern Cibola County with small portions within Sandoval, Bernalillo, and Valencia Counties.

Navajo Nation

The Navajo Nation represents the largest tribal area and is located approximately 1.6 km [1 mi] from the closest potential ISL facility and 74 km [46 mi] from the farthest known potential ISL facility. A portion of the Navajo Nation lies within McKinley County in the northwestern portion of the Northwestern New Mexico Uranium Milling Region.

Ramah Navajo Nation

The Ramah Navajo Nation is 37 km [23 mi] from the nearest potential ISL facility and 64 km [40 mi] from the farthest potential ISL facility. The majority of the Ramah Navajo Nation lies within western Cibola County.

Zuni Indian Reservation

The Zuni Indian Reservation is 37 km [23 mi] from the nearest potential ISL facility and 80 km [50 mi] from the farthest potential ISL facility. The majority of the Zuni Indian Reservation lies within southwest McKinley County.




Figure 6.1-4. Affected Minority and Low-Income Population for the Nebraska-South Dakota-Wyoming Uranium Milling Region




Figure 6.1-5. Affected Minority and Low-Income Populations for the Northwestern New Mexico Uranium Milling Region

Each of these six tribal areas has a Native American population of greater than 95 percent (compared to the state level of 9.5 percent) and is classified as a low-income population based on 2000 Census information. Where reported, unemployment levels on the reservations are greater than 60 percent (Laguna, Navajo, and Zuni).

Town of Grants

The Town of Grants, located in Cibola County, is about 16 km [10 mi] from the closest potential ISL facility and 85 km [53 mi] from the farthest potential ISL facility. Grants has a Hispanic population of greater than 50 percent.

Sandoval County

A small portion of Sandoval County is included within the eastern border of the Northwestern New Mexico Uranium Milling Region. The southwestern border of Sandoval County is about 37 km [23 mi] from the closest potential ISL facility and 108 km [67 mi] from the furthest ISL facility. The total population of the county is 29.4 percent Hispanic and 16.3 percent Native American. However, the southwestern portion of the county that is nearest to the Grant's Uranium Milling District is expected to have a lower percentage of Native American population than the county as a whole.

McKinley County

McKinley County includes most of the potential ISL facilities identified to date (NRC, 2008) and has a Native American population of almost 75 percent, as compared to the state level of 9.5 percent. McKinley County contains portions of three of the reservations identified in Table 6.1-1. These comprise approximately 35 percent of the area in the county. The percentage of individuals below poverty level in McKinley County (36 percent) and Gallup (21 percent) also identify low-income populations. The Core-Based Statistical Area of Gallup is located 29 km [18 mi] from the nearest potential ISL facility and 101 km [63 mi] from the farthest potential ISL facility. It is located in McKinley County, but outside of the tribal lands.

Cibola County

With the exception of the Navajo Nation, Cibola County contains portions of all of the tribal reservations identified in Table 6.1-1, and they comprise almost 50 percent of the county by area. Cibola County has a Native American population of greater than 40 percent, and the percentage of individuals living below the poverty level in Cibola County (25 percent) and Grants (21.9 percent) indicates low-income populations.

The socioeconomic information from the 2000 Census indicates that all of the existing or potential ISL facilities are located in areas of low income. The census data for the Wyoming East Uranium Milling Region did not identify a minority population. The other milling regions used for this analysis identified Native American or Hispanic populations that may be impacted if an individual ISL facility is located in their proximate area.

6.2 Wyoming West Uranium Milling Region

The affected minority and low-income populations for the Wyoming West Uranium Milling Region are in the Wind River Indian Reservation and the towns of Ethete, Arapahoe, and

Fort Washakie (see Figure 6.1-2). The closest potential ISL facility to the Wind River Indian Reservation is at least 16 km [10 mi] away. Based on current information, the tribal populations on the Wind River Indian Reservation could be located within a 80 km [50 mi] radius of potential ISL facilities and could raise specific environmental justice concerns. The low-income population in the area also triggers an environmental justice analysis for existing and potential facilities located in this area.

General cultural information indicates tribal populations in the Great Plains still use hunting and wild plant gathering, to a limited extent, to supplement family food resources that today are derived primarily from tribal and federal assistance programs or wage labor on and off the reservation. In addition, herbs gathered for subsistence, medicinal, and ritual/ceremonial uses remain important to maintaining traditional cultural practices. Traditional use areas claimed by the tribes are places in which traditional subsistence practices and the procurement of animals and plants for ritual, ceremonial, medicinal, and other traditional needs should be assessed on a site-specific basis. Disruption in the availability of or access to areas in which traditional subsistence and ritual/ceremonial practices can be performed should be considered as having the potential to differentially affect the ability of the tribes in this region to practice their traditional lifeways. No culturally significant places listed in the National Register of Historic Places or the state register are located in the Wyoming West Uranium Milling Region (see Section 4.2.8).

NRC concludes that environmental reviews for ISL facilities located in the Wyoming West Uranium Milling Region would need an environmental justice analysis based on this demographic data. Using current available information, NRC has concluded there are no known cultural factors that would change the Chapters 4 and 5 analyses and conclusions of the potential environmental or health impacts from ISL facility activities for tribal or low-income populations compared to the general population for the following resource areas: land use, transportation, geology and soils, meteorology/climate/air quality, noise, visual/scenic resources, and socioeconomics in the Wyoming West Uranium Milling Region.

NRC also concludes that site-specific information is needed to complete the environmental justice analysis in the following resource areas: water resources, historic and cultural resources, ecological resources, and public and occupational health. Site-specific cultural information should be used to evaluate whether the analyses and conclusions in Chapters 4 and 5 should be supplemented before determining whether the minority or low-income populations in the area would receive a disproportionately high and adverse environmental or health impact from the ISL facility activities.

For further site-specific analyses, staff will consider, among other things:

- Subsistence—In areas where there is a significant consumption of native plants and animals, a subsistence consumption analysis of fish, wildlife, and other natural resources should be done to evaluate the estimated "dose" discussed in the occupational and public health sections.
- Cultural—Site-specific historic and cultural information should be gathered because of the proximity of tribal populations.

The NRC staff would conduct an environmental justice analysis based on the methodologies in the appropriate NRC guidance for site-specific environmental reviews.

6.3 Wyoming East Uranium Milling Region

No minority populations were identified in the Wyoming East Uranium Milling Region using 2000 Census data and the criteria from NRC (2004, 2003). Albany County was identified as a low-income population (Figure 6.1-3). At its closest point, Albany County would be about 8 km [5 mi] from the closest potential ISL facility at Shirley Basin. However, northern Albany County is predominantly rural (see Section 3.3.1) with no population centers or towns identified by the U.S. Census Bureau in the portion of the county that lies within the Wyoming East Uranium Milling Region. For this reason, no environmental justice considerations would be expected for the portion of Albany County that is located within the Wyoming East Uranium Milling Region.

NRC concludes that for ISL facilities located in the Wyoming East Uranium Milling Region, no minority and low-income population will experience a disproportionately high and adverse impact. However, NRC would review environmental justice on a site-specific basis to confirm the GEIS conclusion remains valid. Based on NRC's information, the area in northern Albany County that is nearest potential ISL facilities is sparsely populated. There are no known cultural factors that would change the Chapters 4 and 5 analyses and conclusions of the potential environmental or health impacts from ISL facility activities on this low-income population compared to the general population in this region.

6.4 Nebraska-South Dakota-Wyoming Uranium Milling Region

As identified in Table 6.1-1, the closest affected minority and low-income population for the Nebraska-South Dakota-Wyoming Uranium Milling Region is the Pine Ridge Indian Reservation and the towns of Oglala and Pine Ridge in South Dakota (Figure 6.1-4). The Pine Ridge Indian Reservation is 48 km [30 mi] from the closest existing, and potential, ISL facilities at Crow Butte in Dawes County, Nebraska. Based on current information, the tribal populations on the Pine Ridge Indian Reservation could be located within a 80 km [50 mi] radius of potential ISL facilities and could raise specific environmental justice concerns. The low-income population in the area also triggers an environmental justice analysis for existing and potential facilities located in this area.

General cultural information indicates tribal populations in the Great Plains still use hunting and wild plant gathering, to a limited extent, to supplement family food resources that today are derived primarily from tribal and federal assistance programs or wage labor on and off the reservation. In addition, herbs gathered for subsistence, medicinal, and ritual/ ceremonial uses remain important to maintaining traditional cultural practices. Traditional use areas claimed by the tribes are places in which traditional subsistence practices and the procurement of animals and plants for ritual, ceremonial, medicinal, and other traditional needs should be assessed on a site-specific basis. Disruption in the availability of, or access to, areas in which traditional subsistence and ritual/ceremonial practices can be performed should be considered as having the potential to differentially affect the ability of the tribes in this region to practice their traditional lifeways.

Historically, the land in the area of the Black Hills is seen by tribes in Montana, Wyoming, and South Dakota to have provided both sustenance (for fishing, hunting, and plant food gathering) and spiritual value (i.e., as a place in which important personal and tribal rituals and ceremonies were customarily performed and are still performed today). Devils Tower, or Bear Lodge as it is known to many of the tribes in the region, is located in northeastern Wyoming at the western fringe of the Black Hills in the Nebraska-South Dakota-Wyoming Uranium Milling Region. It is

the site of annual ritual and ceremonial events by tribal members in the month of June. Native American tribes in the region believe that preserving and maintaining access to sacred lands is essential to both cultural and spiritual aspects of traditional Native American societies of the northern plains (Iverson, 1985). The cultural significance of these areas should also be considered during the environmental justice analysis for licensing applications in this region.

In addition, availability of affordable housing with water, electricity, plumbing, and sewer service is a concern at the Pine Ridge Indian Reservation in Shannon County, South Dakota (Housing Assistance Council, 2002; Steele, 2007). Inadequate availability of housing may be a concern with regard to overcrowding and should be evaluated in the environmental justice analysis for the socioeconomic resource area.

NRC concludes that environmental reviews for ISL facilities located in the Nebraska-South Dakota-Wyoming Uranium Milling Region would need an environmental justice analysis based on this demographic data. Using current available information, NRC has concluded there are no known cultural factors that would change the Chapters 4 and 5 analyses or conclusions of the potential environmental or health impacts from ISL facility activities for tribal or low-income populations compared to the general population for the following resource areas in the Nebraska-South Dakota-Wyoming Uranium Milling Region: land use, transportation, geology and soils, meteorology/climate/air quality, noise, and visual/scenic resources.

NRC also concludes that site-specific information is needed to complete the environmental justice analysis in the following resource areas: water resources, historic and cultural resources, ecological resources, public and occupational health, socioeconomics, and visual/scenic resources. Site-specific cultural information should be used to evaluate whether the analysis and conclusions in Chapters 4 and 5 should be supplemented before determining whether the minority or low-income populations in the area would receive a disproportionately high and adverse environmental or health impact from the ISL facility activities.

For further site-specific analyses, staff would consider, among other things:

- Subsistence—In areas where there is a significant consumption of native plants and animals, a subsistence consumption analysis of fish, wildlife, and other natural resources should be conducted to evaluate the estimated "dose" discussed in the occupational and public health sections.
- Cultural—Site-specific historic and cultural information should be gathered because of the proximity of tribal populations.

The NRC staff would conduct an environmental justice analysis based on the methodologies in the appropriate NRC guidance for site-specific environmental reviews.

6.5 Northwestern New Mexico Uranium Milling Region

Based on 2000 Census information and the NRC environmental justice criteria (NRC, 2004, 2003), affected minority and/or low-income populations for the Northwestern New Mexico Uranium Milling Region include Acoma Pueblo, Laguna Pueblo, the Navajo Nation, the Ramah Navajo Indian Reservation, the Tohajiilee Indian Reservation, and the Zuni Indian Reservation (Figure 6.1-4). In addition, minority and low-income populations are identified for Cibola County, McKinley County, the Gallup Core-Based Statistical Area, and the town of Grants. The affected

communities are located throughout the region and are close to potential ISL facilities, based on current information. For example, at least one potential facility would be located within about 1.6 km [1 mi] of the border of the Navajo Nation (Figure 6.1-4) and another would be located near the community of Crownpoint. The location of minority and low-income populations triggers an environmental justice analysis for existing and potential facilities located in this area.

In particular, sensitive communities in proximity to a potential ISL facility would also receive potentially disproportionately high and adverse impacts with regard to water resources in the Northwestern New Mexico Uranium Milling Region. As described in Section 3.5.4, these impacts could include (1) sedimentation in surface waters, (2) degradation of water quality in the ore-bearing aquifer, (3) degradation of groundwater quality near well fields if lixiviant unexpectedly travels from the production zone and beyond the boundaries of the well field, and (4) vertical excursions where barren or pregnant lixiviant migrates into other aquifers above or below the production zone. As described in Section 4.5.4 and Chapters 7 and 8, licensees are required to obtain underground injection control permits and implement monitoring programs and remediation actions to mitigate these potential impacts. In addition, aquifer restoration upon completion of uranium recovery is designed to reduce potential impacts to groundwater quality and use. Site-specific analysis of environmental justice concerns with respect to sensitive communities would be necessary for individual license applications. These site-specific environmental reviews would include consultations with local communities or jurisdictions to evaluate key concerns with respect to water resources.

Land use impacts could result in environmental justice considerations if a potential ISL facility is located near tribal lands or abuts private lands, allottees, or residences, particularly in the checkerboard region where land ownership is complicated. As described in Section 4.5.1, impacts from all phases could (1) change and disturb land uses; (2) restrict access and/or establish right-of-way for access; (3) affect mineral rights and land use by allottees and others; (4) restrict livestock grazing areas and revoke grazing permits; (5) restrict recreational activities; and (6) alter ecological, cultural, and historical resources. Site-specific analysis of environmental justice concerns for sensitive communities would be necessary for individual license applications. These site-specific environmental reviews would include consultations with local communities or jurisdictions to evaluate key land ownership and jurisdictional issues.

Because of the large area covered by tribal lands in the Northwestern New Mexico Uranium Milling Region, there may be disproportionately high and adverse affects related to historical, cultural, and visual resources. As described in Section 3.5.8, there are a large number of cultural and historical sites in the Northwestern New Mexico Uranium Milling Region that could be affected by land-disturbing activities, such as grading roads, installing wells, and constructing surface facilities and well field infrastructure. Impacts to a community's historical and cultural resources may also occur if activities at an ISL facility prevent or limit access to a culturally significant site or affect the visual landscape. The Mount Taylor Traditional Cultural Property listing on June 14, 2008 (Los Angeles Times, 2008) is one example of a culturally significant area that would need to be evaluated for disproportionate potential impacts. As described in Section 4.5.8, site-specific analysis of environmental justice concerns with respect to cultural resources and sensitive communities would be necessary for individual license applications. These site-specific environmental reviews would include consultations with local communities or jurisdictions to evaluate key concerns with respect to water resources.

Western Puebloan Tribes (Acoma and Zuni)

The Acoma and Zuni foster and encourage the continuance of traditional subsistence practices including agriculture and, to a limited extent, herding (Garcia-Mason, 1979; Ladd, 1979). The Acoma and Zuni traditionally reside in clustered settlements or villages. Both tribes view game hunting and the gathering of wild plant foods and herbs for subsistence, medicinal, and ritual/ceremonial uses as central to their traditional cultural practices (Dozier, 1970; Dutton, 1976; Green, 1979; Ladd, 1979).

Traditional agricultural practices in the arid Southwest rely on the availability of arable land with access to reliable sources of water from rainfall and runoff at Zuni and from irrigation at Acoma (Dozier, 1970; Garcia-Mason, 1979). Summer precipitation in the arid upland Southwest is characterized by high spatial and temporal variability. As a result, successful traditional agricultural practice distributes fields in a variety of areas where rainfall, runoff, and other techniques help to maximize the potential for sufficient rainfall to occur in at least one of the fields. Traditional hunting and gathering of wild plant food resources also contribute to annual subsistence to a limited extent. Farming, hunting, and gathering are used to supplement store-bought food items purchased with funds obtained through tribal and federal assistance programs, by working for federal and tribal governments on the reservation, or from wage labor away from the reservation.

Because of Acoma and Zuni reliance on traditional forms of agriculture and hunting and gathering of wild foods to supplement their food resources, disruption in the availability and access to areas in which these traditional subsistence practices can be performed, or disruptions in the ability to gather animal and plant foods, should be considered as having the potential to differentially affect the ability of the Acoma and Zuni tribal members to practice traditional lifeways. In addition, specific types of plants and animals are obtained for use in ritual and ceremonial and, in the case of plants, medicinal contexts. Restriction of access to the places in which these resources might be obtained or in which they have traditionally been obtained should also be considered as a differentially adverse effect to the practice of traditional Acoma and Zuni lifeways.

Navajo Tribe

Traditional Navajo subsistence relies on a mix of small agricultural fields and herding of sheep and goats (Kluckhohn and Leighton, 1974; Bailey and Bailey, 1986). The traditional Navajo settlement pattern is characterized by extended family household clusters, traditionally termed and outfitted (Kluckhohn and Leighton, 1974), that reside in proximity to one another. Several such related households are often spatially dispersed across the landscape. In traditional Navajo practice, agricultural fields are tended by individual households, whereas sheep and goats from related households are combined into larger flocks that graze over wide areas of open range belonging to the combined related households (Downs, 1964; Witherspoon, 1983; Bailey and Bailey, 1986). Goats and sheep, in addition to supplying meat and milk for consumption, also provide wool and mohair for sale and for use in making traditional textiles that are then sold to supplement family income (Adams, 1971; Aberle, 1983). Traditional households often maintain one or more horses and occasionally cattle as well. The horses and cattle are often grazed on the open range wherever sufficient forage is available. Subsistence farming, sheep and goat grazing, and to a far more limited extent, hunting and wild plant gathering, are used to supplement family food resources obtained through tribal and federal assistance programs or wage labor on and off the reservation (Aberle, 1983; Bailey and Bailey, 1986).

Like the Zuni and Acoma tribes, disruption in the availability of or access to areas in which traditional subsistence practices can be performed should be considered as having the potential to differentially affect the ability of the Navajo to practice traditional lifeways. Animals are hunted and plants are gathered for nonsubsistence use as well. Both animals and plants are used for traditional ritual, ceremonial, medicinal, and other needs. Restriction of access to the places in which these resources might be obtained or in which they have traditionally been obtained should also be considered as a differentially adverse effect to the practice of traditional Navajo lifeways.

NRC concludes that environmental reviews for ISL facilities located in the Northwestern New Mexico Uranium Milling Region would need an environmental justice analysis based on this demographic data. Using current available information, NRC has concluded there are no known cultural factors that would change the Chapters 4 and 5 analyses or conclusions of the potential environmental or health impacts from ISL facility activities for tribal or low-income populations compared to the general population for the following resource areas in the Northwestern New Mexico Uranium Milling Region: transportation, meteorology/climate/air quality, noise, or socioeconomic.

NRC also concludes that site-specific information is needed to complete the environmental justice analysis in the following resource areas: water resources, historic and cultural resources, ecological resources, public and occupational health, visual/scenic resources, and land use. Site-specific cultural information should be used to evaluate whether the analyses and conclusions in Chapters 4 and 5 should be revised before determining whether the minority or low-income populations in the area would receive a disproportionately high and adverse environmental or health impact from the ISL facility activities.

For further site-specific analyses, staff would consider, among other things:

- Subsistence—In areas where there is a significant consumption of native plants and animals, a subsistence consumption analysis of fish, wildlife, and other natural resources should be done to evaluate the estimated "dose" discussed in the occupational and public health sections.
- Cultural—Site-specific historic and cultural information should be gathered because of the proximity of tribal populations.

6.6 Summary

Based on 2000 Census information and criteria from NRC guidance (NRC, 2004, 2003), a number of sensitive populations were identified (Table 6.1-1). NRC concludes potential environmental justice concerns exist in three of the identified uranium milling regions. All of the identified milling regions are located in low-income areas. Environmental reviews for ISL facilities located in the Wyoming East Uranium Milling Region do not need an environmental justice analysis, because demographic data failed to identify a minority or low-income population that has the potential to receive disproportionately high and adverse environmental or health impacts compared to the general population in the area. Minority populations and tribal lands were identified in (1) the Wyoming West, (2) the Northwestern New Mexico, and (3) the Nebraska-South Dakota-Wyoming Uranium Milling Regions. This situation triggers NRC's obligation to conduct an environmental justice analysis in these three regions.

While the GEIS does not identify impacts that are disproportionately high and adverse for a minority or low-income area, it does identify resource areas that could raise environmental justice concerns and notes where site-specific information is needed to complete the environmental justice analysis. For example, resource areas are identified where there are no known cultural factors that would change the Chapters 4 and 5 analyses or conclusions of the potential environmental or health impacts from ISL facility activities for tribal or low-income populations compared to the general population for specific resource areas in each region.

Other regional resource areas were identified that need site-specific information to evaluate whether the analyses and conclusions in Chapters 4 and 5 should be revised when determining whether the minority or low-income populations in the area would receive a disproportionately high and adverse environmental or health impact from the ISL facility activities. In those cases, the revised impact analysis would be used in the environmental justice analysis to determine whether there is a disproportionately high and adverse environmental or health impact on these minority or low-income populations.

6.7 References

Adams, W.Y. "Navajo Ecology and Economy: A Problem in Cultural Values." *Apachean Culture History and Ethnology.* K. Basso and M. Opler, eds. pp. 77–81. Anthropological Papers of the University of Arizona Number 21, Tucson Arizona: University of Arizona Press. 1971.

Aberle, D.F. "Navajo Economic Development." In, *Handbook of North American Indians, Volume 10, Southwest.* Alfonso Ortiz,ed. Washington DC: Smithsonian Institution. pp. 641–658. 1983.

Bailey, G. and R.G. Bailey. *A History of the Navajos: The Reservation Years*. Santa Fe, New Mexico: School of American Research Press. 1986.

CEQ. "Environmental Justice Guidance Under the National Environmental Policy Act." Washington, DC: CEQ. December 1997.

Downs, J.F. The Navajo. New York: Holt, Rinehart, and Winston. 1971.

Downs, James F. "Animal Husbandry in Navajo Society and Culture." *University of California Publications in Anthropology, Volume 1*. Berkeley, California: University of California Press. 1964.

Dozier, E.P. *The Pueblo Indians of North America*. Prospect Heights, Illinois: Waveland Press. 1970.

Dutton, B.P. The Pueblos. Englewood Cliffs, New Jersey: Prentice-Hall. 1976.

EPA. "Final Guidance for Incorporating Environmental Justice Concerns in EPA's NEPA Compliance Analyses." Washington, DC: EPA. April 1998.

Garcia-Mason, V. "Acoma Pueblo." *Handbook of North American Indians, Volume 9, Southwest.* Alfonso Ortiz, ed. Washington DC: Smithsonian Institution. pp. 450–466. 1979.

Green, J., ed. *Zuni: Selected Writings of Frank Hamilton Cushing*. Lincoln, Nebraska: University of Nebraska Press. 1979.

Housing Assistance Council. "Taking Stock: Rural People, Poverty, and Housing at the Turn of the 21st Century." Washington, DC: Housing Assistance Council. December, 2002.

Kluckhohn, C. and D. Leighton. *The Navajo*. Revised Edition. Cambridge, Massachusetts: Harvard University Press. 1974.

Ladd, E.J. "Zuni Economy." In *Handbook of North American Indians, Volume 9, Southwest.* Alfonso Ortiz, ed. Washington DC: Smithsonian Institution. pp. 492–498. 1980.

Los Angeles Times. "Tribes Get Mt. Taylor Listed as Protected." Los Angeles Times, June 15, 2008. http://articles.latimes.com/2008/jun/15/nation/na-mountain 15> (19 February 2009)

NRC. "Expected Uranium Recovery Facility Applications/Restarts/Expansions: Updated 04/24/2009." 2009. http://www.nrc.gov/info-finder/materials/uranium/2008-ur-projects-list-public.pdf (28 April 2009).

NRC. "Policy Statement on the Treatment of Environmental Justice Matters in NRC Regulatory and Licensing Actions." *Federal Register* 69. pp. 52040–52048. August 24, 2004.

NRC. NUREG-1748, "Environmental Review Guidance for Licensing Actions Associated with NMSS Programs." Final Report. Washington, DC: NRC. August 2003.

NRC. NUREG–1508, "Final Environmental Impact Statement to Construct and Operate the Crown Point Uranium Solution Mining Project, Crown Point, New Mexico." Washington, DC: NRC. February 1997.

Office of the President. "Executive Order 12898 on Federal Actions to Address Environmental Justice in Minority Populations and Low-Income Populations With Accompanying Memorandum." Washington, DC: Office of the President. February 11, 1994.

Steele, J.Y.B. "Testimony of the Honorable John Yellow Bird Steele, President of the Oglala Sioux Tribe, Oversight Hearing on Indian Housing Before the Senate Committee on Indian Affairs, March 22, 2007." Washington, DC: U.S. Senate. March 2007.

U.S. Census Bureau. "American FactFinder." 2000. http://factfinder.census.gov/home/saff/main.html?_lang=en (18 October 2007 and 25 February 2008).

Witherspoon, G. "Navajo Social Organization". *Handbook of North American Indians, Volume 10, Southwest.* Alfonso Ortiz, ed. Washington DC: Smithsonian Institution. pp. 524–535. 1983.

7 POTENTIAL BEST MANAGEMENT PRACTICES, MITIGATION MEASURES, AND MANAGEMENT ACTIONS TO MITIGATE ADVERSE ENVIRONMENTAL IMPACTS

7.1 Introduction

This chapter describes potential best management practices, mitigation measures, and management actions that a licensee or facility operator might use to reduce potential adverse impacts associated with construction, operation, aquifer restoration, and decommissioning of an *in-situ* leach (ISL) milling facility. The Council on Environmental Quality (CEQ) defines mitigation as (40 CFR 1508.20):

- Avoiding the impact altogether by not taking a certain action or parts of an action.
- Minimizing impacts by limiting the degree or magnitude of the action and its implementation.
- Rectifying the impact by repairing, rehabilitating, or restoring the affected environment.
- Reducing or eliminating the impact over time by preservation and maintenance operations during the life of the action.
- Compensating for the impact by replacing or providing substitute resources or environments.

Potential mitigation measures can include general best management practices and more site-specific management actions.

7.2 Best Management Practices

How Are Adverse Impacts Mitigated?

Best Management Practices are techniques, methods, processes, activities, or incentives that are more effective at delivering a particular outcome. Best management practices can also be defined as efficient and effective ways of meeting a given objective based on repeatable procedures that have proven themselves over time. Well-designed best management practices combine existing managerial and scientific knowledge with knowledge about the resource being protected. The Wyoming Department of Environmental Quality (WDEQ) defines best practicable technology as "A technology based process determined by WDEQ as justifiable in terms of existing performance and achievability (in relation to health and safety) which minimizes, to the extent safe and practicable, disturbances and adverse impacts of the operation on human or animal life, fish, wildlife, plant life and related environmental values." (WDEQ, 2007).

Management Actions are active measures a licensee or facility operator implements to reduce potential adverse impacts to a specific resource area. These site-specific actions are sometimes related to environmental (or adaptive) management systems (CEQ, 2007).

Best management practices are processes, techniques, procedures, or considerations that can be used to cost-effectively avoid or reduce the potential environmental impacts. While best management practices are not regulatory requirements, they can overlap and support such requirements. Best management practices would not replace any U.S. Nuclear Regulatory Commission (NRC) requirements or other local, state, or federal regulations.

7.3 Management Actions

Management actions are those that the licensee specifically implements to reduce potential adverse impacts. These actions include compliance with applicable government agency

stipulations or specific guidance, coordination with government agencies or interested parties, and monitoring of relevant ongoing and future activities. If appropriate, corrective actions could be implemented to limit the degree or magnitude of a specific action leading to an adverse impact (reducing or eliminating the impact over time by preservation and maintenance operations) and repairing, rehabilitating, or restoring the affected environment.

Licensees may also minimize potential adverse impacts through specific management actions. These may be part of a broad, more formalized environmental (or adaptive) management system similar to those described in CEQ (2007), or they may be more focused on a particular impact. In establishing management actions, the licensee should create measurable environmental objectives with measurable goals and targets (for example, pollution prevention goals for reducing waste). The licensee then would implement these programs, procedures, and controls for monitoring and measuring progress; document progress; and, if appropriate, institute corrective actions. These management actions may be established through standard operating procedures that are reviewed and approved by the appropriate local, state, or federal agency (including NRC). NRC may also establish requirements for management actions by identifying license conditions. These conditions are written specifically into the NRC source and byproduct material license and then become commitments that are enforced through periodic NRC inspections. As part of this oversight, the NRC staff evaluates violations of specific license commitments to determine their impact on safety and the environment. Depending on significance, NRC may levy a written notice of violation and, in certain circumstances, a civil penalty such as a fine. In no case will licensees who cannot achieve and maintain adequate levels of safety be permitted to continue to conduct NRC-licensed activities. Specific aspects of inspection and enforcement of the terms and conditions of an NRC license for an ISL facility can only be addressed at the site-specific levels, and each enforcement action is dependent on the circumstances of the case. In addition, licensees will be subject to requirements and inspections associated with other necessary permits issued by other state and federal agencies for an ISL facility (see Sections 1.6 and 1.7).

The management actions should specifically describe how mitigation commitments would be implemented and reflect available information about these actions. In an environmental management system approach, planned mitigation actions can be revised as more specific and detailed information becomes available. Typically, monitoring activities could be conducted during all phases of the project to ensure the mitigation of potential adverse impacts.

7.4 Potential Best Management Practices, Management Actions, and Mitigation Measures

Potential best management practices and mitigation measures that are commonly used to minimize potential adverse impacts are listed in Table 7.4-1. The list is based on historical best management practices and mitigation measures used for existing and planned ISL uranium recovery facilities (NRC, 1997, 1998, 2006a,b; Energy Metals Corporation, U.S., 2007; WDEQ, 2007). The list in Table 7.4-1 is not comprehensive and does not imply that NRC endorses these measures. Because the practices, actions, and measures identified in Table 7.4-1 have been developed for a broad geographic area, each practice or mitigation measure described in the table may not apply to a specific project. The list provides a foundation for developing customized management and mitigation plans for a proposed facility or project.

Table	7.4-1. Summary of Potential Best Management Practices and Management Actions
Environmental Resource	Potential Best Management Practices and Management Actions
Land use	 Limit land disturbance to only what is necessary for operation. Conduct historic and cultural resource surveys prior to land disturbance. Conduct ecological resource surveys prior to land disturbance. Reclaim lands disturbed during the construction process. Decontaminate and decommission facilities. Reclaim lands disturbed by surface facilities no longer needed. Plug and abandon wells.
Transportation	 Use dedicated tanker trucks for transporting uranium-loaded and barren resins from satellite facilities. Use accepted industry codes and standards for handling and transporting hazardous chemicals. Maintain shipping records (bill of lading) to identify nature and quantity of shipped materials. Conduct surveys of truck exterior and cab prior to each shipment of yellowcake or resin. Establish an emergency response plan for yellowcake spill and other potential transportation accidents. Implement safe driving and emergency response training for personnel and truck drivers. Use check-in/check-out or global positioning satellite technology to track shipments. Install communication systems to connect trucks to shipper/receiver/emergency responders.
Geology and soils	 Use structures to temporarily divert and/or dissipate surface runoff from undisturbed areas around the disturbed areas. Retain sediment within the disturbed areas by using silt fencing, retention ponds, and hay bales. Salvage and stockpile topsoil from the central plant facility area and from well field access roads so that wind and/or water erosion can be avoided (e.g., graded stockpiles, temporary vegetative cover, fencing and signs, sedimentation catchments). Fill pipeline and cable trenches with excavated rock and soil soon after completion and regrade to surrounding topography. Reestablish temporary or permanent native vegetation as soon as possible after disturbance. Construct roads to minimize erosion (e.g., surface with a gravel road base, construct stream crossings at right angles with adequate embankment protection and culvert installation, and provide adequate road drainage with runoff control structures and revegetation). Implement a spill prevention and cleanup plan to minimize soil contamination. Collect and monitor soils and sediments for potential contamination including areas used for land application of treated waste water, transport routes for yellowcake and ion exchange resins, and well field areas where spills or leaks are possible.

Table 7.4-1.	Summary of Potential Best Management Practices and Management Actions (continued)
Environmental Resource	Potential Best Management Practices and Management Actions
Surface water	 Follow construction practices to reduce potential impacts as defined by the U.S. Army Corps of Engineers permitting process. Minimize disturbance of surface areas and vegetation, which would minimize changes in surface-water flow and soil porosity that would change infiltration and runoff rates. Minimize physical changes to drainage channels by building bridges or culverts where roadways would intersect areas of intermittent water flow. Use erosion and runoff control features such as proper placement of pipe, grading to direct runoff away from water bodies, and use of riprap at these intersections to make bridges or culverts more effective. Use sediment-trapping devices such as hay or straw bales, fabric fences, and devices to control water flow and discharge to trap sediments moved by runoff. Maintain natural contours as much as possible, stabilize slopes, and avoid unnecessary off-road vehicle travel to minimize erosion. Train employees in the handling, storage, distribution, and use of hazardous materials. Conduct fueling operations and store hazardous materials and other chemicals in bermed areas with proper set back distances from water bodies. Provide rapid response cleanup and remediation capability, techniques, procedures, and training for potential spills. Prepare and implement a Storm Water Pollution Prevention Plan consistent with state and federal standards for construction activities. Implement a spill prevention and cleanup plan to minimize soil contamination. Conduct land application of treated waste water activities in a manner consistent with local climate, soil, and vegetation conditions to ensure excess irrigation does not run off into surface water.
Groundwater	 Recycle water collected in subsurface areas for use in dust suppression and other activities. Implement measures to minimize water use during operations. Minimize surface disturbance, which will minimize changes in surface-water flow and subsequent infiltration. Implement a spill prevention and cleanup plan to minimize soil contamination. Provide rapid response cleanup and remediation capability, techniques, procedures, and training for potential spills. Monitor to detect and define unanticipated surface spills, releases, or similar events that may infiltrate into the groundwater system. Manage water balance to ensure hydraulic flow into production zone. Monitoring well pressures to detect leaks. Install monitoring wells in well field and near surface impoundments to monitor for potential lixiviant that travels beyond the production zone or for process solution leaks from impoundments. Manage pumping and injection to control and recover excursions. Monitor closest private domestic, livestock, and agricultural wells as appropriate during operations.

Table 7.4-1.	Summary of Potential Best Management Practices and Management Actions (continued)
Environmental Resource	Potential Best Management Practices and Management Actions
Ecology	 Use measures to control erosion, dust, and particulates that may affect ecological resources from construction, operation, aquifer restoration, and decommissioning. Use dust suppression measures to minimize wind and other erosion and aid recovery on disturbed areas. Conduct pre-construction surveys to evaluate important ecological resources and habitats and to determine the reclamation potential of sites. Implement measures to relocate or avoid sensitive species. Minimize groundbreaking or land-clearing activities during the critical nesting period for migratory birds. Collect data to plan to restore disturbed areas and minimize impacts to sensitive habitats before ground-disturbing activities. Phase construction to the extent practicable. Limit grading activities to the phase immediately under construction, and limit ground disturbance to areas necessary for project-related construction activities. Revegetate with appropriate native species to minimize potential for invasive species. Use weed control as necessary.
Air quality	 Reduce fugitive dust emissions using standard dust control measures (e.g., water application, speed limits). Reduce maximum fugitive dust by coordinating dust-producing activities. Use fossil-fuel vehicles that meet applicable emission standards. Reclaim or re-vegetate disturbed areas. Reduce diesel particulate matter emissions using measures such as particle traps and other technological or operational methods. Ensure that diesel-powered construction equipment is properly tuned and maintained. Use ultra-low sulfur diesel fuel. Use newer, cleaner equipment. Avoid leaving equipment unnecessarily idling or operating.
Noise	 Avoid construction activities at night. Use sound controls on operating equipment and facilities. Use personal hearing protection for workers in high noise areas.
Historic and cultural resources	 Consult with appropriate state and tribal historic preservation officers. Ensure that onsite employees complete cultural resource sensitivity and protection training to reduce the potential for intentional or accidental harm to sites or artifacts. Conduct pre-construction surveys to ensure that work would not affect important archaeological resources. Develop additional mitigation measures such as documenting and collecting resources according to a cultural resource management plan if construction threatens important archaeological resources and modification or relocation of facilities and roads is not feasible.

Table 7.4-1.	Summary of Potential Best Management Practices and Management Actions (continued)
Environmental Resource	Potential Best Management Practices and Management Actions
Visual and Scenic	 Use exterior lighting only where needed to accomplish facility tasks. Limit the height of exterior lighting units. Use shielded or directional lighting to limit lighting only to areas where it is needed.
Socioeconomics	 Purchase materials from local vendors as appropriate. Hire local employees and contractors.
Occupational and public health and safety	 Use ventilation to keep radon levels as low as is reasonably achievable. Use vacuum dryers, bag filters, and vapor filtration to reduce particulate emissions during yellowcake drying. Use high-efficiency particulate air filters or similar controls for particulates. Use personal monitoring devices and respirators as appropriate. Design task procedures to reduce potential accidents. Implement health and safety procedures and administrative controls to minimize worker risks during construction and operations.
Waste and hazardous materials	 Recycle wastewater to reduce the amount of water needed for facilities and the amount of wastewater that could require disposal. Use decontamination techniques that reduce waste generation. Institute preventive maintenance and inventory management programs to minimize waste from breakdowns and overstocking. Recycle nonradioactive materials where appropriate. Encourage the reuse of materials and use of recycled materials. Avoid using hazardous materials when possible. Develop a spill prevention plan for petroleum products and other hazardous materials. Ensure that equipment is available to respond to spills, and identify the location of such equipment. Inspect and replace worn or damaged components. Salvage extra materials and use them for other construction activities or for regrading activities.
Utilities, energy, and materials	 Implement procedures and equipment that would minimize the use of utility services, energy, and materials. Incorporate high-performance and sustainable building criteria into the design and construction of nonnuclear facilities.

7.5 References

CEQ. "Aligning National Environmental Policy Act Processes With Environmental Management Systems. A Guide for NEPA and EMS Practitioners." Washington, DC: CEQ. April 2007.

Energy Metals Corporation, U.S. "Application for USNRC Source Material License Moore Ranch Uranium Project, Campbell County, Wyoming: Environmental Report." Casper, Wyoming: Energy Metals Corporation, U.S. [ADAMS Accession Number: ML072851249] September 2007.

NRC. "Environmental Assessment for the Addition of the Reynolds Ranch Mining Area to Power Resources, Inc.'s Smith Ranch/Highlands Uranium Project Converse County, Wyoming." Source Material License No. SUA–1548. Docket No. 40-8964. Washington, DC: NRC. November 2006a.

NRC. "Environmental Assessment for Amendment to Source Materials License SUA-56: Ground Water Alternate Concentration Limits, Western Nuclear, Inc. Split Rock Uranium Mill Tailings Site, Jeffrey City, Fremont County, Wyoming." Source Materials License SUA-56. Docket No. 40-1162. Washington, DC: NRC. August 2006b.

NRC. "Environmental Assessment for Renewal of Source Material License No. SUA–1534. Crow Butte Resources Incorporated Crow Butte Uranium Project Dawes County, Nebraska." Docket No. 40-8943. Washington, DC: NRC. February 1998.

NRC. NUREG-1508, "Final Environmental Impact Statement To Construct and Operate the Crownpoint Uranium Solution Mining Project, Crownpoint, New Mexico." Washington, DC: NRC. February 1997.

WDEQ. "Guideline No. 4: *In-Situ* Mining." Cheyenne, Wyoming: WDEQ, Land Quality Division. 2007. http://deq.state.wy.us/lqd/quidelns/Guide4=3-00.pdf> (18 December 2007).

8 ENVIRONMENTAL MONITORING ACTIVITIES

8.1 Introduction

Monitoring programs, in general, are developed for *in-situ* leach (ISL) facilities to verify compliance with standards for the protection of worker health and safety in operational areas and for protection of the public and environment beyond the facility boundary. Worker safety monitoring programs are developed as part of a radiological protection program summarized in Section 2.7. This chapter discusses environmental monitoring programs that address the environment beyond the operational areas.

Monitoring programs provide data on operational and environmental conditions so that prompt corrective actions can be implemented when adverse conditions are detected. In this regard, monitoring helps to limit potential environmental impacts at ISL facilities. Required monitoring programs can be modified to address unique site-specific characteristics by the addition of license conditions resulting from the conclusions of the U.S. Nuclear Regulatory Commission's (NRC) safety and environmental reviews.

The discussion of monitoring programs in this section is organized by the following general categories:

- Radiological monitoring (Section 8.2)
- Physiochemical monitoring (Section 8.3)
- Ecological monitoring (Section 8.4)

Descriptions of typical monitoring programs are provided in this chapter. Other NRC guidance documents (NRC, 2007a, 2003, 1980) provide more detailed descriptions.

8.2 Radiological Monitoring

NRC regulations at 10 CFR Parts 20 and 40 address radiological effluents and exposures to the public. NRC requires that licensees have an effluent and environmental monitoring program that complies with these rules. An effluent and environmental monitoring program includes a number of monitoring sites where direct radiation measurements are made and surface waters, groundwater, sediments, soils, and the air are sampled for radionuclides. Licensees must document the sampling and monitoring results and maintain records for a specified period of time. In addition, under 10 CFR 40.65, licensees must submit the results of the effluent and environmental monitoring program to NRC twice a year.

General radiological monitoring practice is described in NRC (1980). Although this regulatory guidance was developed for conventional uranium mills, both NRC and the Wyoming Department of Environmental Quality (WDEQ) (NRC, 2003, WDEQ, 2007) have recommended it for ISL facilities. Other acceptable approaches to radiological monitoring are described in a series of NRC guidance documents listed in NRC (2003, Section 5.7).

8.2.1 Airborne Radiation Monitoring Program

For offsite air monitoring, licensees must establish monitoring stations and environmental sampling areas. Sampling locations are selected based on the proposed facility, nearest

residences, and population centers. As described in NRC (1980), offsite air quality is typically monitored for particulates and radon at a variety of locations near the facility, including the following:

- At least three locations at or near the site boundary;
- At the nearest residence or occupiable structure within 10 km [6 mi] of the site with the highest predicted airborne radionuclide concentrations;
- At least one residence or occupiable structure where predicted doses exceed 5 percent of the standards in 40 CFR Part 190;
- A remote location representing background conditions.

The guidance recommends sampling locations be the same as those used to establish preoperational baseline conditions; filters be changed at least weekly, depending on dust conditions; and radon-222 be monitored continuously for at least 1 week per month (NRC, 1980, Section 2.1).

8.2.2 Direct Radiation Monitoring

Direct radiation or gamma monitoring is needed to establish a baseline external radiation level before ISL operations begin, and thereafter to determine if there has been an increase in external radiation exposure from ISL facility operations. Devices to measure direct radiation levels typically are co-located with air sampling stations, but also need to be located to measure the potential direct radiation exposure to the public (NRC, 1980).

8.2.3 Soils and Sediments Monitoring

Soils and sediments are typically monitored annually, both onsite and offsite (NRC, 1980). For consistency, soil sampling locations are generally the same as those for the airborne radiation monitoring program (see Section 8.2.1), and sediment samples should be collected from surface water locations (see Section 8.3.3). Sampling is conducted both at the surface and across a soil-depth profile to a depth of about 1 m [3 ft] or until rock is encountered. These sampling programs may include surveys for gamma radiation, as well as sampling for natural uranium, thorium-230, and lead-210.

As an example of soil and sediment monitoring, the operator of the Crow Butte ISL uranium facility in Dawes County, Nebraska, implemented a soil monitoring program that involves sampling surface soil at the plant site before and after topsoil removal, at evaporation pond sites before excavation, and at air sampling stations (NRC, 1998).

8.2.4 Vegetation, Food, and Fish Monitoring

If a potentially significant exposure pathway is identified, vegetation (forage), food, and fish samples may be collected and analyzed for radionuclides in accordance with NRC sampling location and sampling frequency guidance (NRC, 1980, Section 2). Vegetation should be sampled three times during the growing season, and livestock grazing within 3 km [5 mi] of the site are sampled at the time of slaughter.

8.2.5 Surface Water Monitoring

Water and bed-sediment samples from perennial streams, standing water bodies (ponds, lakes, etc.) and water samples from springs within and near the ISL facility are tested periodically to determine whether contaminants are leaving the facility through surface runoff.

Sampling frequency and distribution are site specific and established by license condition. For example, at the Crow Butte ISL uranium facility in Dawes County, Nebraska, the effluent monitoring program requires one upstream and one downstream sample for each stream passing through the well field area, as well as quarterly sampling from each water impoundment area in the well field area (NRC, 1998).

8.2.6 Groundwater Monitoring

Environmental monitoring of groundwater for radiological constituents at an ISL facility is similar to chemical constituent groundwater monitoring discussed in Section 8.3.1; however, the areal extent of environmental monitoring can go beyond the well field, as needed, based on site-specific conditions. As discussed in NUREG–1569 (NRC, 2003a, Section 2.2.3), the applicant is required, as part of site characterization, to survey and report locations of all privately owned wells within 3.3 km [2 mi] of the permit area and their current uses and production rates to assess potential impacts on these wells due to the ISL operations. Required environmental monitoring programs (NRC, 1980) include routine monitoring of all downgradient public wells that could be used for drinking water, livestock watering, or crop irrigation.

8.3 Physiochemical Monitoring

Environmental monitoring for chemical constituents at ISL facilities, as needed to comply with environmental requirements or license conditions, is expected to overlap with radiological monitoring activities discussed in Section 8.2 (e.g., sampling of surface water, sediments, soils). The chemical analyses are established on a site- and process-specific basis, and include, but are not limited to, the measurements of sulfate or bicarbonate (or total alkalinity), pH, uranium, iron, aluminum, and heavy metals. Unique and important aspects of physiochemical monitoring at ISL facilities primarily include the groundwater and well field monitoring activities discussed in this section.

8.3.1 Well Field Groundwater Monitoring

The ISL production process directly affects groundwater near the operating well field. For this reason, groundwater conditions are extensively monitored both before and during operations.

8.3.1.1 Pre-Operational Groundwater Sampling

Typically, a licensee must establish baseline groundwater quality before beginning uranium production in a well field. This is done to characterize water quality in monitoring wells that are used to detect lixiviant excursions from the production zone, to recover excursions, and to establish standards for aquifer restoration after uranium recovery ends. General criteria for establishing baseline water quality are described in NRC (2003, Section 2.7)

Baseline water quality can be established through examining records and reports for existing local water wells and by sampling wells developed for the ISL program before production

begins. Although it will vary with deposit and aquifer geometry, a typical sampling to establish baseline conditions is about one production or injection well for every 1.6 ha [4 acres], all wells in the monitoring ring, and wells in aquifers above and below the confining layers for the production zone. Wells are sampled periodically for 25 or more major, minor, and trace elements and other parameters such as pH, specific conductivity, and total dissolved solids (see Table 8.2-1). Sampling should ensure that a stable baseline water quality is established. To determine baseline water quality conditions, at least four sets of samples, spaced sufficiently to indicate seasonal variability, should be collected and analyzed for each listed constituent (NRC, 1997, 1998, 2003).

Table 8.2-1. Typical Baseline Water Quality Parameters and Indicators for Groundwater*				
	Physical Indicators			
Specific Conductivity	Total Dissolved Solids†	pH‡		
	Major Elements and Ions	S		
Alkalinity	Chloride	Sodium		
Bicarbonate	Magnesium	Sulfate		
Calcium	Nitrate			
Carbonate	Potassium			
	Trace and Minor Elements			
Arsenic	Iron	Selenium		
Barium	Lead	Silver		
Boron	Manganese	Uranium		
Cadmium	Mercury	Vanadium		
Chromium	Molybdenum	Zinc		
Copper	Nickel			
Fluoride	Radium-226§			
Radiological Parameters				
Gross Alpha∥	Gross Beta			

^{*}Based on U.S. Nuclear Regulatory Commission (NRC). NUREG–1569, "Standard Review Plan for *In-Situ* Leach Uranium Extraction License Applications—Final Report." Table 2.7.3-1. Washington, DC: NRC. June 2003.

8.3.1.2 Groundwater Quality Monitoring

For early detection of potential horizontal and vertical excursions of lixiviants from the production zone, monitoring wells are situated around the well fields, in the aquifers overlying and underlying the ore-bearing production aquifers within the well field. Monitoring well placement is based on what is known about the nature and extent of the confining layer and presence of drill holes, hydraulic gradient, and aquifer transmissivity and well abandonment procedures used in the region. For example, monitoring wells should be placed downgradient from the production zone to detect excursion plumes. Monitoring wells completed in the uranium bearing horizon must be in hydraulic communication with the production zone to be effective (i.e., groundwater can easily flow between the production zone and the monitoring wells). Additional, more closely spaced wells may be necessary if there are preferred flow paths

[†]Laboratory only.

[‡]Field and laboratory determination.

[§]If site initial sampling indicates the presence of thorium-232, then radium-228 should be considered in the baseline sampling, or an alternative may be proposed.

[|]Excluding radon, radium, and uranium.

in the aquifer (preferred flow paths are identified in the subsurface drilling program discussed in Section 2.11.4). If an excursion is detected, additional monitoring wells may also be installed to delineate the extent of the excursion (NRC, 1998).

The ability of a monitoring well to detect groundwater excursions is influenced by several factors, such as the thickness of the aquifer monitored, the distance between the monitoring wells and the well field, the distance between adjacent monitoring wells, the frequency of groundwater sampling, and the magnitude of changes in chemical indicator parameters (see bulleted list below) that are monitored to determine whether an excursion has occurred.

The spacing, distribution, and the number of monitoring wells at a given ISL facility are site specific and established by license condition. For example, at the Smith Ranch ISL uranium facility, Wyoming, the monitoring wells for detecting horizontal excursions are located approximately 150 m [500 ft] beyond the well field perimeter, with a maximum spacing of 150 m [500 ft] between wells (NRC, 2006). At the proposed ISL facility at Crownpoint, New Mexico, the applicant proposed that wells completed in the production zone (Westwater Canyon formation) encircle each well field 140 m [460 ft] from the outermost production or injection wells with 140 m [460 ft] between each monitoring well (NRC, 1997).

Spacing for monitoring wells to detect vertical excursions in overlying and underlying aquifers at uranium ISL facilities is variable and ranges from 1 well per 1.2 ha [3 acres] to 1 well per 2 ha [5 acres] (NRC, 2006; 1998; 1997; Mackin, et al., 2001). In some cases, hydrologic conditions are such that underlying aquifers may not need to be monitored. For example, at the Crow Butte ISL facility in Dawes County, Nebraska, the underlying confining layer is very thick (more than 300 m [1,000 ft]), and the underlying aquifer is not used as source of water (NRC, 1998).

Generally, a small group of parameters provides early warning of an excursion. These indicators are based on lixiviant chemistry and groundwater geochemistry (NRC, 2003, Section 5.7.8). The best excursion indicators are measurable and more highly concentrated in the lixiviant during ISL operations than in the natural groundwater. Typical excursion indicators include the following:

- Chloride (CI). Chloride does not interact strongly with the minerals in the aquifer (a
 conservative tracer), is easily measured, and CI concentration significantly increases
 during the ISL process because of ion exchange reactions in the milling circuit.
- Specific conductivity. Lixiviants have higher total dissolved solids than the local groundwater and therefore, have a higher specific conductivity. Elevated specific conductivity measurements, therefore, may indicate an excursion has taken place. If conductivity is used to estimate total dissolved solids, measurements will be normalized to a reference temperature (usually 25 °C [77 °F]) because of the temperature dependence of conductivity (Staub, et al., 1986; Deutsch, et al., 1985).
- Total alkalinity (carbonate plus bicarbonate plus hydroxide). This is appropriate for ISL operations where sodium bicarbonate or carbon dioxide is used in the lixiviant.

Cations such as calcium and sodium are usually found at significantly higher levels in lixiviants, but these elements tend to interact more strongly with the minerals in the aquifer. This interaction tends to delay the arrival of calcium and sodium at a monitoring well. For this reason, calcium and sodium should generally not be used as excursion indicators. Similarly,

some major ions such as sulfate are present in significantly higher concentrations in the lixiviants, but complex reduction-oxidation chemistry may complicate the interpretation of the results (NRC, 2003, Section 5.7.8).

An excursion is detected when the concentrations of one or more of the excursion indicators exceed the upper control limit (UCL) concentrations. These UCLs are typically developed for the chosen excursion indicators by analyzing the baseline groundwater quality for a given well field. The UCLs should be set high enough that false positives (false alarms from natural fluctuations in water quality) are not a frequent problem, but not so high that groundwater quality significantly degrades by the time an excursion is identified. Each UCL also must be greater than the baseline concentration for its respective excursion indicator. ASTM D6312 (ASTM International, 1998) and NRC (2003, Section 5.7.8) discuss appropriate statistical methods that can be used to establish UCLs.

The monitoring wells are sampled at least every 2 weeks during well field operations to verify that ISL solutions are contained within the operating well field. NRC (2003, Section 5.7.8) provides basic guidelines for monitoring frequency and response to an excursion detection. As an example, at the Crow Butte ISL uranium recovery facility in Dawes County, Nebraska, baseline water quality was established within the ore zone and in the first aguifer overlying the ore zone prior to uranium recovery. These water quality data are used to determine groundwater monitoring UCLs for five excursion parameters (chloride, sulfate, sodium, conductivity, and alkalinity) (NRC, 1998). The UCLs were calculated as 20 percent above the maximum baseline standards from three samples taken from a well. During well field production, the operator takes samples every 2 weeks from the monitoring wells. A lixiviant excursion is assumed only when two UCLs in any monitoring well are exceeded or if a single UCL at a monitoring well is exceeded by 20 percent. If there is a lixiviant excursion, the operator must notify NRC within 24 hours to institute corrective actions, increase the sampling frequency to weekly, and prepare an excursion report for NRC. If the actions taken in response to the excursion are not effective by the time the 60-day excursion report is submitted, the licensee must stop injecting lixiviant into the well field until aquifer cleanup is complete or provide an increase in surety amount agreeable to NRC that would cover the expected full cost of correcting or cleaning up the excursion (NRC, 1998, 2003). The surety may also be revised to cover the anticipated increase in aquifer restoration costs (NRC, 2003).

8.3.2 Well Field and Pipeline Flow and Pressure Monitoring

The operator typically will monitor injection and production well flow rates to manage the water balance for the entire well field (NRC, 2006). For example, at the proposed Reynolds Ranch expansion for the Smith Ranch/Highlands Uranium Project in Converse County, Wyoming, the operator proposed to monitor the flow rate of each production and injection well by monitoring individual flow meters in each well field header house (NRC, 2006, Section 6). Production well flow rates would be monitored daily and injection well flow rates at least every 3 days.

Additionally, the pressure of each production well and the production trunk line in each well field header house is monitored daily and compared to a maximum surface pressure that is calculated to maintain well integrity. Unexpected losses of pressure may indicate equipment failure, a leak, or a problem with well integrity.

8.4 Ecological Monitoring

Depending on the ecological resources in the area of a facility, the operator may be required to monitor other environmental resources such as plant or animal species.

Ecological monitoring may include surveys of habitat, species counts, or other measures of the health of endangered, threatened, and sensitive species. In addition, surveys may be used to determine whether planned activities are resulting in establishing invasive species populations. Specific survey requirements typically are established through consultations with Federal agencies such as the U.S. Fish and Wildlife Service or State agencies such as the Wyoming Department of Environmental Quality or the New Mexico Environmental Department. Surveys typically cover all phases and areas of planned activity for the life of the project (Energy Metals Corporation, U.S., 2007, Section 6.3). To understand potential impacts on seasonal breeding, timing may be important for some species. For example, in accordance with Wyoming Department of Environmental Quality requirements, Power Resources Inc. conducts a raptor survey in late April or early May of each year to identify any new nests and to address whether known nests are being used (NRC, 2007b). These surveys are conducted to protect against unforeseen conditions where raptors would be nesting in close proximity to operations.

8.5 References

ASTM International. "Standard Guide for Developing Appropriate Statistical Approaches for Ground-Water Detection Monitoring." Designation D6312–98. West Conshohocken, Pennsylvania: ASTM International. 1998.

Council on Environmental Quality. "Aligning National Environmental Policy Act Processes With Environmental Management Systems. A Guide for NEPA and EMS Practitioners." Washington, DC: Council on Environmental Quality. April 2007.

Deutsch, W.J., W.J. Martin, L.E. Eary, and R.J. Serne. NUREG/CR–3709, "Method of Minimizing Ground-Water Contamination From *In-Situ* Leach Uranium Mining." Washington, DC: NRC. 1985.

Energy Metals Corporation, U.S. "Application for USNRC Source Material License Moore Ranch Uranium Project, Campbell County, Wyoming: Environmental Report." Casper, Wyoming: Energy Metals Corporation, U.S. [ADAMS Accession Number: ML072851249] September 2007.

Mackin, P.C., D. Daruwalla, J. Winterle, M. Smith, and D.A. Pickett. NUREG/CR–6733, "A Baseline Risk-Informed Performance-Based Approach for *In-Situ* Leach Uranium Extraction Licensees." Washington, DC: NRC. September 2001.

NRC. "Regulatory Guide 4.15, Quality Assurance for Radiological Monitoring Programs (Normal Operations)—Effluent Streams and the Environment, Revision 2." Washington, DC: NRC. July 2007a.

NRC. "Environmental Assessment for the Construction and Operation of *In-Situ* Leach Satellite SR-2, Amendment No. 12 to Source Materials License No. SUA-1548, Power Resources, Inc.'s Smith Ranch/Highlands Uranium Project Converse County, Wyoming." Source Material License No. SUA-1548. Docket No. 40-8964. Washington, DC: NRC. December 2007b.

NRC. "Environmental Assessment for the Addition of the Reynolds Ranch Mining Area to Power Resources, Inc.'s Smith Ranch/Highlands Uranium Project Converse County, Wyoming." Source Material License No. SUA–1548. Docket No. 40-8964. Washington, DC: NRC. November 2006.

NRC. NUREG–1569, "Standard Review Plan for *In-Situ* Leach Uranium Extraction License Applications—Final Report." Washington, DC: NRC. June 2003.

NRC. "Environmental Assessment for Renewal of Source Material License No. SUA–1534. Crow Butte Resources Incorporated Crow Butte Uranium Project Dawes County, Nebraska." Docket No. 40-8943. Washington, DC: NRC. February 1998.

NRC. NUREG–1508, "Final Environmental Impact Assessment To Construct and Operate the Crownpoint Uranium Solution Mining Project, Crownpoint, New Mexico." Docket No. 40-8968. Washington, DC: NRC. February 1997.

NRC. "Regulatory Guide 4.14, Radiological Effluent and Environmental Monitoring at Uranium Mills, Revision 1." Washington, DC: NRC. 1980.

Staub, W.P., R.E. Williams, F. Anastasi, N.E. Hinkle, J. Osiensky, and D. Rogness. NUREG/CR–3967, "An Analysis of Excursions at Selected *In-Situ* Uranium Mines in Wyoming and Texas." Washington, DC: NRC. 1986.

WDEQ. "In-Situ Mining Permit Application Requirements Handbook. Application Content Requirements—Adjudication and Baseline Information." Cheyenne, Wyoming: WDEQ, Land Quality Division. March 2007.

9 CONSULTATIONS

This Generic Environmental Impact Statement (GEIS) takes a programmatic look at the environmental impacts of *in-situ* leach (ISL) uranium mining on the four regions described in Section 1.4. For the purpose of the GEIS, the programmatic aspects of the consultation process are described in this chapter. Each site-specific review would include its own consultation process with the relevant agencies including, but not limited to, state and tribal historic preservation offices [National Historic Preservation Act, Section 106 (NHPA)], U.S. Fish and Wildlife Service (Endangered Species Act, Section 7), and tribal consultations with appropriate Native American communities. The U.S. Nuclear Regulatory Commission (NRC) consultation process stresses early interaction in an effort to gather information to prepare an environmental review. In particular, 10 CFR 51.28(a)(3–5) specifically requires NRC to extend invitations to affected (state, local, tribal and federal government) agencies to meet as part of the scoping process for an environmental impact statement (EIS).

National Historic Preservation Act

NRC uses its National Environmental Policy Act (NEPA) process to coordinate Section 106 of the NHPA, which requires that Federal agencies "take into account the effects of their undertakings on historic properties and afford the Council (Advisory Council on Historic Preservation) a reasonable opportunity to comment on such undertakings." Typically, NRC licensing actions can be defined as undertakings based on 36 CFR 800.16(y) because the proposed actions consider applications and licensing amendments that require a "Federal permit, license or approval." NRC performs an evaluation of the proposed action to determine whether the activity has a potential to effect historic properties. NRC initiates consultation with relevant agencies including the State Historic Preservation Office and/or the Tribal Historic Preservation Office, reports the conclusions of its evaluation, and seeks concurrence with its findings.

For the purpose of the GEIS, the proposed action considers the impact of construction, operation, aquifer restoration, and decommissioning of ISL facilities in four geographical regions in the western United States. Because the actual undertaking would occur when site-specific applications are submitted, the GEIS does not include Section 106 consultations. The site specific environmental reviews would identify the area of potential effect and list any historic properties. Each site-specific environmental review would address the potential impact of the proposed action on the appropriate historic properties.

Threatened and Endangered Species

The Endangered Species Act (ESA) of 1973 was enacted to protect critically imperiled species from extinction as a "consequence of economic growth and development untendered by adequate concern and conservation." Section 7 of the ESA directs all federal agencies to use their existing authorities to conserve threatened and endangered species and, in consultation with the U.S. Fish and Wildlife Service, to ensure that their actions do not jeopardize listed species or destroy or adversely modify critical habitat. Section 7 applies to management of federal lands as well as other federal actions that may affect listed species, such as federal approval of private activities through the issuance of federal permits, licenses, or other actions.

NRC uses its NEPA process to coordinate Section 7 consultations under the ESA. The staff perform an evaluation to identify the action area, determine whether listed species or critical habitat exist in the action area, and evaluate the potential impact on any listed species or critical habitat. For the purpose of this GEIS, the NRC staff identified endangered species in the four regions. Consultation would be initiated with the U.S. Fish and Wildlife Service to determine whether critical habitats exist for species of concern on a site-specific basis. At the end of the consultation process, NRC would notify the U.S. Fish and Wildlife Service of its conclusions and document them in the site-specific environmental analysis.

State Consultation

As a part of the environmental review process, NRC consults with the affected states and solicits comments on the environmental impact of the proposed action. This consultation is designed to address issues raised by state and local agencies and to reduce any duplication of effort in complying with federal, state, and local environmental requirements. Because the GEIS contains a regional, programmatic evaluation, state consultations are not reported, as these would be would be conducted during the site-specific review. As discussed in Section 1.8, NRC will use the GEIS to prepare a supplemental EIS for new license applications and a site-specific environmental assessment or EIS for applications to renew or amend existing ISL licenses. As part of the environmental review for new applications, NRC may conduct a scoping process, consistent with its regulations at 10 CFR 51.26(d), 51.28, and 51.29. During the scoping and information gathering process for a site-specific environmental review, the NRC staff typically contacts appropriate state and local agencies for initial, informal discussion about the proposed action and potential impacts. Additionally, NRC will publish the draft supplemental EIS for public comment in accordance with 10 CFR 51.73 and 51.117. Part of the NRC state consultations would include informing affected state governments when these opportunities for involvement are initiated for specific licensing actions. The NRC staff will address state comments received on the draft supplemental EIS prior to making a final licensing decision. For site-specific reviews of license renewal or amendment requests that result in the preparation of an environmental assessment, NRC would submit a copy of the draft environmental assessment to the state for review and comment.

Tribal Consultation

NRC consults with the affected tribes as part of carrying out the intent behind Executive Order 13175 "Consultation and Coordination With Indian Tribal Governments" and requirements under 10 CFR 51.28(a)(5). Formal and informal consultations through the environmental review process can fulfill these responsibilities. Because the GEIS contains a regional, programmatic evaluation, tribal consultations are not reported, as these would be conducted during the site-specific review. NRC would consult with affected tribal governments to consider topics of concern regarding specific ISL proposals, including potentially affected places of cultural significance, land disturbance, health, and groundwater use and restoration. NRC has developed a strategy for outreach to Native American tribes to facilitate an open dialogue with tribes on topics of mutual interest regarding future uranium recovery licensing actions. This strategy is available on the NRC website (NRC, 2000). As discussed in Section 1.8, NRC will use the GEIS to prepare a supplemental EIS for new license applications and a site-specific environmental assessment or EIS for applications to renew or amend existing ISL licenses. As part of the environmental review for new applications, NRC may conduct a scoping process, consistent with its regulations at 10 CFR 51.26(d), 51.28, and 51.29. Additionally,

NRC will publish the draft supplemental EIS for public comment in accordance with 10 CFR 51.73 and 51.117. Part of the NRC tribal consultations would include informing affected tribal governments when these opportunities for involvement are initiated for specific licensing actions. The NRC staff will address public comments received on the draft supplemental EIS prior to making a final licensing decision. For site-specific review of license renewal or amendment requests that result in the preparation of an environmental assessment, NRC would submit a copy of the draft environmental assessment to affected tribes for review and comment.

For applications for new ISL facilities that have potential cultural and resource impacts on the Navajo Nation, NRC has committed to consultations with the Navajo Nation, through the Navajo Nation Department of Justice (U.S. Department of the Interior, 2008). These consultations for site-specific environmental reviews would take into account topics identified by NRC and the tribal agencies (e.g., Navajo Nation Environmental Protection Agency).

Reference

U.S. Department of the Interior. "Health and Environmental Impacts of Uranium Contamination in the Navajo Nation: Five-Year Plan." Washington, DC: U.S. Department of the Interior, Bureau of Indian Affairs. 2008.

10 SUMMARY OF ENVIRONMENTAL CONSEQUENCES

The environmental resources in the four geographic regions where current *in-situ* leach (ISL) facilities are located and where future ISL facilities may be located are discussed in Chapter 3. Based on the description of the ISL process and the historical information on ISL facilities in Chapter 2, the potential environmental impacts are described and analyzed in Chapter 4. In this chapter, for each of the four uranium milling regions considered within this GEIS, the potential environmental impacts are summarized for construction, operation, aquifer restoration, and decommissioning at an ISL facility for each environmental resource.

In the Impact Findings column of the table that follows, the impacts are categorized by the significance levels described in Chapter 1:

- **SMALL**—The environmental effects would not be detectable or are so minor that they would neither destabilize nor noticeably alter any important attribute of the resource considered.
- **MODERATE**—The environmental effects would be sufficient to alter noticeably, but not destabilize, important attributes of the resource considered.
- **LARGE**—The environmental effects would be clearly noticeable and are sufficient to destabilize important attributes of the resource considered.

As described in Section 1.8, for each new ISL license application, NRC will conduct an independent site-specific environmental review to meet its responsibilities under the National Environmental Policy Act, drawing on the information and conclusions in the GEIS as appropriate.

Summary	•
9	
t Environmental	1
≦ro	
ňm	
ent	
_	
Con)
seq	
uer	
Consequences	

		Table 10-1. Summary of Impacts for the Wyoming West Uranium Milling Region
Topic/ Resource	GEIS Section	Impact Findings
Land Use	4.2.1	CONSTRUCTION—Land use impacts could occur from land disturbances (including alterations of ecological cultural or historic resources) and access restrictions (including limitations of other mineral extraction activities, grazing activities, or recreational activities). Land disturbances during construction would be temporary and limited to small areas within permitted areas. Well sites, staging areas, and trenches would be reseeded and restored. Unpaved access roads would remain in use until decommissioning. Competing access to mineral rights could be either delayed for the duration of the <i>in-situ</i> leaching (ISL) project or be intermixed with ISL operations (e.g., oil and gas exploration). Changes to land use access including grazing restrictions and impacts on recreational activities would be limited due to the small size of restricted areas, temporary nature of restrictions, and availability of other land for these activities. Ecological, historical, and cultural resources could be affected, but would be protected by careful planning and surveying to help identify resources and avoid or mitigate impacts. For all land use aspects except ecological, historical and cultural resources, the potential impacts would be SMALL. Due to the potential for unidentified resources to be altered or destroyed during excavation, drilling, and grading, the potential impacts to ecological, historical or cultural resources would be SMALL to LARGE, depending on local conditions. OPERATION—The types of land use impacts for operational activities would be expected to be similar to construction impacts regarding access restrictions because the infrastructure would be in place. Additional land disturbances would not occur from conducting operational activities. Because access restriction and land disturbance related impacts would be similar to, or less than, expected for construction, the overall potential impacts to land use from operational activities would be expected to be SMALL. AQUIFER RESTORATION—Due to the use of the same i

	Т	able 10-1. Summary of Impacts for the Wyoming West Uranium Milling Region (continued)
Topic/	GEIS	
Resource	Section	Impact Findings
	Section 4.2.2	Impact Findings CONSTRUCTION—Low magnitude traffic generated by ISL construction relative to local traffic counts would not significantly increase traffic or accidents on many of the roads in the region. Existing low traffic roads could be MODERATEly impacted by the additional worker commuting traffic during periods of peak employment. The potential impact would be more pronounced in areas with lower traffic counts. MODERATE dust, noise, and incidental wildlife or livestock kill impacts would be possible on, or near, site access roads (dust in particular for unpaved access roads)—SMALL to MODERATE. OPERATION—Low magnitude traffic relative to local traffic counts on most roads would not significantly increase traffic, or accidents. Existing low traffic roads could be moderately impacted by commuting traffic during periods of peak employment including dust, noise, and possible incidental wildlife or livestock kill impacts on, or near site access roads. High consequences would be possible for a severe accident involving transportation of hazardous chemicals in a populated area. However, the probability of such accidents occurring would be low, owing to the limited number of shipments, comprehensive regulatory controls, and use of best management practices. For radioactive material shipments (yellowcake product, ion exchange resins, waste materials), compliance with transportation regulations would limit radiological risk for normal operations. Consequently, there is low radiological risk associated with accident conditions. Emergency response protocols would help mitigate long-term consequences of severe accidents involving release of uranium—SMALL to MODERATE. AQUIFER RESTORATION—The magnitude of transportation activities would be lower than for construction and operations, with the exception of workforce commuting which could have moderate impacts on, or in the vicinity of, existing low traffic roads—SMALL to MODERATE.
		DECOMMISSIONING—The types of transportation activities and therefore types of impacts would be similar to those discussed for construction and operations except the magnitude of transportation activities (e.g., number and types of waste and supply shipments, no yellowcake shipments) from decommissioning could be lower than for operations. Accident risks would be bounded by operations yellowcake transportation risk estimates—SMALL.

Summar
of
Environmental
Consequences

	Т	able 10-1. Summary of Impacts for the Wyoming West Uranium Milling Region (continued)
Topic/	GEIS	
Resource	Section	Impact Findings
Geology and	4.2.3	CONSTRUCTION—Disturbance to soil would occur from construction (clearing, excavation, drilling, trenching, road construction). However, such disturbances would be temporary and SMALL (approx. 15 percent of the total site area), and potential impacts would be mitigated by using best management practices. A large portion of the well fields, trenches, and access roads would be restored and reseeded after construction. Excavated soils would be stockpiled, seeded, and stored onsite until needed for reclamation fill. No impacts to subsurface geological strata are likely—SMALL. OPERATION—Temporary contamination or alteration of soils would be likely from operational leaks and spills and possible from transportation, use of evaporation ponds, or land application of treated waste water. However, detection and response techniques, monitoring of treated waste water, and eventual survey and decommissioning of all potentially impacted soils,
Soils	4.2.0	would limit the magnitude of overall impacts to soils—SMALL. AQUIFER RESTORATION—Impacts to geology and soils from aquifer restoration activities would be similar to impacts from operations due to use of the same infrastructure and similar activities conducted (e.g., well field operation, transfer lines, waste water treatment and disposal)—SMALL. DECOMMISSIONING—Impacts to geology and soils from decommissioning would be similar to impacts from construction. Activities to cleanup, recontour, and reclaim disturbed lands during decommissioning would mitigate long-term impacts to soils—SMALL.

	Т	able 10-1. Summary of Impacts for the Wyoming West Uranium Milling Region (continued)
Topic/ Resource	GEIS Section	Impact Findings
Surface Waters	4.2.4.1	CONSTRUCTION—Impacts to surface waters and related habitats from construction (road crossings, filling, erosion, runoff, spills or leaks of fuels and lubricants for construction equipment) would be mitigated through proper planning, design, construction methods, and best management practices. Some impacts directly related to the construction activities would be temporary and limited to the duration of the construction period. U.S. Army Corps of Engineers permits may be required when filling and crossing wetlands. Temporary changes to spring and stream flows from grading and changes in topography and natural drainage patterns could be mitigated through best management practices, or restored after the construction phase. Incidental spills of drilling fluids into local streams would be small and temporary, due to the implementation of mitigation measures. Impacts from construction of roads, parking areas, and buildings on recharge to shallow aquifers would be small, owing to the limited area of impervious surfaces proposed. Infiltration of drilling fluids into the local aquifer would be small, temporary, and localized to a few feet around boreholes—SMALL. OPERATION—Impacts from storm water runoff or direct discharge of process waters (brine reject from reverse osmosis, or spent eluants from an ion exchange system) to surface waters would be regulated by the Wyoming Department of Environmental Quality through the Wyoming Pollutant Discharge Elimination System permit. Expansion of facilities or pipelines during operations would generate impacts similar to construction. Because the aquifers containing uranium ore-bodies would have a weak, if any, connection to local surface water features, such as streams and springs, the impacts of excess net groundwater extraction from local surface water bodies would be SMALL to MODERATE, depending on site-specific characteristics. AQUIFER RESTORATION—Impacts from aquifer restoration would be similar to impacts from operations due to use of in-place infrastructure and similar ac

	Tal	ble 10-1. Summary of Impacts for the Wyoming West Uranium Milling Region (continued)
Topic/ Resource	GEIS Section	Impact Findings
		CONSTRUCTION—Water use impacts would be limited by the small volumes of groundwater used for routine activities such as dust suppression, mixing cements, and drilling support over short and intermittent periods. Contamination of groundwater from construction activities would be mitigated by use of best management practices—SMALL.
Water— Groundwater	4.2.4.2	OPERATION—Potential impacts to shallow aquifers can occur from leaks or spills from surface facilities and equipment. Shallow aquifers are important sources of drinking water in some areas of the Wyoming West Uranium Milling Region. Potential impacts to the ore-bearing and surrounding aquifers include consumptive water use and degradation of water quality (from normal production activities, off-normal excursion events, and deep well injection disposal practices). Consumptive use impacts from withdrawal of groundwater would occur because only 1 to 3 percent of pumped groundwater is not returned to the aquifer (e.g., process bleed). The amount of water lost could be reduced substantially by available treatment methods (e.g., reverse osmosis, brine concentration). Effects of water withdrawal on surface water would be SMALL as the ore zone normally occurs in a confined aquifer. Estimated drawdown effects vary depending on site conditions and water treatment technology applied. Excursions of lixiviant and mobilized chemical constituents could occur from failure of well seals or other operational conditions that result in incomplete recovery of lixiviant. Well seal related excursions would be detected by the groundwater monitoring system and periodic well mechanical integrity testing, and impacts would be mitigated during operation or aquifer restoration. Other excursions could result in plumes of mobilized uranium and heavy metals extending beyond the mineralization zone. The magnitude of potential impacts from vertical excursions would vary depending on site-specific conditions. To reduce the likelihood and consequences of potential excursions at ISL facilities, NRC requires licensees to take preventative measures prior to starting operations including well tests, monitoring, and development of procedures that include excursion response measures and reporting requirements. Impacts associated with alterations of ore body aquifer chemistry would be SMALL because the aquifer would: (1) be confined, (2) not be a potent
		AQUIFER RESTORATION—Potential impacts include consumptive use and potential deep disposal of brine slurries after reverse osmosis, if applicable. The volume of water removed from the aquifer and related impacts would be dependent on site-specific conditions and the type of water treatment technology used at the facility. Groundwater Consumptive use during aquifer restoration could be greater than during ISL operation, if groundwater sweep is implemented during aquifer restoration in which pumped water is not recirculated. Potential environmental impacts associated with water consumption during aquifer restorations are determined by: (1) the restoration techniques chosen, (2) the volume of water to be used, (3) the severity and extent of the contamination, and (4) the current and future use of the production and surrounding aquifers near the ISL facility or at the regional scale—SMALL to MODERATE, depending on site-specific conditions.
		DECOMMISSIONING—Potential impacts from decommissioning would be similar to construction (water use, spills) with an additional potential to mobilize contaminants during demolition and cleanup activities. Contamination of groundwater from decommissioning activities would be mitigated by implementation of an NRC-approved decommissioning plan and use of best management practices—SMALL.

-	_
`	٢
(x

	Tal	ble 10-1. Summary of Impacts for the Wyoming West Uranium Milling Region (continued)
Topic/ Resource	GEIS Section	Impact Findings
Ecology— Terrestrial	4.2.5.1	CONSTRUCTION—Potential terrestrial ecology impacts would include the removal of vegetation from well fields and the milling site, the modification of existing vegetative communities, the loss of sensitive plants and habitats from clearing and grading, and the potential spread of invasive species and noxious weed populations. These impacts would be temporary because restoration and reseeding occur rapidly after the end of construction. Introduction of invasive species and noxious weeds would be possible but could be mitigated by restoration and reseeding after construction. Shrub and tree removal would have a longer restoration period. Wildlife habitat fragmentation, temporary displacement of animal species, and direct or indirect mortalities is possible. Implementation of wildlife surveys and mitigation measures following established guidelines would limit these impacts. The magnitude of impacts depends on whether a new facility is being licensed or an existing facility is being extended—SMALL to MODERATE, depending on site-specific conditions. OPERATION—Habitat could be altered by operations (fencing, traffic, noise), and individual takes could occur due to conflicts between species habitat and operations. Access to crucial wintering habitat and water could be limited by fencing. The Wyoming Game and Fish Department specifies fencing construction techniques to minimize impediments to big game movement. Migratory birds could be affected by exposure to constituents in evaporation ponds, but perimeter fencing, would limit impacts. Temporary contamination or alteration of soils would be from operational leaks and spills and possibly from transportation or land application of treated waste water. However, detection and response techniques, and eventual survey and decommissioning of all potentially impacted soils, would limit the magnitude of overall impacts to terrestrial ecology. Mitigation measures, such as perimeter fencing, netting, alternative sites, and timing stipulations would result from leaks and spills,

Summary of Environment	
of Environmental Consequences	

	Т	able 10-1. Summary of Impacts for the Wyoming West Uranium Milling Region (continued)
Topic/ Resource	GEIS Section	Impact Findings
Ecology— Aquatic	Section 4.2.5.2	CONSTRUCTION—Clearing and grading activities associated with construction could result in a temporary increase in sediment load in local streams, but aquatic species would recover quickly as sediment load decreases. Clearing of riparian vegetation could affect light and temperature of water. Construction impacts to wetlands would be identified and managed through U.S. Army Corps of Engineers permits, as appropriate. Construction impacts to surface waters and aquatic species would be temporary and mitigated by best management practices—SMALL. OPERATION—Impacts could result from spills or releases into surface water. Impacts would be minimized by spill prevention, identification and response programs, and National Pollutant Discharge Elimination System (NPDES) permit requirements—SMALL. AQUIFER RESTORATION—Activities would use existing (in-place) infrastructure, and impacts could result from spills or releases of untreated groundwater. Impacts would be minimized by spill prevention, identification, and response programs, and NPDES permit requirements—SMALL. DECOMMISSIONING—Decommissioning and reclamation activities could result in temporary increases in sediment load in
		local streams, but aquatic species would recover quickly as sediment load decreases. With completion of decommissioning, revegetation, and re-contouring, habitat would be reestablished and impacts would, therefore, be limited—SMALL.

-	_	_
(5
	ı	
ō	_	5

Topic/ Resource GEIS Section Impact Findings	Т	Table 10-1	. Summary of Impacts for the Wyoming West Uranium Milling Region (continued)
CONSTRUCTION—Numerous threatened and endangered species and State Species of Concern are located in the region. Small fragmentation of habitats could occur in addition to potential habitat loss. The magnitude of impacts depends on the size of a new facility or extension to an existing facility and the amount of land disturbance. Inventory threatened or endangered species would be developed during site-specific reviews to identify unique or special habits and Endangered Species Act consultations conducted with the U.S. Fish and Wildlife Service would assist in identifying potential impacts—SMALL to LARGE—depending on site-specific habitat and presence of threatened or endangered species. OPERATION—Impacts could result from individual takes due to conflicts with operations. Small fragmentation of habit could occur, in addition to potential habitat loss. The magnitude of impacts would depend on the size of a new facility extension to an existing facility and the amount of land disturbance. Impacts could potentially result from spills or permitted effluents, but would be minimized by spill prevention measures, identification and response programs, and NPDES permit requirements. Inventory of threatened or endangered species developed during site-specific reviews would identify unique or special habitats, and Endangered Species Act consultations conducted with the U.S. Fish an Wildlife Service would assist in identifying potential impacts—SMALL to LARGE—depending on site-specific habitat and presence of threatened or endangered species.	Topic/ GEIS	Topic/ GEIS	
region. Small fragmentation of habitats could occur in addition to potential habitat loss. The magnitude of impacts depends on the size of a new facility or extension to an existing facility and the amount of land disturbance. Inventor, threatened or endangered species would be developed during site-specific reviews to identify unique or special habitat and Endangered Species Act consultations conducted with the U.S. Fish and Wildlife Service would assist in identifying potential impacts—SMALL to LARGE—depending on site-specific habitat and presence of threatened or endangered species. OPERATION—Impacts could result from individual takes due to conflicts with operations. Small fragmentation of habitat could occur, in addition to potential habitat loss. The magnitude of impacts would depend on the size of a new facility extension to an existing facility and the amount of land disturbance. Impacts could potentially result from spills or permitted effluents, but would be minimized by spill prevention measures, identification and response programs, and NPDES permit requirements. Inventory of threatened or endangered species developed during site-specific reviews would identify unique or special habitats, and Endangered Species Act consultations conducted with the U.S. Fish an Wildlife Service would assist in identifying potential impacts—SMALL to LARGE—depending on site-specific habitat a presence of threatened or endangered species.	Resource Section	Resource Section	Impact Findings
AQUIFER RESTORATION—Impacts could result from individual takes due to conflicts with aquifer restoration activities (equipment, traffic). Existing (in-place) infrastructure would be used during aquifer restoration, so additional land-disturbing activities and habitat fragmentation would not be anticipated. Impacts may result from spills or releas treated or untreated groundwater, but impacts would be minimized by spill prevention measures, identification, and response programs, and NPDES permit requirements. Inventory of threatened or endangered species would be developed during site-specific reviews to identify unique or special habitats, and Endangered Species Act consultation with the U.S. Fish and Wildlife Service would assist in identifying potential impacts—SMALL. DECOMMISSIONING—Impacts resulting from individual takes could occur due to conflicts with decommissioning activities (equipment, traffic). Temporary land disturbance would occur as structures are demolished and removed at the ground surface is re-contoured. Inventory of threatened or endangered species developed during site-specific environmental review of the decommissioning plan would identify unique or special habitats, and Endangered Species consultations with the U.S. Fish and Wildlife Service would assist in identifying potential impacts. With completion of	Ecology— Threatened or Endangered 4.2.5.3	Ecology— Threatened or Endangered Species 4.2.5.3 AQUII (equip land-c treate respondevelowith the green viros	STRUCTION—Numerous threatened and endangered species and State Species of Concern are located in the n. Small fragmentation of habitats could occur in addition to potential habitat loss. The magnitude of impacts nds on the size of a new facility or extension to an existing facility and the amount of land disturbance. Inventory of tened or endangered species would be developed during site-specific reviews to identify unique or special habitats, andangered Species Act consultations conducted with the U.S. Fish and Wildlife Service would assist in identifying tial impacts—SMALL to LARGE—depending on site-specific habitat and presence of threatened or endangered es. RATION—Impacts could result from individual takes due to conflicts with operations. Small fragmentation of habita occur, in addition to potential habitat loss. The magnitude of impacts would depend on the size of a new facility or sion to an existing facility and the amount of land disturbance. Impacts could potentially result from spills or titted effluents, but would be minimized by spill prevention measures, identification and response programs, and ES permit requirements. Inventory of threatened or endangered species developed during site-specific reviews identify unique or special habitats, and Endangered Species Act consultations conducted with the U.S. Fish and fe Service would assist in identifying potential impacts—SMALL to LARGE—depending on site-specific habitat and ince of threatened or endangered species. FER RESTORATION—Impacts could result from individual takes due to conflicts with aquifer restoration activities of threatened or endangered species. FER RESTORATION—Impacts could result from individual takes due to conflicts with aquifer restoration activities of threatened groundwater, but impacts would be minimized by spill prevention measures, identification, and one programs, and NPDES permit requirements. Inventory of threatened or endangered species Act consultations the U.S. Fish and Wildlife Service would assist in identifying p

		ble 10-1. Summary of Impacts for the Wyoming West Uranium Milling Region (continued)
Topic/ Resource	GEIS Section	Impact Findings
		CONSTRUCTION—Fugitive dust and combustion (vehicle and diesel) emissions during land disturbing activities associated with construction would be small, short-term, and reduced through best management practices (e.g., dust supression). For example, estimated fugitive dust emissions during ISc construction are less than 2 percent of the National Ambient Air Quality Standards (NAAQS) for PM _{2.5} and less than 1 percent for PM ₁₀ . For NAAQS attainment areas such as the Wyoming West Uranium Milling Region, nonradiological air quality impacts would be SMALL, and there are no Prevention of Significant Deterioration (PSD) Class I areas in the Wyoming West Uranium Milling Region. Furthermore, if impacts were initially assessed at a higher significance level, permit requirements would impose conditions or mitigation measures to reduce impacts—SMALL. OPERATION—Radiological impacts can result from dust releases from drying of lixiviant pipeline spills, radon releases from well system relief valves, resin transfer, or elution, and gaseous/particulate emissions from yellowcake dryers. Only small amounts of low dose materials would be released based on operational controls and rapid response to spills. Required spill prevention, control, and response procedures would be used to minimize impacts from spills. High Efficiency Particulate Air (HEPA) filters and vacuum dryer designs reduce particulate emissions from operations, and ventilation reduces radon buildup during operations. Compliance with the NRC-required radiation monitoring program would ensure releases are within regulatory limits. Other potential nonradiological emissions during operations include fugitive dust and fuel from equipment, maintenance, transport trucks, and other vehicles. For NAAQS attainment areas such as the Wyoming West Uranium Milling Region, nonradiological air quality impacts would be SMALL, and there are no PSD Class I areas in the Wyoming West Uranium Milling Region. Furthermore, if impacts were initially assessed at a higher significance lev
		similar to, or less than, operations. For NAAQS attainment areas such as the Wyoming West Uranium Milling Region, nonradiological air quality impacts would be SMALL, and there are no PSD Class I areas in the Wyoming West Uranium Milling Region. Furthermore, if impacts were initially assessed at a higher significance level, permit requirements would impose conditions or mitigation measures to reduce impacts—SMALL.
		DECOMMISSIONING—Fugitive dust and combustion (vehicle and diesel) emissions during land-disturbing activities associated with decommissioning would be similar to, or less than, associated with construction, short-term, and reduced through best management practices (e.g., dust suppression). These potential impacts would decrease as decommissioning and reclamation of disturbed areas are completed. For NAAQS attainment areas such as the Wyoming West Uranium Milling Region, nonradiological air quality impacts would be SMALL, and there are no PSD Class I areas in the Wyoming West Uranium Milling Region. Furthermore, if impacts were initially assessed at a higher significance level, permit requirements would impose conditions or mitigation measures to reduce impacts—SMALL.

		able 10-1. Summary of Impacts for the Wyoming West Uranium Milling Region (continued)
Topic/	GEIS	
Resource	Section	Impact Findings
Noise	4.2.7	CONSTRUCTION—Noise generated during construction would be noticeable in proximity to operating equipment, but would be temporary (typically) daytime only). Administrative and engineering controls would be used to maintain noise levels in work areas below Occupational Health and Safety Administration (OSHA) regulatory limits and be mitigated by use of personal hearing protection. Traffic noise during construction (commuting workers, truck shipments to and from the facility, and construction equipment such as trucks, bulldozers, compressors) would be localized, limited to highways in the vicinity of the site, access roads within the site, and roads in well fields. Relative increases in traffic levels would be small for larger roads, but may be moderate for lightly traveled rural roads through less populated communities. Noise may adversely effect on wildlife habitat and their reproductive success in the immediate vicinity of construction activities. Noise levels decrease geometrically with distance, and at distances more than 300 m [1,000 ft], ambient noise levels return to background levels. Wildlife generally avoid construction noise areas. The two uranium districts within the Wyoming West Uranium Milling Region are generally more than 16 km [10 mi] from the closest community—SMALL to MODERATE. OPERATION—Noise-generating activities in the central uranium processing facility would be indoors, minimizing offsite sound levels. Well field equipment (e.g., pumps, compressors) would also be expected to be contained within structures (e.g., header houses, satellite facilities), minimizing sound levels to offsite receptors. Administrative and engineering controls would be used to maintain noise levels in work areas below OSHA regulatory limits, and be mitigated by use of personal hearing protection. Traffic noise from community workers, truck shipments to and from the facility, and facility equipment would be localized, limited to highways in the vicinity of the site, access roads within the site, and roads in well fi

	Т	able 10-1. Summary of Impacts for the Wyoming West Uranium Milling Region (continued)
Topic/	GEIS	
Resource	Section	Impact Findings
Historical and Cultural	4.2.8	CONSTRUCTION—Potential impacts during ISL facility construction could include loss of, or damage and temporary restrictions on access to, historical, cultural, and archaeological resources. The eligibility evaluation of cultural resources for listing in the National Register of Historic Places (NRHP) under criteria in 36 CFR 60.4(a)—(d) and/or as Traditional Cultural Properties (TCPs) is conducted as part of the site-specific review and NRC licensing procedures undertaken during the National Environmental Policy Act (NEPA) review process. The evaluation of impacts to any historic properties designated as TCPs and tribal consultations regarding cultural resources and TCPs also occur during the site-specific licensing application and review process. Consultations to determine whether significant cultural resources would be avoided or mitigated occurs during consultations with State Historic Preservation Offices (SHPOs), other governmental agencies, and Native American tribes, including Tribal Historic Preservation Offices (THPOs), as part of the site-specific review process. Additionally, as needed, the NRC license applicant would be expected to be required, under conditions in its NRC license, to adhere to procedures regarding the discovery of previously undocumented cultural resources during initial construction. These procedures typically require the licensee to stop work and to notify the appropriate federal, tribal, and state agencies with regard to mitigation measures—SMALL or MODERATE to LARGE, depending on site-specific conditions. OPERATION—Because less land disturbance occurs during the operations phase, potential impacts to historical, cultural, and archaeological resources would be less than during construction. Conditions in the NRC license requiring adherence to procedures typically require the licensee to stop work and to notify the appropriate federal, tribal, and state agencies with regard to mitigation measures—SMALL or MODERATE to LARGE depending on site-specific conditions. DECOMMISSIONING—B

_	
0	
느	
\overline{A}	

Table 10-1. Summary of Impacts for the Wyoming West Uranium Milling Region (continued)						
Topic/ Resource	GEIS Section	Impact Findings				
Visual and Scenic	4.2.9	CONSTRUCTION—Visual impacts result from equipment (drill rig masts, cranes), dust/diesel emissions from construction equipment, and hillside and roadside cuts. Most of the Wyoming West Uranium Milling Region is classified as Visual Resource Management (VRM) Class II through IV, and no VRM Class I or PSD Class I areas are located in the region. Most potential visual impacts during construction would be temporary as equipment is moved, and would be mitigated by implementing best management practices (e.g., dust suppression). Because of the generally rolling topography of the region, most visual impacts during construction would not be expected to be visible from more than about 1 km [0.6 mi]. The two uranium districts in the region are located more than 16 km [10 mi] from the closest VRM Class II area, and the visual impacts associated with ISL construction would be consistent with the predominant VRM Class III and IV—SMALL. OPERATION—Visual impacts during operations would be expected to be less than those associated with construction. Most of the well field surface infrastructure has a low profile, and most piping and cables would be buried. The tallest structures would include the central uranium processing facility (10 m [30 ft]) and power lines (6 m [20 ft]). Because of the generally rolling topography of the region, most visual impacts during operations would not be visible from more than about 1 km [0.6 mi]. Irregular layout of well field surface structures such as wellhead protection and header houses would reduce visual contrast. Best management practices, design (e.g., painting buildings), and landscaping techniques would be used to mitigate potential visual impact. The two uranium districts in the region are located more than 16 km [10 mi] from the closest VRM Class II area, and the visual impacts associated with ISL construction would be consistent with the predominant VRM Class III and IV—SMALL. DECOMMISSIONING—Because similar equipment would be used and activities conducted, potential visual impact				

7	$\overline{}$
`	~
-	_
(ກ

	Table 10-1. Summary of Impacts for the Wyoming West Uranium Milling Region (continued)					
Topic/	Topic/ GEIS					
Resource Section Impact Findings						
Public and Occupational Health and Safety	4.2.11	CONSTRUCTION—Worker safety would be addressed by standard construction safety practices. Fugitive dust would result from construction activities and vehicle traffic, but would likely be of short duration, and not result in a radiological dose. Diesel emissions would not be expected to be a concern for worker or public health, because the releases would be of short duration and are readily dispersed into the atmosphere—SMALL. OPERATION—Potential occupational radiological impacts from normal operations would be caused primarily by exposure to radon gas from the well field, ion-exchange resin transfer operations, and venting during processing activities. Workers would also be exposed to airborne uranium particulates from dryer operations and maintenance activities. Potential public exposures to radiation would occur from the same radon releases and uranium particulate releases (i.e., from facilities without vacuum dryer technology). Both worker and public radiological exposures would be limited by NRC regulations at 10 CFR Part 20 which require licensees to implement an NRC-approved radiation monitoring and protection program. (Measured and calculated doses for workers and the public are commonly a fraction of regulated limits.) Nonradiological worker safety matters would be addressed through commonly applied occupational health and safety regulations and practices. Radiological accident risks could involve processing equipment failures leading to yellowcake slurry spills, or radon gas or uranium particulate releases. Consequences of accidents to workers and the public are generally low, with the exception of a dryer explosion, which could result in worker dose above NRC limits. The likelihood of such an accident would be low, and therefore, the risk would also be low. Potential nonradiological accidents impacts include high-consequence chemical release events (e.g., ammonia) for both workers and nearby populations. The likelihood of such release events would be low, based on historical operating experience at NR				

	. Summary of Impacts for the Wyoming West Uranium Milling Region (continued)
Topic/ GEIS	
Resource Section CONS fields a OPER eluant washed would osmos surface addres liquid value site-sp survey dispos general distance estimates AQUIF dispos operate condition restoral removes dispos management DECO evapo A precapacitics of the condition of the cond	Impact Findings STRUCTION—The relatively small scale of construction activities (Section 2.3) and incremental development of well at ISL facilities would generate low volumes of construction waste—SMALL. AATION—Operational wastes primarily result from liquid waste streams including process bleed, flushing of depleted to to limit impurities, resin transfer wash, filter washing, uranium precipitation process wastes (brine), and plant down water. State permitting actions, NRC license conditions, and NRC inspections ensure the proper practices be used to comply with safety requirements to protect workers and the public. Waste treatment such as reverse is and radon settling would help in segregating wastes and minimizing disposal volumes. Potential impacts from be discharge and deep well injection would be limited by the applicable permitting processes. NRC regulations as constructing, operating, and monitoring for leakage from evaporation ponds used to store and reduce volumes of wastes. Potential impacts from land application of treated wastewater would be addressed by NRC review of exaction waste disposal impacts from land application of treated wastewater would be addressed by NRC review of pecific conditions prior to approval, routine monitoring, and inclusion of irrigated land areas in decommissioning ys. Offsite waste disposal impacts would be SMALL for radioactive wastes as a result of required preoperational sal agreements. Impacts for hazardous and municipal waste would be SMALL due to the volume of wastes ated. For remote areas with limited available disposal capacity, such wastes may need to be shipped greater cost to facilities that have capacity. However, the volume of wastes generated, and magnitude of the shipments are ated to be low—SMALL. FER RESTORATION—Waste management activities during aquifer restoration would utilize the same treatment and sal options implemented for operations. Therefore, impacts associated with aquifer restoration would be similar to tional impacts. While the amount of was

	_
(5
	<u>'</u>
Č	သ

Table 10-2. Summary of Impacts for the Wyoming East Uranium Milling Region		
Topic/ Resource	GEIS Section	Impact Findings
Land Use	4.3.1	CONSTRUCTION—Land use impacts could occur from land disturbances (including alterations of ecological cultural or historic resources) and access restrictions (including limitations of other mineral extraction activities, grazing activities, or recreational activities). A higher percentage of private land ownership occurs in this region than in the Wyoming West Uranium Milling Region, and could increase the potential for land use conflicts with private land owners. Land disturbances during construction would be temporary and limited to small areas within permitted site. Well sites, staging areas, and trenches would be reseeded and restored, but unpaved access roads would remain in use until decommissioning is complete. Competing access to mineral rights could be either delayed for the duration of the ISL project or be intermixed with ISL operations (e.g., oil and gas exploration). Changes to land use access including grazing restrictions and impacts on recreational activities would be limited due to the small size of restricted areas, temporary nature of restrictions, and availability of other land for these activities. Ecological, historical, and cultural resources could be affected but would be protected by careful planning and surveying to help identify resources and avoid or mitigate impacts. For all land use aspects except ecological, historical and cultural resources, the potential impacts would be SMALL. Due to the potential for unidentified resources to be altered or destroyed during excavation, drilling, and grading, the potential impacts to ecological, historical or cultural resources would be SMALL to LARGE, depending on local conditions. OPERATION—The types of land use impacts for operational activities would be similar to construction impacts regarding access restrictions because the infrastructure would be in place. Additional land disturbances would not occur from conducting operational activities. Because access restriction and land disturbance related impacts would be SMALL. AQUIFER RESTORATIO

ഗ	
\subset	
=	
⊐	
mmu	
$\stackrel{\sim}{\sim}$	
<u>u</u>	
J	
_	
으	
_	
E.	
ш.	
ږ	
≤.	
=:	
0	
\supset	
⋾	
ب	
ัด	
en	
nent	
nental	
vironmental o	
\sim	
nental Consequences	

		e 10-2. Summary of Impacts for the Wyoming East Uranium Milling Region (continued)
Topic/ Resource	GEIS Section	Impact Findings
Transportation	4.3.2	CONSTRUCTION—Low magnitude traffic generated by ISL construction relative to local traffic counts would not significantly increase traffic or accidents on many of the roads in the region. Existing low traffic roads could be moderately impacted by the additional worker commuting traffic during periods of peak employment. The impact would be more pronounced in areas with lower traffic counts. MODERATE dust, noise, and incidental wildlife or livestock kill impacts would be possible on, or near, site access roads (dust in particular for unpaved access roads)—SMALL to MODERATE. OPERATION—Low magnitude traffic relative to local traffic counts on most roads would not significantly increase traffic or accidents. Existing low traffic roads could be moderately impacted by commuting traffic during periods of peak employment including dust, noise, and possible incidental wildlife or livestock kill impacts on, or near, site access roads High consequences are possible for a severe accident involving transportation of hazardous chemicals in a populated area. However, the probability of such accidents occurring would be low, owing to the limited number of shipments, comprehensive regulatory controls, and use of best management practices. For radioactive material shipments (yellowcake product, ion exchange resins, waste materials) compliance with transportation regulations would limit radiological risk for normal operations. Low radiological risk is estimated for accident conditions. Emergency response protocols would help mitigate long-term consequences of severe accidents involving release of uranium—SMALL to MODERATE.
		AQUIFER RESTORATION—The magnitude of transportation activities would be lower than for construction and operations, with the exception of workforce commuting which could have moderate impacts on, or near, existing low traffic roads—SMALL to MODERATE.
		DECOMMISSIONING—The types of transportation activities, and therefore, types of impacts would be similar to those discussed for construction and operations except the magnitude of transportation activities (e.g., number and types of waste and supply shipments, no yellowcake shipments) from decommissioning could be lower than for operations. Accident risks would be bounded by operations yellowcake transportation risk estimates—SMALL.

Table 10-2. Summary of Impacts for the Wyoming East Uranium Milling Region (continued)		
Topic/	GEIS	
Resource	Section	Impact Findings
Geology and Soils	4.3.3	CONSTRUCTION—Disturbance to soil would occur from construction (clearing, excavation, drilling, trenching, road construction). However, such disturbances would be temporary and SMALL (approx. 15 percent of the total site area), and potential impacts would be mitigated by using best management practices. A large portion of the well fields, trenches, and access roads would be expected to be restored and reseeded after construction. Excavated soils would be stockpiled, seeded, and stored onsite until needed for reclamation fill. No impacts to subsurface geological strata are likely—SMALL. OPERATION—Temporary contamination or alteration of soils would be likely from operational leaks and spills and possible from transportation, use of evaporation ponds, or land application of treated waste water. However, detection and response techniques, monitoring of treated waste water, and eventual survey and decommissioning of all potentially impacted soils would limit the magnitude of overall impacts to soils—SMALL. AQUIFER RESTORATION—Impacts to geology and soils from aquifer restoration activities would be similar to impacts from operations due to use of the same infrastructure and similar activities conducted (e.g., well field operation, transfer lines, waste water treatment and disposal)—SMALL. DECOMMISSIONING—Impacts to geology and soils from decommissioning would be similar to impacts from construction. Activities to clean up, re-contour, and reclaim disturbed lands during decommissioning would mitigate long-term impacts to soils—SMALL.

Summary
으
of Environmental
Consequences

		ble 10-2. Summary of Impacts for the Wyoming East Uranium Milling Region (continued)
Topic/	GEIS	love and File discuss
Resource	Section	Impact Findings
Surface Waters	4.3.4.1	CONSTRUCTION—Impacts to surface waters and related habitats from construction (road crossings, filling, erosion, runoff, spills or leaks of fuels and lubricants for construction equipment) would be expected to be mitigated through prope planning, design, construction methods, and best management practices. The average annual surface runoff is similar to or slightly less than that in the Wyoming West Uranium Milling Region. As a result, runoff-related impacts will be similar. Some impacts directly related to the construction activities would be expected to be temporary and limited to the duration of the construction period. U.S. Army Corps of Engineers permits may be required when filling and crossing wetlands. Temporary changes to spring and stream flows from grading, changes in topography, and natural drainage patterns woul be mitigated through best management practices, and restored after the construction phase. Incidental spills of drilling fluids into local streams would be small and temporary due to implementation of mitigation measures. Impacts from construction of roads, parking areas, and buildings on recharge to shallow aquifers would be small, owing to the limited area of impervious surfaces proposed. Infiltration of drilling fluids into the local aquifer would be SMALL, temporary, and localized to a few feet around boreholes—SMALL, depending on site-specific characteristics OPERATION—Impacts from storm water runoff or direct discharge of process waters (brine reject from reverse osmosis, or spent eluants from an ion exchange system) to surface waters would be regulated by the Wyoming Department of Environmental Quality through the Wyoming Pollutant Discharge Elimination System permit. The increased areal runoff projections for this region would result in a potential increase of runoff-related impacts. Expansion of facilities or pipelined during operations would generate impacts similar to construction. Because the aquifers containing uranium ore-bodies would have a weak, if any, connection to local

	Table 10-2. Summary of Impacts for the Wyoming East Uranium Milling Region (continued)		
	EIS	lunnast Findings	
Resource Sec	ction 3.4.2	CONSTRUCTION—Water use impacts would be limited by the small volumes of groundwater used for routine activities such as dust suppression, mixing cements, and drilling support over short and intermittent periods. Contamination of groundwater from construction activities would be mitigated by best management practices—SMALL. OPERATION—Potential impacts to shallow aquifers can occur from leaks or spills from surface facilities and equipment. Shallow aquifers are important sources of drinking water in some areas of the Wyoming East Uranium Milling Region. Potential impacts to the ore-bearing and surrounding aquifers include consumptive water use and degradation of water quality (from normal production activities, off-normal excursion events, and deep well injection disposal practices). Consumptive use impacts from withdrawal of groundwater would be SMALL because only 1 to 3 percent of pumped groundwater would not be returned to the aquifer (e.g., process bleed). The amount of water lost could be reduced substantially by currently available treatment methods (e.g., reverse osmosis, brine concentration). Effects of water withdrawal on surface water would be SMALL, as the ore zone normally occurs in a confined aquifer. Estimated drawdown effects vary depending on site conditions and water treatment technology applied. Excursions of lixiviant and mobilized chemical constituents could occur from a failure of twell seals or other operational conditions that result in incomplete recovery of lixiviant. Well-seal-related excursions would be detected by the groundwater monitoring system, and periodic well integrity testing, and impacts would be mitigated during operation or aquifer restoration. Other excursions could result in plumes of mobilized uranium and heavy metals extending beyond the mineralization zone. The magnitude of potential impacts from vertical excursions at ISL facilities, NRC requires licensees to take preventive measures prior to starting operations including well tests, monitoring, and development of pro	

	Table 10-2. Summary of Impacts for the Wyoming East Uranium Milling Region (continued)			
Topic/	GEIS			
Resource	Section	Impact Findings		
		CONSTRUCTION—Potential terrestrial ecology impacts would include the removal of vegetation from well fields and the milling site, the modification of existing vegetative communities, the loss of sensitive plants and habitats from clearing and grading, and the potential spread of invasive species and noxious weed populations. These impacts would be temporary because restoration and reseeding occur rapidly after the end of construction. Introduction of invasive species and noxious weeds would be possible but could be mitigated by restoration and reseeding after construction. Shrub and tree removal would have a longer restoration period. Construction noise could affect reproductive success of sage-grouse leks by interfering with mating calls. Temporary displacement of animal species would also be possible. Crucial wintering and year-long ranges are important to survival of big game and sage grouse. Wildlife habitat fragmentation, temporary displacement of animal species, and direct or indirect mortalities is also possible. Implementation of wildlife surveys and mitigation measures following established guidelines would limit these impacts. The magnitude of impacts depends on whether a new facility is being licensed or an existing facility is being extended—SMALL to MODERATE, depending on site-specific habitat.		
Ecology— Terrestrial	4.3.5.1	OPERATION—Habitat could be altered by operations (fencing, traffic, noise), and individual takes could occur due to conflicts between species habitat and operations. Access to crucial wintering habitat and water could be limited by fencing. However, the Wyoming Game and Fish Department specifies fencing construction techniques to minimize impediments to big game movement. Migratory birds could be affected by exposure to constituents in evaporation ponds, but perimeter fencing, and netting would limit impacts. Temporary contamination or alteration of soils would be from operational leaks and spills and possible from transportation or land application of treated waste water. However, detection and response techniques and eventual survey and decommissioning of all potentially impacted soil, would limit the magnitude of overall impacts to terrestrial ecology. Mitigation measures such as perimeter fencing, netting, alternative sites, and timing stipulations would reduce overall impacts—SMALL.		
		AQUIFER RESTORATION—Impacts include habitat disruption, but existing (in-place) infrastructure would be used during aquifer restoration, with little additional ground disturbance. Migratory birds could be affected by exposure to constituents in evaporation ponds, but perimeter fencing, and netting would limit impacts. Contamination of soils could result from leaks and spills or land application of treated waste water. However, detection and response techniques, and eventual survey and decommissioning of all potentially impacted soils, would limit the magnitude of overall impacts to terrestrial ecology. Mitigation measures such as perimeter fencing, netting, alternative sites, and timing stipulations would reduce overall impacts—SMALL.		
		DECOMMISSIONING—During decommissioning and reclamation, there would be a temporary disturbance to land (e.g., excavating soils, buried piping, removal of structures). However, revegetation and re-contouring would restore habitat altered during construction and operations. Wildlife would be temporarily displaced, but are expected to return after decommissioning and reclamation are completed and vegetation and habitat are reestablished—SMALL.		

	Table 10-2. Summary of Impacts for the Wyoming East Uranium Milling Region (continued)		
Topic/ Resource	GEIS Section	Impact Findings	
Ecology—Aquatic	4.3.5.2	CONSTRUCTION—Clearing and grading activities associated with construction could result in a temporary increase in sediment load in local streams, but aquatic species would recover quickly as sediment load decreases. Clearing of riparian vegetation could affect light and temperature of water. Construction impacts to wetlands would be identified and managed through U.S. Army Corps of Engineers permits, as appropriate. Construction impacts to surface waters and aquatic species would be temporary and mitigated by best management practices—SMALL. OPERATION—Impacts could result from spills or releases into surface water. Impacts would be minimized by spill prevention, identification and response programs, and National Pollutant Discharge Elimination System (NPDES) permit requirements—SMALL.	
		AQUIFER RESTORATION—Activities would use existing (in-place) infrastructure, and impacts could result from spills o releases of untreated groundwater. Impacts would be minimized by spill prevention, identification, and response programs, and NPDES permit requirements—SMALL.	
		DECOMMISSIONING—Decommissioning and reclamation activities could result in temporary increases in sediment load in local streams, but aquatic species would recover quickly as sediment load decreases. With completion of decommissioning, revegetation, and re-contouring, habitat would be reestablished and impacts would, therefore, be limited—SMALL.	

	Table 10-2. Summary of Impacts for the Wyoming East Uranium Milling Region (continued)		
Topic/ Resource	GEIS Section	Impact Findings	
Ecology— Threatened or Endangered Species		CONSTRUCTION—Numerous threatened and endangered species and State Species of Concern are located in the region. Small fragmentation of habitats could occur, in addition to potential habitat loss. The magnitude of impacts depends on the size of a new facility or extension to an existing facility and the amount of land disturbance. Inventory of threatened or endangered species would be developed during site-specific reviews to identify unique or special habitats, and Endangered Species Act consultations conducted with the U.S. Fish and Wildlife Service would assist in identifying potential impacts—SMALL to LARGE—depending on site-specific habitat and presence of threatened or endangered species.	
	4.3.5.3	OPERATION—Impacts could result from individual takes due to conflicts with operations. Small fragmentation of habitats would occur, in addition to potential habitat loss. The magnitude of impacts would depend on the size of a new facility or extension to an existing facility and the amount of land disturbance. Impacts could potentially result from spills or permitted effluents, but would be minimized by spill prevention measures, identification and response programs, and NPDES permit requirements. Inventory of threatened or endangered species developed during site-specific reviews would identify unique or special habitats, and Endangered Species Act consultations conducted with the U.S. Fish and Wildlife Service would assist in identifying potential impacts—SMALL.	
		AQUIFER RESTORATION—Impacts could result from individual takes due to conflicts with aquifer restoration activities (equipment, traffic). Existing (in-place) infrastructure would be used during aquifer restoration, so additional land-disturbing activities and habitat fragmentation would not be anticipated. Impacts may result from spills or releases of treated or untreated groundwater, but impacts would be minimized by spill prevention measures, identification, and response programs, and NPDES permit requirements. Inventory of threatened or endangered species would be developed during site-specific reviews to identify unique or special habitats, and Endangered Species Act consultations with the U.S. Fish and Wildlife Service would assist in identifying potential impacts—SMALL.	
		DECOMMISSIONING—Impacts resulting from individual takes could occur due to conflicts with decommissioning activities (equipment, traffic). Temporary land disturbance would occur as structures are demolished and removed and the ground surface is re-contoured. Inventory of threatened or endangered species developed during site-specific environmental review of the decommissioning plan would identify unique or special habitats, and Endangered Species Act consultations with the U.S. Fish and Wildlife Service would assist in identifying potential impacts. With completion of decommissioning, revegetation, and re-contouring, habitat would be reestablished and impacts would, therefore, be limited—SMALL.	

	Т	able 10-2. Summary of Impacts for the Wyoming East Uranium Milling Region (continued)
Topic/	GEIS	Impact Findings
Topic/ Resource	_	Impact Findings CONSTRUCTION—Fugitive dust and combustion (vehicle and diesel) emissions during land-disturbing activities associated with construction would be small, short-term, and reduced through best management practices (e.g., dust suppression). For example, estimated fugitive dust emissions during ISL construction are less than 2 percent of the National Ambient Air Quality Standards (NAAQS) for PM2,5 and less than 1 percent for PM10. For NAAQS attainment areas such as the Wyoming East Uranium Milling Region, nonradiological air quality impacts would be SMALL, and there are no Prevention of Significant Deterioration (PSD) Class I areas in the Wyoming East Uranium Milling Region. Furthermore, if impacts were initially assessed at a higher significance level, permit requirements would impose conditions or mitigation to reduce impacts—SMALL. OPERATION—Radiological impacts can result from dust releases from drying of lixiviant pipeline spills, radon releases from well system relief valves, resin transfer, or elution, and gaseous/particulate emissions from yellowcake dryers. Only small amounts of low dose materials would be expected to be released based on operational controls and rapid response to spills. Required spill prevention, control, and response procedures would be used to minimize impacts from spills. High Efficiency Particulate Air (HEPA) filters and vacuum dryer designs reduce particulate emissions from operations and ventilation reduces radon buildup during operations. Compliance with the NRC-required radiation monitoring program ensures releases would be within regulatory limits. Other potential nonradiological emissions during operations include fugitive dust and fuel from equipment, maintenance, transport trucks, and other vehicles. For NAAQS attainment areas such as the Wyoming East Uranium Milling Region, nonradiological air quality impacts were initially assessed at a higher significance level, permit requirements would impose condition, or mitigation measures to reduce impacts—SMALL. AQU
		DECOMMISSIONING—Fugitive dust and combustion (vehicle and diesel) emissions during land-disturbing activities associated with decommissioning would be similar to, or less than that associated with construction, short-term, and reduced through best management practices (e.g., dust suppression). These impacts would decrease as decommissioning and reclamation of disturbed areas are completed. For NAAQS attainment areas such as the Wyoming East Uranium Milling Region, nonradiological air quality impacts would be SMALL, and there are no PSD Class I areas in the Wyoming East Uranium Milling Region. Furthermore, if impacts were initially assessed at a higher significance level, permit requirements would impose conditions or mitigation measures to reduce impacts—SMALL.

Summ
ary o
of Envir
ronmen
าtal (
Environmental Consequences
lences

	T	able 10-2. Summary of Impacts for the Wyoming East Uranium Milling Region (continued)
Topic/ Resource	GEIS Section	Impact Findings
Resource	Section	CONSTRUCTION—Noise generated during construction would be noticeable in proximity to operating equipment, but would be temporary (typically daytime only). Administrative and engineering controls would be used to maintain noise levels in work areas below Occupational Health and Safety Administration (OSHA) regulatory limits and be mitigated by use of personal hearing protection. Traffic noise during construction (commuting workers, truck shipments to and from the facility, and construction equipment such as trucks, bulldozers, compressors) would be localized, limited to highways in the vicinity of the site, access roads within the site, and roads in well fields. Relative increases in traffic levels would be small for larger roads, but may be moderate for lightly traveled rural roads through less populated communities. Noise may adversely affect wildlife habitat and their reproductive success in the immediate vicinity of construction activities. Noise levels decrease geometrically with distance, and at distances more than 300 m [1,000 ft], ambient noise levels would return to background levels. Wildlife generally avoid construction noise areas. The three uranium districts in the Wyoming East Uranium Milling Region are located in undeveloped rural areas, generally 16 km [10 mi] from the closest communities—SMALL to MODERATE.
Noise	4.3.7	OPERATION—Noise-generating activities in the central uranium processing facility would be indoors, minimizing offsite sound levels. Well field equipment (e.g., pumps, compressors) would also be expected to be contained within structures (e.g., header houses, satellite facilities) minimizing sound levels to offsite receptors. Administrative and engineering controls would be used to maintain noise levels in work areas below OSHA regulatory limits and be mitigated by use of personal hearing protection. Traffic noise from commuting workers, truck shipments to and from the facility, and facility equipment would be expected to be localized, limited to highways in the vicinity of the site, access roads within the site, and roads in well fields. Relative increases in traffic levels would be SMALL for larger roads, but may be MODERATE for lightly traveled rural roads through less populated communities. Most noise would be generated indoors and mitigated by regulatory compliance and best management practices. Noise from trucks and other vehicles is typically of short duration. Noise usually is not discernable to offsite receptors at distances of more than 300 m [1,000 ft]. The three uranium districts in the Wyoming East Uranium Milling Region are located in undeveloped rural areas, generally 16 km [10 mi] from the closest communities—SMALL to MODERATE.
		AQUIFER RESTORATION—Noise generation is expected to be less than during construction and operations. Pumps and other well field equipment contained in buildings, minimize sound levels to offsite receptors. Existing operational infrastructure would be used, and traffic levels would be less than during construction and operations; however, relative increases to existing traffic levels from commuting may be more significant for lightly traveled rural roads through smaller communities. Noise usually is not discernable to offsite receptors at distances of more than 300 m [1,000 ft]. The three uranium districts in the Wyoming East Uranium Milling Region are located in undeveloped rural areas, generally 16 km [10 mi] from the closest communities—SMALL to MODERATE.
		DECOMMISSIONING—Noise generated during decommissioning would be noticeable only in proximity to operating equipment and be temporary (typically daytime only). Administrative and engineering controls would be used to maintain noise levels in work areas below OSHA regulatory limits and be mitigated by use of personal hearing protection. Noise levels during decommissioning would be expected to be less than during construction and would diminish as less and less equipment is used and truck traffic is reduced. Noise usually is not discernable to offsite receptors at distances of more than 300 m [1,000 ft]. The three uranium districts in the Wyoming East Uranium Milling Region are located in undeveloped rural areas, generally 16 km [10 mi] from the closest communities—SMALL to MODERATE.

Table 10-2. Summary of Impacts for the Wyoming East Uranium Milling Region (continued)

Topic/

GEIS

Table 10-2. Summary of Impacts for the Wyoming East Uranium Milling Region (continued)			
Topic/	GEIS	loon and Pin dia on	
Resource	Section	Impact Findings	
Visual and Scenic	4.3.9	CONSTRUCTION—Visual impacts result from equipment (drill rig masts, cranes), dust/diesel emissions from construction equipment, and hillside and roadside cuts. Most of the Wyoming East Uranium Milling Region is classified as Visual Resource Management (VRM) Class II through IV, and no VRM Class I or PSD Class I areas are located in the region. Most potential visual impacts during construction would be temporary as equipment is moved, and would be mitigated by implementing best management practices (e.g., dust suppression). Because of the generally rolling topography of the region, most visual impacts during construction would not be visible from more than about 1 km [0.6 mi]. The uranium districts in the region are located more than 8 km [5 mi] from the closest VRM Class III area, and the visual impacts associated with ISL construction would be consistent with the predominant VRM Class III and IV—SMALL. OPERATION—Visual impacts during operations would be expected to be less than those associated with construction. Most of the well field surface infrastructure has a low profile, and most piping and cables would be buried. The tallest structures would include the central uranium processing facility {10 m [30 ft]} and power lines {6 m [20 ft]}. Because of the generally rolling topography of the region, most visual impacts during operations would not be expected to be visible from more than about 1 km [0.6 mi]. Irregular layout of well field surface structures such as wellhead protection and header houses would reduce visual contrast. Best management practices, design (e.g., painting buildings), and landscaping techniques would be used to mitigate potential visual impact. The three uranium districts in the region are located more than 8 km [5 mi] from the closest VRM Class II area, and the visual impacts associated with ISL construction would be expected to be consistent with the predominant VRM Class III and IV—SMALL. DECOMMISSIONING—Because similar equipment would be used and activities conducted, potential vis	

			depending on timing of construction with other stages of the ISL lifecycle. During construction of surface facilities and well fields, the general practice has been to use local contractors (drillers, construction) if available. A local multiplier of 0.7 would indicate a maximum of about 140 ancillary jobs could be created. For example, local building materials and building supplies would be used to the extent practical. Most employees would live in larger communities with access to more services. Some construction employees, however, would commute from outside the county to the ISL facility, and skilled employees (e.g., engineers, accountants, managers) would come from outside the local work force. Some of these employees would temporarily relocate to the project area and contribute to the local economy through purchasing goods and services and taxes. Because of the small relative size and temporary nature of the ISL construction workforce, net impacts would be SMALL to MODERATE, depending on proximity to less populated areas such as those in Niobrara or Albany Counties.
10-30	Socioeconomics	4.3.10	OPERATION—Employment levels for ISL facility operations would be similar to, or less than for construction, with total peak employment depending on timing and overlap with other stages of the ISL lifecycle. Use of local contract workers and local building materials would diminish after the construction stage. Additional revenues would be generated by federal, state, and local taxes on the facility and the uranium produced. Because of similar employment levels, other socioeconomic impacts would be expected to be similar to construction—SMALL to MODERATE, depending on proximity to less populated areas such as those in Niobrara or Albany Counties.
			AQUIFER RESTORATION—Because much of the same (in-place) infrastructure would be used, employment levels would be similar to, or less than, for operations, with total peak employment depending on timing and overlap with other stages of the ISL lifecycle. Use of local contract workers and local building materials would diminish after the construction stage. Because of similar employment levels, other socioeconomic impacts would be similar to construction—SMALL to MODERATE, depending on proximity to less populated areas such as those in Niobrara or Albany Counties.
			DECOMMISSIONING—A skill set similar to the construction workforce would be involved in dismantling surface structures, removing pumps, plugging and abandoning wells, and reclaiming/re-contouring the ground surface. Employment levels and

proximity to less populated areas such as those in Niobrara or Albany Counties.

Table 10-2. Summary of Impacts for the Wyoming East Uranium Milling Region (continued)

Impact Findings

CONSTRUCTION—Potential impacts to socioeconomics would result predominantly from employment at an ISL facility and demands on the existing public and social services, tourism/recreation, housing, infrastructure (schools, utilities), and the local work force. Total peak employment would be about 200 people, including company employees and local contractors,

use of local contractor support during decommissioning would be similar to or less than what would be required for construction. Employment would be temporary as decommissioning activities are limited in duration. Because of similar employment levels, other socioeconomic impacts would be similar to construction—SMALL to MODERATE, depending on

GEIS

Section

Topic/ Resource

Topic/ Resource	GEIS Section	Impact Findings
		Impact Findings CONSTRUCTION—Worker safety would be addressed by standard construction safety practices. Fugitive dust would result from construction activities and vehicle traffic but would likely be of short duration, and would not result in a radiological dose. Diesel emissions would not be a concern for worker or public health, because the releases would be of short duration and readily dispersed into the atmosphere—SMALL. OPERATION—Potential occupational radiological impacts from normal operations would be caused primarily by exposure to radon gas from the well field, ion exchange resin transfer operations, and venting during processing activities. Workers would also be exposed to airborne uranium particulates from dryer operations and maintenance activities. Potential public exposures to radiation would occur from the same radon releases and uranium particulate releases (i.e., from facilities without vacuum dryer technology). Both worker and public radiological exposures would be limited by NRC regulations at 10 CFR Part 20 which require licensees to implement an NRC-approved monitoring and radiation protection program. (Measured and calculated doses for workers and the public are commonly a fraction of regulated limits.) Nonradiological worker safety matters would be addressed through commonly applied occupational health and safety regulations and practices. Radiological accident risks could involve processing equipment failures leading to yellowcake slurry spills, or radon gas or uranium particulate releases. Consequences of accidents to workers and the public are generally low, with the exception of a dryer explosion, which could result in worker dose above NRC limits. The likelihood of such an accident
Safety		would be low, and therefore, the risk would also be low. Potential nonradiological accidents impacts include high-consequence chemical release events (e.g., ammonia) for both workers and nearby populations. The likelihood of such release events would be low, based on historical operating experience at NRC-licensed facilities, which is partly the result o operators following commonly applied chemical safety and handling protocols—SMALL to MODERATE. AQUIFER RESTORATION—Because the activities during aquifer restoration overlap with similar operational activities (e.g., operation of well fields, waste water treatment and disposal) the types of impacts on public and occupational health and safety would be similar to operational impacts. The reduction of some operational activities (e.g., yellowcake production and drying, remote ion exchange) further limits the relative magnitude of potential worker and public health and safety hazards—SMALL. DECOMMISSIONING—Worker and public health and safety would be addressed in a required decommissioning plan. This

			OPERATION—Operational wastes primarily result from liquid waste streams including process bleed, flushing of depleted eluant to limit impurities, resin transfer wash, filter washing, uranium precipitation process wastes (brine), and plant washdown water. State permitting actions, NRC license conditions, and NRC inspections ensure the proper practices would be used to comply with safety requirements to protect workers and the public. Waste treatment such as reverse osmosis and radon settling would help in segregating wastes and minimizing disposal volumes. Potential impacts from surface discharge and deep well injection would be limited by the applicable permitting processes. NRC regulations address constructing, operating, and monitoring for leakage from evaporation ponds used to store and reduce volumes of liquid wastes. Potential impacts from land application of treated wastewater would be addressed by NRC review of site-specific conditions prior to approval, routine monitoring, and inclusion of irrigated land areas in decommissioning surveys. Offsite waste disposal impacts would be SMALL for radioactive wastes as a result of required preoperational disposal agreements. Impacts for hazardous and municipal waste would be SMALL due to the volume of wastes generated. For remote areas with limited available disposal capacity, such wastes may need to be shipped greater distances to facilities that have capacity. However, the volume of wastes generated, and magnitude of the shipments, are estimated to be low—SMALL.
10-32	Waste Management	4.3.12	AQUIFER RESTORATION—Waste management activities during aquifer restoration would utilize the same treatment and disposal options implemented for operations. Therefore, impacts associated with aquifer restoration would be similar to operational impacts. While the amount of waste water generated during aquifer restoration is dependent on site-specific conditions, the potential exists for additional generation of wastewater volume and associated treatment wastes during the restoration period. However, this would be offset to some degree by the reduction in production capacity from the removal of a well field. NRC review of future ISL facility applications would verify that sufficient water treatment and disposal capacity (and the associated agreement for disposal of byproduct material) are addressed. As a result, waste management impacts from aquifer restoration would be low—SMALL.
			DECOMMISSIONING—Radioactive wastes from decommissioning ISL facilities (including contaminated excavated soil, evaporation pond bottoms, process equipment) would be disposed of as byproduct material at an NRC-licensed facility. A preoperational agreement with a licensed disposal facility to accept radioactive wastes ensures sufficient disposal capacity would be available for byproduct wastes generated by decommissioning activities. Safe handling, storage, and disposal of

decommissioning radioactive, chemical, and solid wastes would be small-SMALL.

Table 10-2. Summary of Impacts for the Wyoming East Uranium Milling Region (continued)

fields at ISL facilities would generate low volumes of construction waste—SMALL.

Impact Findings CONSTRUCTION—The relatively small scale of construction activities (Section 2.3) and incremental development of well

decommissioning wastes would be addressed in a required decommissioning plan, subject to NRC review. This plan details how a 10 CFR Part 20-compliant radiation safety program would be implemented during decommissioning, to ensure safety of workers and the public and to comply with applicable safety regulations would be complied with. Overall, volumes of

GEIS

Section

Topic/ Resource

Summary
ᅌ
of Environmental (
Consequences

	Table 1	0-3. Summary of Impacts for the Nebraska-South Dakota-Wyoming Uranium Milling Region
Topic/	GEIS	
Resource	Section	Impact Findings
Land Use	4.4.1	CONSTRUCTION—Land use impacts could occur from land disturbances (including alterations of ecological cultural or historic resources) and access restrictions (including limitations of other mineral extraction activities, grazing activities, or recreational activities). A higher percentage of private land ownership occurs in this region than in the Wyoming West Uranium Milling Region, and could increase the potential for land use conflicts with private land owners. Land disturbances during construction would be temporary and limited to specific areas within permitted area. Well sites, staging areas, and trenches would be reseeded and restored. Unpaved access roads would remain in use until decommissioning. Competing access to mineral rights could be either delayed for the duration of the ISL project or be intermixed with ISL operations (e.g., oil and gas exploration). Changes to land use access including grazing restrictions and impacts on recreational activities would be limited due to the small size of restricted areas, temporary nature of restrictions, and availability of other land for these activities. Ecological, historical, and cultural resources could be affected, but would be protected by careful planning and surveying to help identify resources and avoid or mitigate impacts. For all land use aspects except ecological, historical and cultural resources, the potential impacts would be SMALL. Due to the potential for unidentified resources to be altered or destroyed during excavation, drilling, and grading, the potential impacts to ecological, historical or cultural resources would be SMALL to LARGE, depending on local conditions. OPERATION—The types of land use impacts for operational activities would be similar to construction impacts regarding access restrictions because the infrastructure would be in place. Additional land disturbances would not occur from conducting operational activities. Because access restriction and land disturbance related impacts would be similar to, or less than, expected for

	Table 10-3. \$	Summary of Impacts for the Nebraska-South Dakota-Wyoming Uranium Milling Region (continued)
Topic/ Resource	GEIS Section	Impact Findings
Transportation	4.4.2	CONSTRUCTION—Low magnitude traffic generated by ISL construction relative to local traffic counts would not significantly increase traffic or accidents on many of the roads in the region. Existing low traffic roads could be moderately impacted by the additional worker commuting traffic during periods of peak employment. This impact would be more pronounced in the Nebraska-South Dakota-Wyoming Uranium Milling Region owing to the relatively lower traffic counts in this region, in comparison to the other milling regions. Moderate dust, noise, and incidental wildlife or livestock kill impacts would be possible on, or near, site access roads (dust in particular for unpaved access roads)—SMALL to MODERATE. OPERATION—Low magnitude traffic relative to local traffic counts on most roads would not significantly increase traffic or accidents. Existing low traffic roads could be moderately impacted by commuting traffic during periods of peak employment including dust, noise, and possible incidental wildlife or livestock kill impacts on, or near, site access roads. High consequences would be possible for a severe accident involving transportation of hazardous chemicals in a populated area. However, the probability of such accidents occurring would be low owing to the limited number of shipments, comprehensive regulatory controls, and use of best management practices. For radioactive material shipments (yellowcake product, ion exchange resins, waste materials), compliance with transportation regulations would limit radiological risk for normal operations. Low radiological risk is estimated for accident conditions. Emergency response protocols would help mitigate long-term consequences of severe accidents involving release of uranium—SMALL to MODERATE.
		AQUIFER RESTORATION—The magnitude of transportation activities would be lower than for construction and operations, with the exception of workforce commuting, which could have moderate impacts on, or near, existing low traffic roads—SMALL to MODERATE.
		DECOMMISSIONING—The types of transportation activities and, therefore, types of impacts would be similar to those discussed for construction and operations except the magnitude of transportation activities (e.g., number and types of waste and supply shipments, no yellowcake shipments) from decommissioning could be lower than for operations. Accident risks would be bounded by operations yellowcake transportation risk estimates—SMALL

Summary
으
of Environmental Consequence
Consequences

	able 10-3. 3	Summary of Impacts for the Nebraska-South Dakota-Wyoming Uranium Milling Region (continued)
Topic/ Resource	GEIS Section	Impact Findings
Geology and Soils	4.4.3	CONSTRUCTION—Disturbance to soil would occur from construction (clearing, excavation, drilling, trenching, road construction). However, such disturbances would be temporary and SMALL (approx. 15 percent of the total site area), and potential impacts would be mitigated by using best management practices. A large portion of the well fields, trenches, and access roads would be restored and reseeded after construction. Excavated soils would be stockpiled, seeded, and stored onsite until needed for reclamation fill. No impacts are expected to subsurface geological strata—SMALL. OPERATION—Temporary contamination or alteration of soils would be likely from operational leaks and spills and possible from transportation, use of evaporation ponds, or land application of treated wastewater. However, detection and response techniques, monitoring of treated waste water, and eventual survey and decommissioning of all potentially impacted soils, would limit the magnitude of overall impacts to soils—SMALL. AQUIFER RESTORATION—Impacts to geology and soils from aquifer restoration activities would be similar to impacts from operations due to use of the same infrastructure and similar activities conducted (e.g., well field operation, transfer lines, waste water treatment and disposal)—SMALL. DECOMMISSIONING—Impacts to geology and soils from decommissioning would be similar to impacts from construction. Activities to clean up, recontour, and reclaim disturbed lands during decommissioning would mitigate long-term impacts to soils—SMALL.

			design, construction methods, and best management practices. This region has the same or lower surface runoff (areal flow) than the Wyoming West Uranium Milling Region, and for that reason, runoff-related impacts will be similar or lower. Some impacts directly related to the construction activities would be temporary and limited to the duration of the construction period. U.S. Army Corps of Engineers permits may be required when filling and crossing wetlands. Temporary changes to spring and stream flows from grading and changes in topography and natural drainage patterns could be mitigated through best management practices, or restored after the construction phase. Incidental spills of drilling fluids into local streams would be SMALL and temporary, due to the implementation of mitigation measures. Impacts from construction of roads, parking areas, and buildings on recharge to shallow aquifers would be SMALL, temporary, and localized to a few feet around boreholes—SMALL to MODERATE depending on site-specific characteristics.	
10-36	Surface Waters	4.4.4.1	OPERATION—Impacts from storm water runoff or direct discharge of produced waters (brine reject from reverse osmosis, or spent eluants from an ion exchange system) to surface waters would be regulated by individual states through the National Pollutant Discharge Elimination System permits. Increased runoff compared to the Wyoming West Uranium Milling Region could potentially contribute to a slight increase in runoff-related impacts. Expansion of facilities or pipelines during operations would generate impacts similar to construction. Because the aquifers containing uranium orebodies would have a weak, if any, connection to local surface water features, such as streams and springs, the impacts of excess net groundwater extraction from local surface water bodies would be SMALL —SMALL to MODERATE depending on site-specific characteristics.	
			AQUIFER RESTORATION—Impacts from aquifer restoration would be similar to impacts from operations due to use of in-place infrastructure and similar activities conducted (e.g., well field operation, transfer lines, water treatment, stormwater runoff)—SMALL to MODERATE depending on site-specific characteristics.	

waters—SMALL to MODERATE depending on site-specific characteristics.

Table 10-3. Summary of Impacts for the Nebraska-South Dakota-Wyoming Uranium Milling Region (continued)

Impact Findings CONSTRUCTION—Impacts to surface waters and related habitats from construction (road crossings, filling, erosion, runoff, spills or leaks of fuels and lubricants for construction equipment) would be mitigated through proper planning,

DECOMMISSIONING—Impacts from decommissioning would be similar to impacts from construction. Activities to clean up, re-contour, and reclaim disturbed lands during decommissioning would mitigate long-term impacts to surface

Topic/

Resource

GEIS Section

Topic/ Resource CONSTRUCTION—Water use impacts would be limited by the small volumes of groundwater used for routine activities such as dust suppression, mixing cerents, and drilling support over short and intermittent periods. Contamination of groundwater from construction activities would be mitigated by use of best management practices—SMALL. OPERATION—Potential impacts to shallow aquifers can occur from leaks or spills from surface facilities and equipment. Shallow aquifers are important sources of drinking water in some areas of the Nebraska-South Dakota-Wyoming Uranium Milling Region. Potential impacts to the ore-bearing and surrounding aquifers include consumptive water use and degradation of water quality (from normal production activities, off-normal excursion events, and deep well injection disposal practices). Consumptive use impacts from withdrawal of groundwater would be SMALL as advantaged on surface substantially by currrently available treatment methods (e.g., reverse osmosis, brine concentration). Effects of water withdrawal on surface water would be SMALL, as for exportant production activities, off-normally occurs in a confined aquifer. Estimated drawdown effects vary depending on site conditions and water treatment technology applied. Excursions that cause incomplete recovery of lixiviant. Well-seal-related excursions would be detected by the groundwater monitoring system, and periodic well integrity testing, and impacts would be mitigated during operation or aquifer restoration. Other excursions could result in plumes of mobilized uranium and heavy metals extending beyond the mineralization zone. The magnitude of potential impacts from vertical excursions at SL facilities, NRC requires licensees to take preventative measures prior to starting operations including well tests, monitoring, and development of procedures that include excursion response measures and reporting requirements. Impacts associated with alterations of ore body aquifer chemistry would be SMALL because the aquifer would
CONSTRUCTION—Water use impacts would be limited by the small volumes of groundwater used for routine activities such as dust suppression, mixing cements, and drilling support over short and intermittent periods. Contamination of groundwater from construction activities would be mitigated by use of best management practices—SMALL. OPERATION—Potential impacts to shallow aquifers can occur from leaks or spills from surface facilities and equipment. Shallow aquifers are important sources of drinking water in some areas of the Nebraska-South Dakota-Wyoming Uranium Milling Region. Potential impacts to the ore-bearing and surrounding aquifers include consumptive water use and degradation of water quality (from normal production activities, off-normal excursion events, and deep well injection disposal practices). Consumptive use impacts from withdrawal of groundwater would be SMALL because only 1 to 3 percent of pumped groundwater is not returned to the aquifer (e.g., process bleed). The amount of water lost could be reduced substantially by currently available treatment methods (e.g., reverse osmosis, brine concentration). Effects of water withdrawal on surface water would be SMALL, as the ore zone normally occurs in a confined aquifer. Estimated drawdown effects vary depending on site conditions and water treatment technology applied. Excursions of lixiviant and mobilized chemical constituents could occur from a failure of well seals or other operational conditions that cause incomplete recovery of lixiviant. Well-seal-related excursions would be detected by the groundwater monitoring system, and periodic well integrity testing, and impacts would be mitigated during operation or aquifer restoration. Other excursions could result in plumes of mobilized uranium and heavy metals extending beyond the mineralization zone. The magnitude of potential impacts from vertical excursions would vary depending on site-specific conditions. To reduce the likelihood and consequences of potential excursions as 1SL facilities, NRC req
restoration in which pumped water is not recirculated. Potential environmental impacts associated with water consumption during aquifer restorations would be determined by: (1) the restoration techniques chosen, (2) the volume of water to be used, (3) the severity and extent of the contamination, and (4) the current and future use of the production and surrounding aquifers in the vicinity of the ISL facility or at the regional scale—SMALL to MODERATE depending on site-specific conditions. DECOMMISSIONING—Potential impacts from decommissioning would be similar to construction (water use, spills) with

	Table 10-3.	Summary of Impacts for the Nebraska-South Dakota-Wyoming Uranium Milling Region (continued)
_Topic/	GEIS	
Resource	Section	Impact Findings
	4.4.5.1	CONSTRUCTION—Potential terrestrial ecology impacts would include the removal of vegetation from well fields and the milling site, the modification of existing vegetative communities, the loss of sensitive plants and habitats from clearing and grading; and the potential spread of invasive species and noxious weed populations. These impacts would be temporary because restoration and reseeding occur rapidly after the end of construction. Introduction of invasive species and noxious weeds would be possible but could be mitigated by restoration and reseeding after construction. Shrub and tree removal would have a longer restoration period. Construction noise could affect reproductive success of sage-grouse leks (in the Wyoming part of the region) by interfering with mating calls. Temporary displacement of animal species would also be possible. Crucial wintering and year-long ranges are important to survival of big game and sage grouse. Wildlife habitat fragmentation, temporary displacement of animal species, and direct or indirect mortalities would be possible. Implementation of wildlife surveys and mitigation measures following established guidelines would limit impacts. The magnitude of impacts depends on whether a new facility is being licensed or an existing facility is being extended—SMALL to MODERATE, depending on site-specific habitat.
Ecology— Terrestrial		OPERATION—Habitat could be altered by operations (fencing, traffic, noise), and individual takes could occur due to conflicts between species habitat and operations. Access to crucial wintering habitat and water could be limited by fencing. However, the Wyoming Game and Fish Department specifies fencing construction techniques to minimize impediments to big game movement. Migratory birds could be affected by exposure to constituents in evaporation ponds, but perimeter fencing and netting would limit impacts. Temporary contamination or alteration of soils would be likely from operational leaks and spills or from land application of treated wastewater. However, detection and response techniques, and eventual survey and decommissioning of all potentially impacted soils, would limit the magnitude of overall impacts to terrestrial ecology. Mitigation measures such as perimeter fencing, netting, alternative sites, and timing stipulations would reduce overall impacts—SMALL.
		AQUIFER RESTORATION—Impacts include habitat disruption, but existing (in-place) infrastructure would be used during aquifer restoration, with little additional ground disturbance. Migratory birds could be affected by exposure to constituents in evaporation ponds, but perimeter fencing, and netting would limit impacts. Contamination of soils could result from leaks and spills or land application of treated wastewater. However, detection and response techniques, and eventual survey and decommissioning of all potentially impacted soils, would limit the magnitude of overall impacts to terrestrial ecology. Mitigation measures such as perimeter fencing, netting, alternative sites, and timing stipulations would reduce overall impacts—SMALL.
		DECOMMISSIONING—During decommissioning and reclamation, there would be a temporary disturbance to land (e.g., excavating soils, buried piping, removal of structures). However, revegetation and re-contouring would restore habitat altered during construction and operations. Wildlife would be temporarily displaced, but are expected to return after decommissioning and reclamation are completed and vegetation and habitat are reestablished—SMALL.

Summary	
으	
П	1
≦c	
Ĭ	
of Environmental (
=	<u> </u>
Cons	
*	
equen	
nc	
ces	

Table 10-3.	Summary of Impacts for the Nebraska-South Dakota-Wyoming Uranium Milling Region (continued)
Topic/ GEIS Resource Section	Impact Findings
Ecology—Aquatic 4.4.5.2	CONSTRUCTION—Clearing and grading activities associated with construction could result in a temporary increase in sediment load in local streams, but aquatic species would recover quickly as sediment load decreases. Clearing of riparian vegetation could affect light and temperature of water. Construction impacts to wetlands would be identified and managed through U.S. Army Corps of Engineers permits, as appropriate. Construction impacts to surface waters and aquatic species would be temporary and mitigated by best management practices—SMALL. OPERATION—Impacts could result from spills or releases into surface water. Impacts would be minimized by spill prevention, identification and response programs, and National Pollutant Discharge Elimination System (NPDES) permit requirements—SMALL. AQUIFER RESTORATION—Activities would use existing (in-place) infrastructure, and impacts could result from spills or releases of untreated groundwater. Impacts would be minimized by spill prevention, identification, and response programs, and NPDES permit requirements—SMALL. DECOMMISSIONING—Decommissioning and reclamation activities could result in temporary increases in sediment load in local streams, but aquatic species would recover quickly as sediment load decreases. With completion of decommissioning, revegetation, and re-contouring, habitat would be reestablished and impacts would, therefore, be

	Table 10-3. S	Summary of Impacts for the Nebraska-South Dakota-Wyoming Uranium Milling Region (continued)
Topic/	GEIS	
Resource	Section	Impact Findings
Ecology— Threatened or Endangered Species	4.4.5.3	CONSTRUCTION—Numerous threatened and endangered species and State Species of Concern are located in the region. Small fragmentation of habitats could occur in addition to potential habitat loss. The magnitude of impacts depends on the size of a new facility or extension to an existing facility and the amount of land disturbance. Inventory of threatened or endangered species would be developed during site-specific reviews to identify unique or special habitats, and Endangered Species Act consultations conducted with the U.S. Fish and Wildlife Service would assist in identifying potential impacts—SMALL to LARGE—depending on site-specific habitat and presence of threatened or endangered species. OPERATION—Impacts could result from individual takes due to conflicts with operations. Small fragmentation of habitats would occur in addition to potential habitat loss. The magnitude of impacts would depend on the size of a new facility or extension to an existing facility and the amount of land disturbance. Impacts could potentially result from spills or permitted effluents, but would be minimized by spill prevention measures, identification and response programs, and NPDES permit requirements. Inventory of threatened or endangered species developed during site-specific reviews would identify unique or special habitats, and Endangered Species Act consultations conducted with the U.S. Fish and Wildlife Service would assist in identifying potential impacts—SMALL—depending on site-specific habitat and presence of threatened or endangered species. AQUIFER RESTORATION—Impacts could result from individual takes due to conflicts with aquifer restoration activities (equipment, traffic). Existing (in-place) infrastructure would be used during aquifer restoration, so additional land-disturbing activities and habitat fragmentation would not be anticipated. Impacts may result from spills or releases of treated or untreated groundwater, but impacts would be minimized by implementing spill prevention measures, identification and res

	Table 10-3.	Summary of Impacts for the Nebraska-South Dakota-Wyoming Uranium Milling Region (continued)
Topic/	GEIS	
Resource	Section	
Resource Air Quality	Section 4.4.6	Impact Findings CONSTRUCTION—Fugitive dust combustion (vehicle and diesel) emissions during land-disturbing activities associated with construction would be small, short-term, and reduced through best management practices (e.g., dust suppression). For example, estimated fugitive dust emissions during ISL construction are less than 2 percent of the National Ambient Air Quality Standards (NAAQS) for PM _{2.5} and less than 1 percent for PM ₁₀ . For NAAQS attainment areas such as the Nebraska-South Dakota-Wyoming Uranium Milling Region, nonradiological air quality impacts would be SMALL. A Prevention of Significant Deterioration (PSD) Class I area exists (Wind Cave National Park, Black Hills, South Dakota). More stringent air quality standards would apply to any facility that could potentially impact the air quality of that area. If impacts were initially assessed at a higher significance level, permit requirements would impose conditions or mitigation measures to reduce impacts—SMALL. OPERATION—Radiological impacts can result from dust releases from drying of lixiviant pipeline spills, radon releases from well system relief valves, resin transfer, or elution, and gaseous/particulate emissions from yellowcake dryers. Only small amounts of low dose materials would be released based on operational controls and rapid response to spills. Required spill prevention, control, and response procedures would be used to minimize impacts from spills. High Efficiency Particulate Air (HEPA) filters and vacuum dryer designs reduce particulate emissions from operations and ventilation reduces radon buildup during operations. Compliance with the NRC-required radiation monitoring program ensures releases are within regulatory limits. Other potential nonradiological emissions during operations include fugitive dust and fuel from equipment, maintenance, transport trucks, and other vehicles. For NAAQS attainment areas such as the Nebraska-South Dakota-Wyoming Uranium Milling Region, nonradiological ir quality impacts would be small. A
		level, permit requirements would impose conditions or mitigation measures to reduce impacts—SMALL. AQUIFER RESTORATION—Because the same infrastructure would be used, air quality impacts are expected to be similar to, or less than, operations. For NAAQS attainment areas such as the Nebraska-South Dakota-Wyoming Uranium Milling Region, nonradiological air quality impacts would be SMALL. A PSD Class I area exists at Wind Cave National Park, South Dakota. More stringent air quality standards would apply to any facility that could potentially impact the air quality of that area. If impacts were initially assessed at a higher significance level, permit requirements would impose conditions or mitigation measures to reduce impacts—SMALL. DECOMMISSIONING—Fugitive dust and combustion (vehicle and diesel) emissions during land-disturbing activities associated with decommissioning would be similar to, or less than, construction, short-term, and reduced through use of best management practices (e.g., dust suppression). These impacts would decrease as decommissioning and reclamation of disturbed areas are completed. For NAAQS attainment areas such as the Nebraska-South Dakota-Wyoming Uranium Milling Region, nonradiological air quality impacts would be SMALL. A PSD Class I area exists at Wind Cave National Park, South Dakota. More stringent air quality standards would apply to any facility that could potentially impact the air quality of that area. If impacts were initially assessed at a higher significance level, permit requirements would impose conditions or mitigation measures to reduce impacts—SMALL.

	Table 10-3.	Summary of Impacts for the Nebraska-South Dakota-Wyoming Uranium Milling Region (continued)
Topic/	GEIS	Local First
Resource	Section 4.4.7	Impact Findings CONSTRUCTION—Noise generated during construction would be noticeable in proximity to operating equipment, but would be temporary (typically daytime only). Administrative and engineering controls would be used to maintain noise levels in work areas below Occupational Health and Safety Administrativa (OSHA) regulatory limits and be mitigated by use of personal hearing protection. Traffic noise during construction (commuting workers, truck shipments to and from the facility, and construction equipment such as trucks, buildozers, compressors) would be localized, limited to highways in the vicinity of the site, access roads within the site, and roads in well fields. Relative increases in traffic levels would be small for larger roads, but may be moderate for lightly traveled rural roads through less populated communities. Noise may also adversely affect wildlife habitat and their reproductive success in the immediate vicinity of construction activities. Noise levels decrease geometrically with distance, and at distances more than 300 mt 1,000 ttp. Inoise levels return to background levels. Wildlife generally avoid construction osic areas. The three uranium districts within the Nebraska-South Dakota-Wyoming Uranium Milling Region are generally more than 300 mt 1,000 ttp. Inoise penerating activities in the central uranium processing facility would be indoors, minimizing offsite sound levels. Well field equipment (e.g., pumps, compressors) would also be expected to be contained within structures (e.g., header houses, satellite facilities) minimizing sound levels to offsite receptors. Administrative and engineering controls would be used to highways in the vicinity of the site, access roads within the site, and roads in well fields. Relative increases in traffic levels would be SMALL for larger roads, but may be MODERATE for lighty traveled rural roads through lesses populated communities. Most noise would be generated indopors and mitigated by regulatory compliance and use of best management pract

	+	Summary of Impacts for the Nebraska-South Dakota-Wyoming Uranium Milling Region (continued)
		Insurant Findings
Topic/ Resource Historical and Cultural	GEIS Section	Impact Findings CONSTRUCTION—Potential impacts during ISL facility construction could include loss of, or damage and temporary restrictions on access to, historical, cultural, and archaeological resources. The eligibility evaluation of cultural resources for listing in the National Register of Historic Places (NRHP) under criteria in 36 CFR 60.4(a)—(d) and/or as Traditional Cultural Properties TCPs) is conducted as part of the site-specific review and NRC licensing procedures undertaken during the National Environmental Policy Act (NEPA) review process. The evaluation of impacts to any historic properties designated as TCPs and tribal consultations regarding cultural resources and TCPs also occur during the site-specific licensing application and review process. Consultations to determine whether significant cultural resources would be avoided or mitigated occurs during consultations with State Historic Preservation Offices (SHPOs), other governmental agencies, and Native American Tribes, including Tribal Historic Preservation Offices (SHPOs), other governmental agencies, and Native American Tribes, including Tribal Historic Preservation Offices (SHPOs), other governmental agencies, and Native American Tribes, including Tribal Historic Preservation Offices (SHPOs), other governmental agencies, and Native American Tribes, including Tribal Historic Preservation Offices (SHPOs), other governmental agencies with recording the discovery of previously undocumented cultural resources during initial construction. These procedures regarding the discovery of previously undocumented cultural resources would be less than during construction. Conditions in the NRC license requiring adherence to procedures typically require the licensee to stop work and to notify the appropriate federal, tribal, and state agencies with regard to mitigation measures—SMALL or MODERATE to LARGE depending on site-specific conditions. AQUIFER RESTORATION—Because less land disturbance occurs during the aquifer restoration phase, potential impact

٦	Table 10-3.	Summary of Impacts for the Nebraska-South Dakota-Wyoming Uranium Milling Region (continued)
Topic/	GEIS	
Resource	Section	Impact Findings
Visual and Scenic	4.4.9	CONSTRUCTION—Visual impacts result from equipment (drill rig masts, cranes), dust/diesel emissions from construction equipment, and hillside and roadside cuts. Most of the Nebraska-South Dakota-Wyoming Uranium Milling Region is classified as Visual Resource Management (VRM) Class II through IV. Most potential visual impacts during construction would be temporary as equipment is moved, and would be mitigated by implementing best management practices (e.g., dust suppression). Because of the generally rolling topography of the region, most visual impacts during construction would not be visible from more than 1 km [0.6 mi]. The three uranium districts in the region are located more than 16 km [10 mi] from the closest VRM Class II region and 40 km [25 mi] from the PSD Class I area at Wind Cave National Park in South Dakota. The visual impacts associated with ISL construction would be consistent with the predominant VRM Class III and IV—SMALL. OPERATION—Visual impacts during operations would be expected to be less than those associated with construction. Most of the well field surface infrastructure has a low profile, and most piping and cables would be buried. The tallest structures would include the central uranium processing facility {10 m [30 ft]} and power lines {6 m [20 ft]}. Because of the generally rolling topography of the region, most visual impacts during operations would not be visible from more than about 1 km [0.6 mi]. Irregular layout of well field surface structures such as wellhead protection and header houses would reduce visual contrast. Best management practices, design (e.g., painting buildings), and landscaping techniques would be used to mitigate potential visual impact. The three uranium districts in the region are located more than 16 km [10 mi] from the closest VRM Class II region and 40 km [25 mi] from the PSD Class I area at Wind Cave National Park in South Dakota. The visual impacts associated with ISL construction would be consistent with the predominant VRM Class III and IV—SMALL. DE

	Table 10-3.	Summary of Impacts for the Nebraska-South Dakota-Wyoming Uranium Milling Region (continued)
Topic/ Resource	GEIS Section	Impact Findings
Socioeconomics	4.4.10	CONSTRUCTION—Potential impacts to socioeconomics would result predominantly from employment at an ISL facility and demands on the existing public and social services, tourism/recreation, housing, infrastructure (schools, utilities), and the local work force. Total peak employment would be about 200, people including company employees and local contractors, depending on timing of construction with other stages of the ISL lifecycle. During construction of surface facilities and well fields, the general practice has been to use local contractors (drillers, construction) if available. A local multiplier of 0.7 would indicate a maximum of about 140 ancillary jobs could be created. For example, local building materials and building supplies would be used to the extent practical. Most employees would live in larger communities with access to more services. Some construction employees, however, would commute from outside the county to the ISL facility, and skilled employees (e.g., engineers, accountants, managers) would come from outside the local work force. Some of these employees would temporarily relocate to the project area and contribute to the local economy through purchasing goods and services and taxes. Because of the small relative size and temporary nature of the ISL workforce, net impacts would be SMALL to MODERATE, depending on proximity to less populated communities such as Oglala, Pine Ridge, and Sioux City. OPERATION—Employment levels for ISL facility operations would be similar to, or less than, for construction, with total peak employment depending on timing and overlap with other stages of the ISL lifecycle. Use of local contract workers and local building materials would diminish after the construction stage. Additional revenues would be generated by federal, state, and local taxes on the facility and the uranium produced. Because of similar employment levels, other socioeconomic impacts would be expected to be similar to construction—SMALL to MODERATE, depending on proximity to smaller communities
		AQUIFER RESTORATION—Because much of the same (in-place) infrastructure would be used, employment levels would be similar to, or less than, for operations, with total peak employment depending on timing and overlap with other stages of the ISL lifecycle. Use of local contract workers and local building materials would diminish after the construction stage. Because of similar employment levels, other socioeconomic impacts would be similar to construction—SMALL to MODERATE, depending on proximity to less populated communities such as Oglala, Pine Ridge, and Sioux City.
		DECOMMISSIONING—A skill set similar to the construction workforce would be involved in dismantling surface structures, removing pumps, plugging and abandoning wells, and reclaiming/recontouring the ground surface. Employment levels and use of local contractor support during decommissioning would be similar to or less than what would be required for construction. Employment would be temporary as decommissioning activities are limited in duration. Because of similar employment levels, other socioeconomic impacts would be similar to construction—SMALL to MODERATE, depending on proximity to less populated communities such as Oglala, Pine Ridge, and Sioux City.

		Summary of Impacts for the Nebraska-South Dakota-Wyoming Uranium Milling Region (continued)
Topic/	GEIS	
Resource	Section	Impact Findings
		CONSTRUCTION—Worker safety would be addressed by standard construction safety practices. Fugitive dust would result from construction activities and vehicle traffic, but would likely be of short duration, and not result in a radiological dose. Diesel emissions would not be a concern for worker or public health, because the releases would be of short duration and readily dispersed into the atmosphere—SMALL. OPERATION—Potential occupational radiological impacts from normal operations would be caused primarily by exposure to radon gas from well field, ion-exchange resin transfer operations, and venting during processing activities. Workers would also be exposed to airborne uranium particulates from dryer operations and maintenance activities. Potential public exposures to radiation would occur from the same radon releases and uranium particulate releases (i.e., from facilities without vacuum dryer technology). Both worker and public radiological exposures would be addressed by NRC regulations at 10 CFR Part 20 which require licensees to implement an NRC-approved radiation protection program. (Measured and calculated doses for workers and the public are commonly a fraction of regulated limits.) Nonradiological worker safety matters would be addressed through commonly applied occupational health and safety regulations and practices. Radiological accident risks could involve processing equipment failures leading to yellowcake slurry spills, or radon gas or uranium particulate releases. Consequences of accidents to workers and the public are generally low, with the exception of a dryer explosion, which could result in worker dose above NRC limits. The likelihood of such an accident would be low, and therefore the risk would also be low. Potential nonradiological accidents impacts include, high-consequence chemical release events (e.g., ammonia) for both workers and nearby populations. The likelihood of such result of operators following commonly applied chemical safety and handling protocols—SMALL to MODERATE. AQUIF
		safety hazards—SMALL. DECOMMISSIONING—Worker and public health and safety would be addressed in a required decommissioning plan. This plan details how a 10 CFR Part 20 compliant radiation safety program would be implemented during decommissioning, to ensure safety of workers and the public and would comply with applicable safety regulations—SMALL.

Table 10-3. Summary of Impacts for the Nebraska-South Dakota-Wyoming Uranium Milling Region (continued)		
Topic/	GEIS	
Resource	Section	Impact Findings
Waste Management	4.4.12	CONSTRUCTION—The relatively small scale of construction activities (Section 2.3) and incremental development of well fields at ISL facilities would generate low volumes of construction waste—SMALL. OPERATION—Operational wastes primarily result from liquid waste streams including process bleed, flushing of depleted eluant to limit impurities, resin transfer wash, filter washing, uranium precipitation process wastes (brine), and plant washdown water. State permitting actions, NRC license conditions, and NRC inspections ensure the proper practices would be used to comply with safety requirements to protect workers and the public. Waste treatment such as reverse osmosis and radon settling would help in segregating wastes and minimizing disposal volumes. Potential impacts from surface discharge and deep well injection would be limited by the applicable permitting processes. NRC regulations address constructing, operating, and monitoring for leakage from evaporation ponds used to store and reduce volumes of liquid wastes. Potential impacts from land application of treated wastewater would be addressed by NRC review of site-specific conditions prior to approval, routine monitoring, and inclusion of irrigated land areas in decommissioning surveys. Offsite waste disposal impacts would be SMALL for radioactive wastes as a result of required preoperational disposal agreements. Impacts for hazardous and municipal waste would be SMALL due to the volume of wastes generated. For remote areas with limited available disposal capacity, such wastes may need to be shipped greater distances to facilities that have capacity. However, the volume of wastes generated and magnitude of the shipments are estimated to be low—SMALL. AQUIFER RESTORATION—Waste management activities during aquifer restoration would utilize the same treatment and disposal options implemented for operations. Therefore, impacts associated with aquifer restoration would be similar to operational exists for additional generation of wastewater volume and associate

Table 10-4. Summary of Impacts for the Northwestern New Mexico Uranium Milling Region			
Topic/ Resource	GEIS Section	Impact Findings	
Land Use	4.5.1	CONSTRUCTION—Land use impacts could occur from land disturbances (including alterations of ecological cultural or historic resources) and access restrictions (including limitations of other mineral extraction activities, grazing activities, or recreational activities). A higher percentage of private land and Native American land ownership occurs in this region than in the Wyoming West Uranium Milling Region, and a more complex patchwork of land ownership could increase the potential for land use conflicts with private and other land owners. Land disturbances during construction would be temporary, but limited to specific locations within the permitted site. Well sites, staging areas, and trenches would be reseeded and restored after construction. Unpaved access roads would remain in use until decommissioning is completed. Competing access to mineral rights could be either delayed for the duration of the ISL project or be intermixed with ISL operations (e.g., oil and gas exploration). Changes to land use access including grazing restrictions and impacts on recreational activities would be limited due to the small size of restricted areas, temporary nature of restrictions, and availability of other land for these activities. Ecological, historical, and cultural resources could be affected but would be minimized due to careful planning and surveying to help identify resources and avoid or mitigate impacts. For all land use aspects except ecological, historical and cultural resources, the potential impacts would be SMALL. Due to the potential for unidentified resources to be altered or destroyed during excavation, drilling, and grading, the potential impacts to ecological, historical or cultural resources would be SMALL to LARGE, depending on local conditions. OPERATION—The types of land use impacts for operational activities would be similar to construction impacts regarding access restrictions because the infrastructure would be in place. Additional land disturbances would not occur from conducting operational a	

,	Summary)
	♀	
	of Environmental (
	Consequences)

	Table 10-4. Summary of Impacts for the Northwestern New Mexico Uranium Milling Region (continued)		
Topic/	GEIS		
Resource	Section	Impact Findings	
Transportation	4.5.2	CONSTRUCTION—Low magnitude traffic generated by ISL construction relative to local traffic counts would not significantly increase traffic or accidents on many of the roads in the region. Existing low traffic roads could be MODERATELY impacted by the additional worker commuting traffic during periods of peak employment. The impact would be more pronounced in areas of low traffic counts. MODERATE dust, noise, and incidental wildlife or livestock kill impacts would be possible on, or near, site access roads (dust in particular for unpaved access roads)—SMALL to MODERATE. OPERATION—Low magnitude traffic relative to local traffic counts on most roads would not significantly increase traffic or accidents. Existing low traffic roads could be moderately impacted by commuting traffic during periods of peak employment including dust, noise, and possible incidental wildlife or livestock kill impacts on, or near, site access roads. High consequences would be possible for a severe accident involving transportation of hazardous chemicals in a populated area. However, the probability of such accidents occurring would be low, owing to the limited number of shipments, comprehensive regulatory controls, and use of best management practices. For radioactive material shipments (yellowcake product, ion exchange resins, waste materials), compliance with transportation regulations would limit radiological risk for normal operations. Consequently, there is low radiological risk associated with accident conditions. Emergency response protocols would help mitigate long-term consequences of severe accidents involving release of uranium—SMALL to MODERATE.	
		AQUIFER RESTORATION—The magnitude of transportation activities would be lower than for construction and operations, with the exception of workforce commuting, which could have moderate impacts on, or near, existing low traffic roads—SMALL to MODERATE.	
		DECOMMISSIONING—The types of transportation activities and, therefore, types of impacts would be similar to those discussed for construction and operations except the magnitude of transportation activities (e.g., number and types of waste and supply shipments, no yellowcake shipments) from decommissioning could be lower than for operations. Accident risks would be bounded by operations yellowcake transportation risk estimates—SMALL.	

Table 10-4. Summary of Impacts for the Northwestern New Mexico Uranium Milling Region (continued)		
Topic/	GEIS	
Resource	Section	Impact Findings
Geology and Soils	4.5.3	CONSTRUCTION—Disturbance to soil would occur from construction (clearing, excavation, drilling, trenching, road construction). However, such disturbances would be temporary and SMALL (approx. 15 percent of the total site area), and potential impacts would be mitigated by using best management practices. A large portion of the well fields, trenches, and access roads would be restored and reseeded after construction has been completed. Excavated soils would be stockpiled, seeded, and stored on site until needed for reclamation fill. No impacts are expected to subsurface geological strata—SMALL. OPERATION—Temporary contamination or alteration of soils would be likely from operational leaks and spills and possible from transportation, use of evaporation ponds, or land application of treated wastewater. However, detection and response techniques, monitoring of treated wastewater, and eventual survey and decommissioning of all potentially impacted soils would limit the magnitude of overall impacts to soils—SMALL. AQUIFER RESTORATION—Impacts to geology and soils from aquifer restoration activities would be similar to impacts from operations due to use of the same infrastructure and similar activities conducted (e.g., well field operation, transfer lines, waste water treatment and disposal)—SMALL. DECOMMISSIONING—Impacts to geology and soils from decommissioning would be expected to be similar to impacts from construction. Activities to clean up, re-contour, and reclaim disturbed lands during decommissioning would mitigate long-term impacts to soils—SMALL.

Summary	•
9	
П	1
≦ro	
m	
of Environmental (
=	
Consequences	,

	Table 10-4. Summary of Impacts for the Northwestern New Mexico Uranium Milling Region (continued)		
Topic/	GEIS		
Resource S	Section	Impact Findings	
Surface Waters	4.5.4.1	CONSTRUCTION—Impacts to surface waters and related habitats from construction (road crossings, filling, erosion, runoff, spills or leaks of fuels and lubricants for construction equipment) would be mitigated through proper planning, design, construction methods, and best management practices. This region experiences less runoff per given area (areal flow per square mile) than the Wyoming West Uranium Milling Region. As a result, the potential for runoff-related impacts would be less. Some impacts directly related to the construction activities would be temporary and limited to the duration of the construction period. U.S. Army Corps of Engineers permits may be required when filling and crossing wetlands. Temporary changes to spring and stream flow from grading and changes in topography and natural drainage patterns could be mitigated through best management practices, or restored after the construction phase. Incidental spills of drilling fluids into local streams would be small and temporary, due to implementation of mitigation measures. Impacts from construction of roads, parking areas, and buildings on recharge to shallow aquifers would be small, owing to the limited area of impervious surfaces proposed. Infiltration of drilling fluids into the local aquifer would also be small, temporary, and localized to a few feet around boreholes—SMALL. OPERATION—Impacts from storm water runoff or direct discharge of produced waters (brine reject from reverse osmosis, or spent eluants from an ion exchange system) to surface waters would be regulated by a state or EPA-issued National Pollutant Discharge Elimination System (NPDES) permit. Expansion of facilities or pipelines during operations would generate impacts similar to construction. Because the aquifers containing uranium ore-bodies would have a weak, if any, connection to local surface water features, such as streams and springs, the impacts from operations due to use of in-place infrastructure and similar activities conducted (e.g., well field operation, transfer	

	Table 10-4. Summary of Impacts for the Northwestern New Mexico Uranium Milling Region (continued)		
Topic/	GEIS		
Resource	Section	Impact Findings	
Water— Groundwater	4.5.4.2	CONSTRUCTION—Water use impacts would be limited by the small volumes of groundwater used for routine activities such as dust suppression, mixing cements, and drilling support over short and intermittent periods. Contamination of groundwater from construction activities would be mitigated by use of best management practices—SMALL. OPERATION—Potential impacts to shallow aquifers can occur from leaks or spills from surface facilities and equipment. Shallow aquifers are important sources of drinking water in some areas of the Northwestern New Mexico Uranium Milling Region. Potential impacts to the ore-bearing and surrounding aquifers include consumptive water use and degradation of water quality (from normal production activities, off-normal excursion events, and deep well injection disposal practices). Consumptive use impacts from withdrawal of groundwater would be small because only 1 to 3 percent of pumped groundwater is not returned to the aquifer (e.g., process bleed). The amount of water lost could be reduced substantially by currently available treatment methods (e.g., reverse osmosis, brine concentration). Effects of water withdrawal on surface water would be SMALL, as the ore zone normally occurs in a confined aquifer. Estimated drawdown effects vary depending on site conditions and water treatment technology applied. Excursions of lixiviant and mobilized chemical constituents could occur from failure of well seals or other operational conditions that cause incomplete recovery of lixiviant. Well-seal-related excursions would be detected by the groundwater monitoring system, and periodic well integrity testing, and impacts would be mitigated during operation or aquifer restoration. Other excursions could result in plumes of mobilized uranium and heavy metals extending beyond the mineralization zone. The magnitude of potential impacts from vertical excursions at ISL facilities, NRC requires licensees to take preventative measures prior to starting operations including well tests, monitoring, and development	
		AQUIFER RESTORATION—Potential impacts include consumptive use and potential deep disposal of brine slurries after reverse osmosis, if applicable. The volume of water removed from the aquifer and related impacts would be dependent on site-specific conditions and the type of water treatment technology the facility used. Groundwater Consumptive use during aquifer restoration could be greater than during ISL operation, if groundwater sweep is implemented during aquifer restoration in which pumped water is not recirculated. Potential environmental impacts associated with water consumption during aquifer restorations would be determined by (1) the restoration techniques chosen, (2) the volume of water to be used, (3) the severity and extent of the contamination, and (4) the current and future use of the production and surrounding aquifers in the vicinity of the ISL facility or at the regional scale—SMALL to MODEDERATE, depending on site-specific conditions.	
		DECOMMISSIONING—Potential impacts from decommissioning would be similar to construction (water use, spills) with an additional potential to mobilize contaminants during demolition and cleanup activities. Contamination of groundwater from decommissioning activities would be mitigated by implementation of an NRC-approved decommissioning plan and use of best management practices—SMALL.	

Table 10-4. Summary of Impacts for the Northwestern New Mexico Uranium Milling Region (continued)				
Topic/	Topic/ GEIS			
Resource	Section	Impact Findings		
Ecology— Terrestrial	4.5.5.1	CONSTRUCTION—Potential terrestrial ecology impacts would include the removal of vegetation from well fields and the milling site, the modification of existing vegetative communities, the loss of sensitive plants and habitats from clearing and grading, and the potential spread of invasive species and noxious weed populations. These impacts would be temporary because restoration and reseeding occur rapidly at the completion of construction. Introduction of invasive species or noxious weeds would be possible but could be mitigated by restoration and reseeding after construction. Shrub and tree removal would require a longer restoration period. Construction noise could affect reproductive success of sage-grouse leks by interfering with mating calls. Temporary displacement of animal species would be possible. Critical wintering habitats vital for the survival of local elk populations, are located within the region. Raptors breeding onsite may be impacted by construction activities or mining operations and may be temporarily impacted depending on the time of year construction activities occur. Wildlife habitat fragmentation, temporary displacement of animal species, and direct or indirect mortalities would be possible. Implementation of wildlife surveys and mitigation measures following established guidelines would limit impacts. The magnitude of impacts depends on whether a new facility is being licensed or an existing facility is being extended—SMALL to MODERATE, depending on site-specific habitat affected. OPERATION—Habitat could be altered by operations. Access to crucial wintering habitat and water could be limited by fencing. Migratory birds could be affected by exposure to constituents in evaporation ponds, but perimeter fencing and netting could limit impacts. Temporary contamination or alteration of soils would be likely from operational leaks and spills and possible from transportation or land application of treated wastewater. However, detection and response techniques, and eventual survey and decommissio		

Table 10-4. Summary of Impacts for the Northwestern New Mexico Uranium Milling Region (continued)				
_Topic/				
Resource	GEIS Section	Impact Findings		
Ecology— Aquatic	4.5.5.2	CONSTRUCTION—Clearing and grading activities associated with construction could result in a temporary increase in sediment load in local streams, but aquatic species would recover quickly as sediment load decreases. Clearing of riparian vegetation could affect light and temperature of water. Construction impacts to wetlands would be identified and managed through U.S. Army Corps of Engineers permits, as appropriate. Construction impacts to surface waters and aquatic species would be temporary and mitigated by best management practices—SMALL. OPERATION—Impacts could result from spills or releases into surface water. Impacts would be minimized by spill prevention, identification and response programs, and National Pollutant Discharge Elimination System (NPDES) permit requirements—SMALL. AQUIFER RESTORATION—Activities would use existing (in-place) infrastructure, and impacts could result from spills or releases of untreated groundwater. Impacts would be minimized by spill prevention, identification, and response programs, and NPDES permit requirements—SMALL. DECOMMISSIONING—Decommissioning and reclamation activities could result in temporary increases in sediment load in local streams, but aquatic species would recover quickly as sediment load decreases. With completion of decommissioning, revegetation, and re-contouring, habitat would be reestablished and impacts would, therefore, be limited—SMALL.		

Table 10-4. Summary of Impacts for the Northwestern New Mexico Uranium Milling Region (continued)					
Topic/					
Resource	Section	Impact Findings			
Air Quality	4.5.6	CONSTRUCTION—Fugitive dust and combustion (vehicle and diesel) emissions during land-disturbing activities associated with construction would be small, short-term, and reduced through use of best management practices (e.g., dust suppression). For example, estimated fugitive dust emissions during ISL construction are less than 2 percent of the National Ambient Air Quality Standards (NAAQS) for PM25 and less than 1 percent for PM10. For NAAQS attainment areas such as the Northwestern New Mexico Unanium Milling Region, nonradiological air quality impacts would be SMALL. There are no Prevention of Significant Deterioration (PSD) Class I areas in the Northwestern New Mexico Uranium Milling Region. Furthermore, if impacts were initially assessed at a higher significance level, permit requirements would impose conditions or mitigation measures to reduce impacts—SMALL. OPERATION—Radiological impacts can result from dust releases from drying of lixiviant pipeline spills, radon releases from well system relief valves, resin transfer, or elution, and gaseous/particulate emissions from yellowcake dryers. Only small amounts of low dose materials would be released based on operational controls and rapid response to spills. Required spill prevention, control, and response procedures would be used to minimize impacts from operations and ventilation reduces radon buildup during operations. Compliance with the NRC-required radiation monitoring program ensures releases are within regulatory limits. Other potential nonradiological emissions during operations include fugitive dust and fuel from equipment, maintenance, transport trucks, and other vehicles. For NAAQS attainment areas such as the Northwestern New Mexico Uranium Milling Region, nonradiological air quality impacts would be SMALL. There are no PSD Class I areas in the Northwestern New Mexico Uranium Milling Region. Furthermore, if impacts were initially assessed at a higher significance level, permit requirements would impose conditions or mitigation measures to reduce			

	Table 10-4. Summary of Impacts for the Northwestern New Mexico Uranium Milling Region (continued)				
Topic/	GEIS				
Resource	Section 4.5.7	Impact Findings CONSTRUCTION—Noise generated during construction would be noticeable in proximity to operating equipment, but would be temporary (typically daytime only). Administrative and engineering controls would be used to maintain noise levels in work areas below Occupational Health and Safety Administration (OSHA) regulatory limits and be mitigated by use of personal hearing protection. Traffic noise during construction (commuting workers, truck shipments to and from the facility, and construction equipment such as trucks, bulldozers, compressors) would be localized, limited to highways in the vicinity of the site, access roads within the site, and roads in well fields. Relative increases in traffic levels would be small for larger roads, but may be moderate for lightly traveled rural roads through less populated communities. Noise may adversely affect wildlife habitat and their reproductive success in the immediate vicinity of construction activities. Noise levels decrease geometrically with distance, and at distances more than 300 m [1,000 ft], noise levels return to background levels. Wildlife generally avoid construction noise areas. The uranium districts within the Northwestern New Mexico Uranium Milling Region are generally more than 300 m [1,000 ft] from the closest community—SMALL to MODERATE.			
Noise		OPERATION—Noise-generating activities in the central uranium processing facility would be indoors, minimizing offs sound levels. Well field equipment (e.g., pumps, compressors) would also be expected to be contained within structu (e.g., header houses, satellite facilities) minimizing sound levels to offsite receptors. Administrative and engineering controls would be used to maintain noise levels in work areas below OSHA regulatory limits and be mitigated by use personal hearing protection. Traffic noise from commuting workers, truck shipments to and from the facility, and facili equipment would be localized, limited to highways in the vicinity of the site, access roads within the site, and roads in fields. Relative increases in traffic levels would be SMALL for larger roads, but may be MODERATE for lightly traveler rural roads through less populated communities. Most noise would be generated indoors, and mitigated by regulatory compliance and use of best management practices. Noise from trucks and other vehicles is typically of short duration Noise usually is not discernable to offsite receptors at distances of more than 300 m [1,000 ft]. The uranium districts within the Northwestern New Mexico Uranium Milling Region are generally more than 300 m [1,000 ft] from the closes community—SMALL to MODERATE.			
		AQUIFER RESTORATION—Noise generation is expected to be less than during construction and operations. Pumps and other well field equipment contained in buildings, minimize sound levels to offsite receptors. Existing operational infrastructure would be used, and traffic levels would be less than during construction and operations however, relative increases to existing traffic levels from commuting may be more significant for lightly traveled rural roads through smaller communities. Noise usually is not discernable to offsite receptors at distances of more than 300 m [1,000 ft]. The uranium districts within the Northwestern New Mexico Uranium Milling Region are generally more than 300 m [1,000 ft] from the closest community—SMALL to MODERATE.			
		DECOMMISSIONING—Noise generated during decommissioning would be noticeable only in proximity to operating equipment, and be temporary (typically daytime only). Administrative and engineering controls would be used to maintain noise levels in work areas below OSHA regulatory limits, and be mitigated by use of personal hearing protection. Noise levels during decommissioning would be expected to be less than during construction and would diminish as less and less equipment is used and truck traffic is reduced. Noise usually is not discernable to offsite receptors at distances of more than 300 m [1,000 ft]. The uranium districts within the Northwestern New Mexico Uranium Milling Region are generally more than 300 m [1,000 ft] from the closest community—SMALL.			

Table 10-4. Summary of Impacts for the Northwestern New Mexico Uranium Milling Region (continued)						
Topic/	GEIS					
Resource	Section	Impact Findings				
Historical and Cultural	4.5.8	CONSTRUCTION—Potential impacts during ISL facility construction could include loss of, or damage and temporary restrictions on access to, historical, cultural, and archaeological resources. Prominent cultural resources in the Northwestern New Mexico Uranium Milling Region include culturally significant landscapes such as Mount Taylor. The eligibility evaluation of cultural resources for listing in the National Register of Historic Places (RNHP) under criteria in 36 CFR 60 4(a)—(d) and/or as Traditional Cultural Properties (TCPs) is conducted as part of the site-specific review and NRC licensing procedures undertaken during the National Environmental Policy Act (NEPA) review process. The evaluation of impacts to any historic properties designated as TCPs and tribal consultations regarding cultural resources and TSPs also occurs during the site-specific licensing application and review process. Consultations to determine whether significant cultural resources would be avoided or mitigated occurs during consulting with the State Historic Preservation Office, other governmental agencies, and Native American Tribes, including Tribal Historic Preservation Office, other governmental agencies, and Native American Tribes, including Tribal Historic Preservation Office, other governmental agencies, and Native American Tribes, including Tribal Historic Preservation Office or notify the appropriate federal, tribal, and state agencies with regard to appropriate mitigation measures—SMALL or MODERATE to LARGE, depending on site-specific conditions. OPERATION—Because less land disturbance occurs during the operations phase, potential impacts to historical, cultural, and archaeological resources would be less than during construction. Conditions in the NRC license requiring adherence to procedures typically require the licensee to stop work and to notify the appropriate federal, tribal, and state agencies with regard to mitigation measures—SMALL or MODERATE to LARGE, depending on site-specific conditions. DECOMMISSIONING—Becau				

Table 10-4. Summary of Impacts for the Northwestern New Mexico Uranium Milling Region (continued)				
Topic/	GEIS			
Resource	Section	Impact Findings		
Visual and Scenic	4.5.9	CONSTRUCTION—Visual impacts result from equipment (drill rig masts, cranes), dust/diesel emissions from construction equipment, and hillside and roadside cuts. Most of the Northwestern New Mexico Uranium Milling Region is classified as Visual Resource Management (VRM) Class II through IV. A number of VRM Class II areas surrounding the national monuments (El Morro and El Malpais), the Chaco Culture National Historic Park, and the sensitive areas managed within the Mount Taylor district of the Cibola National Forest would have the greatest potential for impacts to visual resources. Most of these areas, however, are located to the north, south, and east of the potential ISL facilities, at distances of 16 km [10 mi] or more. The facilities would be located in VRM Class III and IV areas. Current understanding indicates that several potential ISL facilities may be located near the Navajo Nation or near Mount Taylor in the San Mateo Mountains. The general visual and scenic impacts associated with ISL facility construction would be temporary and SMALL, but from a Native American perspective, any construction activities would likely result in adverse impacts to the landscape, particularly for facilities located in areas within view of tribal lands and areas of special significance such as Mount Taylor. Most potential visual impacts during construction would be temporary as equipment is moved and would be miligated by implementing best management practices (e.g., dust suppression). Because of the generally rolling topography of the region, most visual impacts during operations would not be visible from more than 1 km [0.6 mi]. The visual impacts during operations would be less than those associated with construction. Most of the well field surface infrastructure has a low profile, and most piping and cables would be buried. The tallest structures would include the central uranium processing facility {10 m [30 ft]} and power lines {6 m [20 ft]}. Because of the generally rolling topography of the region, most visual impact		

Table 10-4. Summary of Impacts for the Northwestern New Mexico Uranium Milling Region (continued)				
Topic/	GEIS			
Resource	Section	Impact Findings		
Socioeconomics	4.5.10	CONSTRUCTION—Potential impacts to socioeconomics would result predominantly from employment at an ISL facility and demands on the existing public and social services, tourism/recreation, housing, infrastructure (schools, utilities), and the local work force. Total peak employment would be about 200 people, including company employees and local contractors, depending on timing of construction with other stages of the ISL lifecycle. During construction of surface facilities and well fields, the general practice has been to use local contractors (drillers, construction) if available. A local multiplier of 0.7 would indicate a maximum of about 140 ancillary jobs could be created. For example, local building materials and building supplies would be used to the extent practical. Most employees would live in larger communities with access to more services. Some construction employees, however, would commute from outside the county to the ISL facility, and skilled employees (e.g., engineers, accountants, managers) would come from outside the local work force. Some of these employees would temporarily relocate to the project area and contribute to the local economy through purchasing goods and services and taxes. Because of the small relative size and temporary nature of the ISL workforce, net impacts would be—SMALL to MODERATE, depending on proximity to less populated communities such as those in Cibola County and the Town of Grants. OPERATION—Employment levels for ISL facility operations would be similar to, or less than, for construction, with total peak employment depending on timing and overlap with other stages of the ISL lifecycle. Use of local contract workers and local building materials would be ginerated by federal, state, and local taxes on the facility and the uranium produced. Because of similar employment levels, other socioeconomic impacts would be similar to construction—SMALL to MODERATE, depending on proximity to less populated communities such as those in Cibola County and the Town of Grants. DECOM		

Table 10-4. Summary of Impacts for the Northwestern New Mexico Uranium Milling Region (continued)				
Topic/ Resource	GEIS Section	Impact Findings		
Public and Occupational Health and Safety	4.5.11	CONSTRUCTION—Worker safety would be addressed by standard construction safety practices. Fugitive dust would result from construction activities and vehicle traffic but would likely be of short duration, and not result in a radiological dose. Diese emissions would not be expected to be a concern for worker or public health, because the releases would be of short duration readily dispersed into the atmosphere—SMALL. OPERATION—Potential occupational radiological impacts from normal operations would be caused primarily by exposure to radon gas from well field, ion-exchange resin transfer operations, and venting during processing activities. Workers would also be exposed to airborne uranium particulates from dryer operations and maintenance activities. Potential public exposures to radiation would occur from the same radon releases and uranium particulate releases (i.e., from facilities without vacuum dryer technology). Both worker and public radiological exposures would be addressed by NRC regulations at 10 CFR Part 20, which require licensees to implement an NRC-approved radiation protection program. (Measured and calculated doses for workers and the public are commonly a fraction of regulated limits.) Nonradiological worker safety matters would be addressed through commonly applied occupational health and safety regulations and practices. Radiological accident risks could involve processing equipment failures leading to yellowcake slurry spills, or radon gas or uranium particulate releases. Consequences of accidents to workers and the public are generally low, with the exception of a dryer explosion which could result in worker dose above NRC limits. The likelihood of such release events would be low, based on historical operating experience at NRC-licensed facilities which is partly the result of operators following commonly applied chemical safety and handling protocols—SMALL to MODERATE. AQUIFER RESTORATION—Because the activities during aquifer restoration overlap with similar operational activities (e.g.,		

Table 10-4. Summary of Impacts for the Northwestern New Mexico Uranium Milling Region (continued)				
Topic/	GEIS	, , , , , , , , , , , , , , , , , , ,		
Resource	Section	Impact Findings		
Waste Management	4.5.12	CONSTRUCTION—The relatively small scale of construction activities (Section 2.3) and incremental development of well fields at ISL facilities would generate low volumes of construction waste—SMALL. OPERATION—Operational wastes primarily result from liquid waste streams including process bleed, flushing of depleted eluant to limit impurities, resin transfer wash, filter washing, uranium precipitation process wastes (brine), and plant wash down water. State permitting actions, NRC license conditions, and NRC inspections ensure the proper practices would be used to comply with safety requirements to protect workers and the public. Waste treatment such as reverse osmosis and radon settling would help in segregating wastes and minimizing disposal volumes. Potential impacts from surface discharge and deep well injection would be limited by the applicable permitting processes. NRC regulations address constructing, operating, and monitoring for leakage from evaporation ponds used to store and reduce volumes of liquid wastes. Potential impacts from land application of treated wastewater would be addressed by NRC review of site-specific conditions prior to approval, routine monitoring, and inclusion of irrigated land areas in decommissioning surveys. Offsite waste disposal impacts would be SMALL for radioactive wastes as a result of required preoperational disposal agreements. Impacts for hazardous and municipal waste would be SMALL due to the volume of wastes generated. For remote areas with limited available disposal capacity, such wastes may need to be shipped greater distances to facilities that have capacity. However, the volume of wastes generated and magnitude of the shipments are estimated to be low—SMALL. AQUIFER RESTORATION—Waste management activities during aquifer restoration would be dependent on site-specific conditions, the potential exists for additional generation of wastewater volume and associated treatment wastes during the restoration prodice. However, this would be offset to some degree by the red		

11 LIST OF PREPARERS

11.1 U.S. Nuclear Regulatory Commission Contributors

Alan B. Bjornsen: Project Manager

M.S., Silviculture, SUNY College of Forestry, 1971

M.S., Forestry, Syracuse University, 1971

B.S., Geology, Wheaton College, 1968

Years of Experience: 36

Ron Linton: Groundwater

M.S., Geology, West Virginia University, 1992

B.S., Geology, James Madison, 1984

Years of Experience: 20

James Park: Lead Project Manager

B.S., Geology, Virginia Polytechnic Institute and State University 1986

M.S., Structural Geology and Rock Mechanics, Imperial College, University of London,

England, 1988

M.Ed., Marymount University, 1999

Years of Experience: 15

A. Christianne Ridge: Groundwater and Public Scoping Comments

Ph.D., Environmental Engineering, University of California at Berkeley, 2004

M.S., Environmental Engineering, Cornell University, 1999

B.A., Physics, Drew University, 1996

Years of Experience: 5

11.2 Southwest Research Institute[®] and Center for Nuclear Waste Regulatory Analyses (CNWRA[®]) Contributors

Hakan Basagaoglu: Analyst—Surface/Groundwater

Ph.D., Civil/Environmental Engineering, University of California-Davis, 2000

M.S., Geological Engineering, Middle East Technical University, 1993

B.S., Geological Engineering, Middle East Technical University, 1991

Years of Experience: 16

Darius Daruwalla: Analyst—Occupational Health and Safety (Nonradiological)

M.S., Chemical Engineering, State University of New York at Buffalo, 1974

B.S., Chemical Engineering, State University of New York at Buffalo, 1971

Years of Experience: 36

Philippe Dubreuilh: Analyst—Land Use

Ph.D., Geology, University of Bordeaux, France, 1982

M.S., Geology, University of Bordeaux, France, 1977

B.S., Geology, University of Bordeaux, France, 1976

Years of Experience: 26

List of Preparers

Amy Glovan: Analyst—Noise, Aesthetics

B.A., Environmental Studies, University of Kansas, 1998

Years of Experience: 10

Patrick LaPlante: Principal Investigator Analyst—Decommissioning, Transportation and Waste Management

M.S., Biostatistics and Epidemiology, Georgetown University, 1994 B.S., Environmental Studies, Western Washington University, 1988

Years of Experience: 20

David Pickett: Analyst—Geochemistry

Ph.D., Geology, California Institute of Technology, 1991 M.S., Geology, California Institute of Technology, 1984

B.A., Geology, Rice University, 1982

Years of Experience: 25

James Prikryl: Analyst—Geology

M.A., Geology, University of Texas at Austin, 1989 B.S., Geology, University of Texas at Austin, 1984

Years of Experience: 23

Ali Simpkins: Project Manager, Analyst—Occupational Health and Safety (Radiological)

M.S., Nuclear Engineering, University of Missouri-Rolla, 1991 B.S., Nuclear Engineering, University of Missouri-Rolla, 1989

Years of Experience: 18

Marla Roberts: Analyst—Database Management and Document Integration

M.S., Geology, University of Texas at San Antonio, 2007

B.A., Geology, Vanderbilt University, 2001

Years of Experience: 7

David Turner: Analyst—Noise, Aesthetics

Ph.D., Geology, University of Utah, 1990 M.S., Geology, University of Utah, 1985

B.A., Music/Geology, College of William and Mary, 1981

Years of Experience: 26

Gary Walter: Analyst—Surface/Groundwater

Ph.D., Hydrology, University of Arizona, 1985

M.A., Geology, University of Missouri, Columbia, 1974 B.A., Chinese and Sociology, University of Kansas, 1969

Years of Experience: 38

Bradley Werling: Analyst—Air Quality

M.S., Environmental Science, University of Texas at San Antonio, 2000

B.S., Chemistry, Southwest Texas State University, 1999

B.A., Engineering Physics, Westmont College, Santa Barbara, 1985

Years of Experience: 22

11.3 CNWRA Subcontractors

Sam Blanco: Analyst—Ecological Resources

M.S., Environmental Sciences, Texas A&M Corpus Christi, 2002

B.S., Biology, Texas A&M Kingsville, 1998

Years of Experience: 21

Larry Canter: Analyst—Cumulative Impacts

Ph.D., Environmental Health Engineering, University of Texas at Austin, 1967

Years of Experience: 40

Susan Courage: Analyst—Socioeconomics and Environmental Justice

M.S., Environmental Science, University of San Antonio at San Antonio, 2003

B.S., Biology, University of San Antonio at San Antonio, 1999

Years of Experience: 8

Edgar K. Huber: Analyst—Cultural and Historic Resources

Ph.D., Anthropology, Washington State University, 1993

M.A., Anthropology, University of Colorado, Denver, 1984

B.A., Anthropology, Metropolitan State College, 1978

Years of Experience: 29

Brian Strye: Analyst—Ecological Resources

M.S., Environmental Sciences, University of San Antonio at San Antonio, 2001

B.S., Biology, University of San Antonio at San Antonio, 1996

Years of Experience: 11

12 GLOSSARY

Agreement State—A state that signed an agreement with the U.S. Nuclear Regulatory Commission (NRC) under Section 274 of the Atomic Energy Act (42 U.S.C. 2021). The state subsequently issues licenses and establishes remedial action requirements under its state laws and according to an alternative to Sections 62 or 81 of the Atomic Energy Act.

Alluvial—Pertaining to or composed of alluvium, or deposited by a stream or running water.

Alluvial fan—An outspread, gently sloping mass of alluvium deposited by a stream.

Alluvium—A general term for detrital deposits made by streams on river beds, floodplains, and alluvial fans.

Anticlinal—Of or pertaining to a generally convex upward fold, whose core contains the stratigraphically older rocks.

Aquifer—Porous water-bearing formation (bed or stratum) of permeable rock, sand, or gravel capable of producing significant quantities of water.

Aquifer Exemption—The process by which protection under the Safe Drinking Water Act for an aquifer, or a portion of an aquifer, that meets the criteria for an underground source of drinking water, has been waived by the EPA based on applicable underground injection control regulations at 40 CFR 146.4. An aquifer may be exempted if it is:

- Not currently being used—and will not be used in the future—as a drinking water source. or
- It is not reasonably expected to supply a public water system due to a high total dissolved solids content

Without an aquifer exemption, certain types of energy production, mining, or waste disposal into underground sources of drinking water would be prohibited.

Aquiclude or Aquitard—Geologic units that are impermeable (aquiclude) or of low permeability (aquitard) adjacent to an aquifer. These units serve to confine groundwater (or uranium recovery solutions) within the exempted aquifer.

Arkosic—Sediments with a considerable amount of the mineral feldspar.

Artesian—Pertaining to groundwater under sufficient hydrostatic pressure to rise above the aquifer containing it.

Ash fall—A rain of airborne volcanic ash falling from an eruption cloud.

Ball mill—A rotating, horizontal cylinder with a diameter almost equal to its length supported by a frame or shaft in which ores are ground using various grinders (such as steel balls, quartz pebbles, or porcelain balls).

Bar—An elongate offshore ridge, bank, or mound of sand or gravel, built by waves and currents, especially at the mouth of a river or at a slight distance from the beach.

Barren solution—A solution in hydrometallurgical treatment that has had valuable constituents removed.

Basin—A low area in the earth's crust, of tectonic origin, in which sediments have accumulated.

Bentonite—A soft, plastic, light-colored clay formed by chemical alteration of volcanic ash.

Bleed solution—A solution drawn to adjust production or to restore groundwater by pumping more fluids from the production zone than are injected, causing fresh groundwater to flow into the production area.

Braided stream—A stream that divides into an interlacing network of branching and reuniting shallow channels separated from each other by islands or channel bars.

Brine solution—A concentrated solution containing dissolved minerals (usually greater than 100,000 mg/liter), especially chloride salts.

Byproduct material—The tailings or wastes produced by extracting or concentrating uranium or thorium from any ore processed primarily for its source material content. See the full definition at 10 CFR Part 40.4. See also **Source Material**.

Calcareous—containing calcium carbonate (CaCO₃).

Carbonaceous—A rock or sediment containing organic matter.

Cenozoic—the latest of the four eras into which geologic time is divided; it extends from the close of the Mesozoic era, about 65 million years ago, to the present. The Cenozoic era is subdivided into Tertiary and Quaternary periods.

Channel—The deepest part of a stream.

Channel-fill deposit—Sediments deposited in a stream channel, where the transporting capacity of the stream is insufficient to remove the material supplied to it.

Clastic—Pertaining to a rock or sediment composed principally of fragments derived from pre-existing rocks or minerals, and transported some distance from their places of origin.

Clay—An earthy, extremely fine-grained sediment or soft rock composed primarily of clay-size particles (e.g., particles with diameters less than 1/256 mm).

Claystone—A cemented clay.

Coastal plain—A low, broad plain that has its margin on the oceanic shore and its strata either horizontal or very gently sloping toward the water.

Colluvium—A general term applied to loose or incoherent deposits, usually at the foot of a slope or cliff and brought there chiefly by gravity.

Confining units—A general term applied to low permeability geologic units above and below an aquifer that confine groundwater to flow within the aquifer.

Conformable—Geologic layers or strata characterized by an unbroken sequence in which the layers are formed one above the other in parallel order by uninterrupted deposition.

Conglomerate—A coarse-grained clastic sedimentary rock composed of fragments larger than 2 mm in diameter.

Continental—A sedimentary deposit laid down on land or in bodies of water not directly connected with the ocean.

Conventional uranium milling—A chemical process used to extract uranium from mined uranium ore. At conventional uranium mills, the ore typically arrives via truck and is crushed and chemically leached with sulfuric acid or alkaline solutions to remove about 90 to 95 percent of the uranium. NRC regulates the milling process (after ore enters the mill), but other agencies regulate the mining processes used to extract the ore.

Cretaceous—The first period of the Mesozoic era (after the Jurassic and before the Tertiary period of the Cenozoic era), thought to have covered the span of time between 144 and 65 million years ago; also, the corresponding system or rocks.

Crystalline—A general term for igneous and metamorphic rocks as opposed to sedimentary.

Cuesta—An asymmetrical ridge, with a long gentle slope on one side conforming with the dip of the underlying strata, and a steep or cliff like face on the other side formed by the outcrop of the resistant beds.

Decantation—The process of separating sediments from liquid by settling solids below and pouring off liquids above.

Decommissioning—The process of closing down a facility followed by reducing residual radioactivity.

Detrital—Minerals occurring in sedimentary rocks, which were derived from pre-existing rocks.

Disseminated—A scattered distribution of generally fine-grained minerals throughout a rock body, in sufficient quantity to make the deposit an ore.

Dome—An uplift or anticlinal structure, circular or elliptical in outline, in which the rocks dip gently away in all directions.

Eocene—An epoch of the Tertiary period (after the Paleocene and before the Oligocene), thought to have covered the span of time between 54.8 and 33.7 million years ago; also, the corresponding worldwide series of rocks.

Effluent—A waste liquid, solid, or gas, in its natural state or partially or completely treated, that is discharged into the environment.

Elution—The process of extracting (or eluting) one material from another by washing with a solvent (eluant) to remove adsorbed material (such as uranium) from an adsorbent such as an ion exchange resin.

Ephemeral—A stream which flows briefly in direct response to precipitation in the immediate vicinity.

Erosion—The wearing-away or soil and rock by weathering, mass wasting, and the action of streams, glaciers, waves, wind, and underground water.

Escarpment—A long, more or less continuous cliff or relatively steep slope, separating two level or gently sloping surfaces, and produced by erosion or faulting.

Excursion—The unintended spread, either horizontally or vertically, of recovery solutions beyond the production zone. Monitoring wells are installed to analyze for appropriate water quality parameters and detect excursions.

Evaporation pond—A containment pond, typically lined, to hold liquid wastes and to concentrate wastewater through evaporation.

Feldspar—A group of abundant rock-forming minerals of the general formula, MAl(Al, Si)₃O₈, where M can be K, Na, Ca, Ba, Rb, Sr, or Fe. Feldspars are the most widespread of any mineral group and constitute 60 percent of the Earth's crust.

Flare—The undetected spread of recovery solutions between the well field and monitoring wells of the production zone. Flare is also a proportionality factor that estimates the amount of aquifer water outside of the pore volume that has been affected by lixiviant flow during the recovery phase. The flare is usually expressed as a horizontal and vertical component to account for differences between the horizontal and vertical hydraulic conductivity of an aquifer material.

Floodplain—That portion of a river valley, adjacent to the channel, which is built of sediments deposited during the present regimen of the stream and is covered with water when the river overflows its banks at flood stages.

Fluvial—Produced by the action of a stream or river.

Formation—A body of rock or strata that consists dominantly of a certain lithologic type or combination of types.

Gangue—The valueless rock or mineral aggregates in an ore; that part of the ore that is not economically desirable but cannot be avoided in mining.

Granite—An igneous rock formed below the earth's surface in which quartz makes up 10 to 50 percent of the rock components.

Granitic—Pertaining to or composed of granite.

Groundwater—Water beneath the surface in the saturated zone that is under atmospheric or artesian pressure.

Heap leach—A method of extracting uranium from ore using a leaching solution. Small ore pieces are placed in a heap on an impervious material (plastic, clay, asphalt) with perforated pipes under the heap. Acidic solution is then sprayed over the ore, dissolving the uranium. The solution in the pipes is collected and transferred to an ion-exchange system for concentration of the uranium.

Heavy metals—Metallic elements, including those required for plant and animal nutrition, in trace concentration, that become toxic at higher concentrations. Examples are mercury, chromium, cadmium, and lead.

Hogback ridge—A sharp-crested ridge formed by the outcropping edges of steeply inclined resistant rocks, and produced by differential erosion.

Holocene—An epoch of the Quaternary period, from the end of the Pleistocene, approximately 8 thousand years ago, to the present time; also, the corresponding series of rocks and deposits.

Horizon—An interface that indicates a particular position in a stratigraphic sequence. Technically it is a surface with no thickness, but in practice it is commonly a distinctive very thin bed.

Humic—Pertaining to or derived from the dark, more or less stable part of the organic matter in soil.

Hydrothermal—Pertaining to a mineral deposit precipitated from a hot solutions.

Igneous—A rock or mineral that solidified from a magma.

Impermeable—A rock, sediment, or soil that is incapable of transmitting fluids under pressure.

Injection—The subsurface discharge of fluids through a well.

Injection zone—A geological formation, group of formations, or part of a formation that receives fluids through a well.

In-situ leaching (ISL)—The in-place recovery of a mineral resource without removing overburden or ore. This is typically accomplished by installing a well and recovering the resource directly from the natural deposit by exposing it to the injection and recovery of a fluid that causes the leaching, dissolution, or recovery of the mineral.

Injection well—A well or a drill hole in an *in-situ* leach operation through which barren solutions enter an underground stratum or ore body by gravity or under pressure.

Interbedded—Rock material or sediments lying between or alternating with others of different character.

Interfinger—To grade or pass from one material into another through a series of interpenetrating wedge-shaped layers.

Interstitial—A mineral deposit in which the minerals fill the pores of the host rock.

Interstratified—See Interbedded.

Intertonguing—The disappearance of sedimentary bodies in laterally adjacent masses owing to splitting into may thin tongues, each of which reaches an independent pinch-out termination.

Ion exchange—A chemical process used to recover uranium from solution by the exchange of dissolved uranium ions between a lixiviant (leach solution) and a solid, either a mineral surface or, more commonly, a synthetic polymer resin.

Isotope—Any two or more forms of an element having identical or very closely related chemical properties and the same atomic number but different atomic weights or mass numbers.

Jurassic—The second period of the Mesozoic era (after the Triassic and before the Cretaceous), thought to have covered the span of time between 206 and 144 million years ago; also, the corresponding system or rocks.

Lacustrine—Pertaining to or produced by a lake or lakes.

Lagoonal—Pertaining to a channel or bay partly or completely separated from the sea by a reef or barrier island, especially the water between an offshore coral reef and the mainland.

Leach—Dissolving of soluble constituents (e.g., uranium) from a rock or ore body by the natural action of percolating water or a lixiviant (leaching solution).

Leachate—The liquid that has percolated through the soil or other medium.

Lenticular—Pertaining to a stratigraphic lens; resembling in shape the cross section of a lens.

Lithologic—The physical character of a rock, such as color, mineralogical composition, and grain size.

Lixiviant—A leachate solution composed of native groundwater and chemicals (such as sodium carbonate/bicarbonate, ammonia, or sulfuric acid) added by the ISL facility operator. In the ISL process, the lixiviant is pumped underground for the purpose of mobilizing (dissolving) uranium from a uranium ore body.

Loam—A rich, permeable soil composed of a mixture of clay, silt, sand, and organic matter.

Marine—A sedimentary deposit laid down or caused by the sea.

Mechanical integrity—The absence of significant leakage within the injection tubing, casing, or packer (known as internal mechanical integrity), or outside of the casing (known as external mechanical integrity). Mechanical integrity tests (MITs) are performed to determine the adequacy of the construction of an injection well. Periodic mechanical integrity tests (MITs) are performed to confirm that a well maintains internal and external mechanical integrity.

Mesa—A flat-topped mountain bounded on a least one side by a steep cliff.

Mesozoic—An era of geologic time, from the end to the Paleozoic to the beginning of the Cenozoic, or from about 248 to about 65 million years ago; also, the rocks formed during that era. It includes the Triassic, Jurassic, and Cretaceous periods.

Metamorphic—A rock derived from pre-existing rocks by mineralogical, chemical, and/or structural changes in response to marked changes in temperature, pressure, shearing stress, and chemical environment.

Meteoric—Pertaining to or derived from the earth's atmosphere, e.g. meteoric water.

Micaceous—Consisting of, containing, or pertaining to mica – a group of minerals of the general formula (K, Na, Ca)(Mg, Fe, Li, Al)₂₋₃(Al, Si)₄O₁₀(OH, F)₂. Micas are prominent rock-forming constituents of igneous and metamorphic rocks.

Mill feed—Uranium ore supplied to a crusher or grinding mill in an ore-dressing process.

Mill tailings—See Tailings.

Miocene—An epoch of the Tertiary period (after the Oligocene and before the Pliocene), thought to have covered the span of time between 23.8 and 5.3 million years ago; also, the corresponding worldwide series of rocks.

Mudstone—A fine-grained sedimentary rock in which the proportion of clay and silt are approximately equal.

Natural levee—A ridge or embankment of sand and silt, built up by a stream on its flood plain along both banks of its channel.

Oligocene—An epoch of the Tertiary period (after the Eocene and before the Miocene), thought to have covered the span of time between 33.7 and 23.8 million years ago; also, the corresponding worldwide series of rocks.

Ore—A naturally occurring mineral that contains an economically valuable constituent, such as uranium, in sufficient concentration and quantity to allow economic production.

Outcrop—That part of a geologic formation or structure that appears at the surface of the earth.

Overbank deposit—Silt and clay deposited from suspension on a flood plain by floodwaters that cannot be contained within the stream channel.

Oxidation—An oxidizing environment is characterized by an excess of free oxygen (either dissolved or as a gas). During oxidation, the atoms in an element lose electrons and the valence state of the element increases. Chemically, oxidation is the opposite process from reduction (see **Reduction**). Oxidized uranium with a 6+ valence state (U⁶⁺ with fewer electrons) is more readily dissolved than reduced uranium (U⁴⁺ with more electrons).

Packer—A mechanical device set immediately above the injection zone that seals the outside of the tubing to the inside of the long string casing. A packer may be a simple mechanically set rubber device or a complex concentric seal assembly.

Paleocene—An epoch of the Tertiary period (after the Cretaceous period and before the Eocene), thought to have covered the span of time between 65 and 54.8 million years ago; also, the corresponding worldwide series of rocks.

Paleosol—A buried soil; a soil of the past.

Paleozoic—An era of geologic time, from the end of the Precambrian to the beginning of the Mesozoic, or from about 543 to about 248 million years ago. Also, the rocks formed during that era.

Paludal—Pertaining to a marsh.

Pennsylvanian—A period of the Paleozoic era (before the Permian), thought to have covered the span of time between 323 and 290 million years ago; also, the corresponding system or rocks.

Permeability—The ease with which fluid flows through a porous rock or sediment. Rock or sediment that allows water to move through at an appreciable rate are called "permeable."

Permian—The last period of the Paleozoic era, thought to have covered the span of time between 290 and 248 million years ago; also, the corresponding system of rocks.

Physiographic province—A region of which all parts are similar in geologic structure and climate and which has had a unified geologic history.

Plateau—A relatively elevated area of comparatively flat land which is commonly limited on a least one side by an abrupt descent to lower ground.

Pleistocene—An epoch of the Quaternary period, after the Pliocene of the Tertiary and before the Holocene; also, the corresponding worldwide series of rocks. It began about 1.8 million years ago and lasted until the start of the Holocene some 8,000 years ago.

Pliocene—An epoch of the Tertiary period (after the Miocene and before the Pleistocene), thought of have covered the span of time between 5.3 and 1.8 million years ago; also, the corresponding worldwide series of rocks.

Pore space or porosity—The collective open spaces of a rock. It is a measure of the amount of liquid or gas that may be absorbed or produced by a particular formation.

Pore volume—A volume equal to the open space in rock or soil. The ISL industry uses this term to define an indirect measurement of a unit volume of aquifer water affected by ISL recovery. It represents the volume of water that fills the void space inside a certain volume of rock or sediment. Pore volume provides a unit reference that an operator can use to describe (1) the amount of lixiviant circulation needed to leach an ore body or (2) the unit number of treated water circulations needed to flow through a depleted ore body to achieve restoration. A pore volume allows an operator to use relatively small-scale studies and scale the results to field-level pilot tests or to commercial well field scales. Typically, a pore volume is calculated by multiplying the surficial area of a well field (the area covered by injection and recovery wells) by the thickness of the production zone being exploited and the estimated or measured porosity of the aquifer material.

Potentiometric surface—An imaginary surface representing the total head of groundwater and defined by the level to which water will rise in a well.

Precambrian—All geologic time, and its corresponding rocks, before the beginning of the Paleozoic.

Pregnant solution—A solution containing a dissolved, extractable mineral that was leached from the ore; uranium leach solution pumped up from the underground ore zone through a production hole. Also called "pregnant lixiviant."

Primacy or primary enforcement authority—The authority delegated by EPA to implement the UIC Program. To receive primacy, a state, territory, or tribe must demonstrate to EPA that its UIC program is at least as stringent as the federal standards; the state, territory, or tribal UIC requirements may be more stringent than the federal requirements. (For Class II, states must demonstrate that their programs are effective in preventing pollution of USDWs.) EPA may grant primacy for all or part of the UIC program, e.g., for certain classes of injection wells.

Production zone—The uranium-bearing portion of a geological formation or part of a formation that is the target of ISL uranium recovery by underground injection and production of lixiviant.

Pyrite—The most widespread and abundant of the sulfide minerals, H₂S.

Quaternary—The second period of the Cenozoic era, following the Tertiary; also, the corresponding system or rocks. It began about 1.8 million years ago and extends to the present. It consists of two epochs: the Pleistocene and the Holocene.

Quartz—Crystalline silica, an important rock-forming mineral, SiO₂.

Quartzose—Containing quartz as a principal constituent.

Production bleed—See Bleed Solution.

Production (or recovery) well—A well or a drill hole in an *in-situ* leach operation through which pregnant (uranium-bearing) solutions are extracted from an underground stratum or uranium deposit.

Radioisotope—An unstable isotope of an element that decays or disintegrates spontaneously, emitting radiation. Approximately 5,000 natural and artificial radioisotopes have been identified.

Radon—A chemically inert radioactive gaseous element formed when radium decays. Exposure to radon may pose a potential health hazard.

Reclamation—The process of restoring the surface environment to acceptable pre-existing conditions. Reclamation includes activities such as surface contouring, equipment removal, well plugging, and revegetation.

Redox—A term commonly used to refer to the oxidation-reduction potential of a chemical system.

Reduction—A reducing environment is characterized by little or no free oxygen (dissolved or as a gas). During reduction, the atoms in an element gain electrons and the valence state of the element decreases. Chemically, reduction is the opposite process from oxidation (see **Oxidation**). Reduced uranium (U⁴⁺ with more electrons) is less dissolvable than oxidized uranium (U⁶⁺ with fewer electrons).

Remote ion exchange (RIX)—A type of ISL uranium recovery operation where pregnant lixiviant from production wells is collected at a small satellite RIX facility. The uranium is stripped from the lixiviant by loading onto ion exchange resins. The loaded resins are then transported by tanker truck to a larger central facility for additional processing and uranium

recovery. RIX operations are used to produce uranium from smaller, more disperse uranium deposits.

Restoration—Returning each constituent in the affected groundwater to its NRC-approved baseline concentration or to an alternate standard approved by NRC.

Reverse osmosis—The act of reversing a diffusion through a semipermeable membrane, typically separating a solvent and a solution, that tends to equalize their concentrations. In ISL facilities, this process is used to treat wastewater to remove dissolved constituents and reduce total dissolved solids.

Rip rap—Cobblestone or coarsely broken rock used for protection against erosion of embankments or gullies.

Roll front—A localized uranium deposit in the form of a roll or interface that separates an oxidized interior from a reduced exterior. The reduced side of this interface is significantly enriched in uranium.

Runoff—The portion of rainfall that is not absorbed by soil, evaporated, or transpired by plants, but finds its way directly into streams or as overland surface flows.

Sand—A loose aggregate of particles having a diameter in the range of 1/16 to 2 mm.

Sandstone—A clastic sedimentary rock composed of grains of sand size set in a matrix of silt or clay and more or less firmly united by a cementing material.

Satellite facility—A remotely located facility for initial processing of uranium bearing solutions [see Remote ion exchange (RIX)].

Scour protection—Using flushing water to protect the trench surface from erosion.

Sediment—Solid fragmental material transported and deposited by wind or water, or chemically precipitated from solution, that forms in layers in loose unconsolidated form.

Sedimentary—Pertaining to or containing sediment, or formed by its deposition.

Shale—A fine-grained detrital sedimentary rock, formed by the compaction of clay, silt, and mud.

Silicified—A rock in which silica, in the form of quartz, chalcedony, or opal, has replaced existing minerals.

Silt—A loose aggregate of rock or mineral particles commonly in the range of 1/16 to 1/256 mm.

Siltstone—A massive mudstone in which silt predominates over clay.

Source material—Uranium or thorium, or any combination thereof, in any physical or chemical form or ores which contain by weight one-twentieth of one percent (0.05%) or more of: uranium, thorium, or any combination thereof. Source material does not include special nuclear material.

Spit—A small point of sand or gravel projecting from the shore into a body of water; a fingerlike extension of the beach.

Stratabound—A type of mineral deposit contained within a single layer of sedimentary rock. Usually refers to a deposit in a permeable rock such as a sandstone bounded by impermeable confining layers such as shelves.

Stratigraphic unit—A body of strata recognized as a unit for description, mapping, and correlation.

Stratigraphic section or sequence—A chronologic succession of sedimentary rocks from older below to younger above, essentially without interruption.

Subsidence—Sinking or downward settling of the earth's surface.

Surety—A type of bond to ensure that funds are available for a specific activity (in this case, dismantling, reclamation, restoration, and remediation of uranium production sites). If the company goes bankrupt, the bonding company pays NRC or the appropriate state the amount of the bond. NRC or the appropriate state must ensure that the amount is adequate for the remediation activities.

Synclinal—Pertaining to a fold of which the core contains the stratigraphically younger rocks; it is generally concave upward.

Tailings—The remaining portion of a metal-bearing ore consisting of finely ground rock and process liquid after some or all of the metal, such as uranium, has been extracted.

Terrace—A relatively level bench or steplike surface breaking the continuity of a slope.

Tertiary—The first period of the Cenozoic era (after the Cretaceous of the Mesozoic era and before the Quaternary), thought to have covered the span of time between 65 million and 1.8 million years ago; also, the corresponding system of rocks. It is divided into five epochs: the Paleocene, Eocene, Oligocene, Miocene, and Pliocene.

Texture—The physical nature of a soil, according to the relative proportions of sand, silt, and clay.

Tiering—For the purposes of the National Environmental Policy Act, tiering is defined by the Council on Environmental Quality in 40 CFR 1508.28. It refers to "the coverage of general matters in broader environmental impact statements (such as national program or policy statements) with subsequent narrower statements or environmental analyses (such as regional or basinwide program statements or ultimately site-specific statements) incorporating by reference the general discussions and concentrating solely on the issues specific to the statement subsequently prepared."

Topography—The general configuration of a land surface including elevations.

Tongue—A minor stratigraphic unit of limited extent, especially a member that extends outward beyond the main body of a formation and disappears laterally.

Transgression—The spread of the sea over land areas.

Triassic—The first period of the Mesozoic era (after the Permian of the Paleozoic era, and before the Jurassic), thought to have covered the span of time between 248 and 206 million years ago; also, the corresponding system of rocks.

Trunkline—Main pipeline that brings together flow from individual wells.

Tuff—A general term for consolidated rocks formed by volcanic explosion or aerial expulsion from a volcanic vent.

Tuffaceous—Rocks or sediments containing particles derived from pre-existing tuff rocks.

Underground injection control (UIC)—The UIC Program is administered by the EPA or by tribal or state agencies that have been granted primacy by EPA. The UIC program is responsible for regulating the construction, operation, permitting, and closure of injection wells that place fluids underground for storage or disposal. Based on EPA regulations, UIC programs identify five different classes of injection wells.

Class I wells—Technologically sophisticated wells that inject wastes into deep, isolated rock formations below the lowermost USDW. Class I wells may inject hazardous waste, non-hazardous industrial waste, or municipal wastewater.

Class II wells—Wells that inject brines and other fluids associated with oil and gas production, or storage of hydrocarbons. Class II well types include salt water disposal wells, enhanced recovery wells, and hydrocarbon storage wells.

Class III wells—Wells that inject fluids associated with solution mining of minerals. Mining practices that use Class III wells include salt solution mining, in-situ leaching of uranium, and sulfur mining using the Frasch process.

Class IV wells—Wells that inject hazardous or radioactive wastes into or above a USDW. These wells are banned unless authorized under a federal or state groundwater remediation project.

Class V wells—Wells not included in Classes I to IV. Class V wells inject non-hazardous fluids into or above a USDW and are typically shallow, on-site disposal systems; however, this class also includes some deeper injection operations. There are approximately 20 subtypes of Class V wells.

Underground Source of Drinking Water (USDW)—An aquifer or portion of an aquifer that supplies any public water system or that contains a sufficient quantity of ground water to supply a public water system, and currently supplies drinking water for human consumption, or that contains fewer than 10,000 mg/l total dissolved solids and is not an exempted aquifer.

Uplift—A structurally high area in the crust, produced by movements that raise the rocks, as in a broad dome or arch.

Uraniferous—A rock or sediment that contains uranium.

Viewshed—The Bureau of Land Management uses this term in the Visual Resource Management process to describe landscape that can be seen under favorable atmospheric conditions from a viewpoint (key observation point) or along a transportation corridor.

Visual resources—The visible physical features of a landscape (topography, water, vegetation, animals, structures, and other features) that constitute the scenery of an area.

Visual resource management (VRM) classes—

Class I—The objective of this class is to maintain a landscape setting that appears unaltered by humans. It is applied to wilderness areas, some natural areas, wild portions of wild and scenic rivers, and other similar situations in which management activities are to be restricted.

Class II—The objective of this class is to design proposed alterations so as to retain the existing character of the landscape. The level of change to the characteristic landscape should be low. Management activities may be seen, but should not attract the attention of the casual observer. Any changes must repeat the basic elements of form, line, color, and texture found in the predominant natural features of the characteristic landscape.

Class III—The objective of this class is to design proposed alterations so as to partially retain the existing character of the landscape. Contrasts to the basic elements (form, line, color, and texture) caused by a management activity may be evident and begin to attract attention in the characteristic landscape; however, the changes should remain subordinate to the existing characteristic landscape.

Class IV—The objective of this class is to provide for management activities that require major modification of the existing character of the landscape. Contrasts may attract attention and be a dominant feature of the landscape in terms of scale; however, changes should repeat the basic elements (form, line, color, and texture) inherent in the characteristic landscape.

Class V or Rehabilitation Area—Change is needed or change may add acceptable visual variety to an area. This class applies to areas where the naturalistic character has been disturbed to a point at which rehabilitation is needed to make it conform to the surrounding landscape. This class would apply to areas where the quality class has been reduced because of unacceptable cultural modification as identified in the scenic evaluation. The contrast is inharmonious with the characteristic landscape. It may also be applied to areas that have the potential for enhancement, where it would add acceptable visual variety to an area or site. It should be considered an interim or short-term classification until one of the other VRM class objectives can be reached through rehabilitation or enhancement. The desired VRM class should be identified.

Volcanic—Pertaining to the activities, structures, or rock types of a volcano.

Volcanic ash—Fine (under 2 mm in diameter) clastic rock material formed by volcanic explosion or aerial expulsion from a volcanic vent.

Volcaniclastic—Pertaining to a clastic rock containing volcanic material.

Well field—The area of an ISL operation that encompasses the array of injection, recovery (or production), and monitoring wells and interconnected piping employed in the leaching process.

Yellowcake—The product of the uranium extraction (milling) process that is mixture of uranium oxides that can vary in proportion and in color from yellow to orange to dark green (blackish) depending at which temperature the material was dried.